

Algebraic categorification and its applications, II

Volodymyr Mazorchuk
(Uppsala University)

Winter School “Geometry and physics”
January 17-24, 2015, Srní, Czech Republic

Complex representations of symmetric groups

S_n — the symmetric group on $\{1, 2, \dots, n\}$

$P_n := \{\lambda = (\lambda_1, \dots, \lambda_k) : \lambda_1 \geq \dots \geq \lambda_k, \lambda_1 + \dots + \lambda_k = n\}$

$\lambda \in P_n$ is called a partition of n , denoted $\lambda \vdash n$

\mathcal{S}^λ — the Specht module associated to λ

Theorem. $\{\mathcal{S}^\lambda : \lambda \vdash n\}$ is a cross-section of isomorphism classes of simple S_n -modules.

Examples:

- ▶ $\mathcal{S}^{(n)}$ is the trivial module
- ▶ $\mathcal{S}^{(1,1,\dots,1)}$ is the sign module
- ▶ $\mathcal{S}^{(n)} \oplus \mathcal{S}^{(n-1,1)}$ is the natural module

Complex representations of symmetric groups

S_n — the **symmetric group** on $\{1, 2, \dots, n\}$

$\mathbf{P}_n := \{\lambda = (\lambda_1, \dots, \lambda_k) : \lambda_1 \geq \dots \geq \lambda_k, \lambda_1 + \dots + \lambda_k = n\}$

$\lambda \in \mathbf{P}_n$ is called a **partition** of n , denoted $\lambda \vdash n$

\mathcal{S}^λ — the **Specht module** associated to λ

Theorem. $\{\mathcal{S}^\lambda : \lambda \vdash n\}$ is a cross-section of isomorphism classes of simple S_n -modules.

Examples:

- ▶ $\mathcal{S}^{(n)}$ is the trivial module
- ▶ $\mathcal{S}^{(1,1,\dots,1)}$ is the sign module
- ▶ $\mathcal{S}^{(n)} \oplus \mathcal{S}^{(n-1,1)}$ is the natural module

Complex representations of symmetric groups

S_n — the **symmetric group** on $\{1, 2, \dots, n\}$

$\mathbf{P}_n := \{\lambda = (\lambda_1, \dots, \lambda_k) : \lambda_1 \geq \dots \geq \lambda_k, \lambda_1 + \dots + \lambda_k = n\}$

$\lambda \in \mathbf{P}_n$ is called a **partition** of n , denoted $\lambda \vdash n$

\mathcal{S}^λ — the **Specht module** associated to λ

Theorem. $\{\mathcal{S}^\lambda : \lambda \vdash n\}$ is a cross-section of isomorphism classes of simple S_n -modules.

Examples:

- ▶ $\mathcal{S}^{(n)}$ is the trivial module
- ▶ $\mathcal{S}^{(1,1,\dots,1)}$ is the sign module
- ▶ $\mathcal{S}^{(n)} \oplus \mathcal{S}^{(n-1,1)}$ is the natural module

Complex representations of symmetric groups

S_n — the **symmetric group** on $\{1, 2, \dots, n\}$

$\mathbf{P}_n := \{\lambda = (\lambda_1, \dots, \lambda_k) : \lambda_1 \geq \dots \geq \lambda_k, \lambda_1 + \dots + \lambda_k = n\}$

$\lambda \in \mathbf{P}_n$ is called a **partition** of n , denoted $\lambda \vdash n$

\mathcal{S}^λ — the **Specht module** associated to λ

Theorem. $\{\mathcal{S}^\lambda : \lambda \vdash n\}$ is a **cross-section** of isomorphism classes of simple S_n -modules.

Examples:

- ▶ $\mathcal{S}^{(n)}$ is the trivial module
- ▶ $\mathcal{S}^{(1,1,\dots,1)}$ is the sign module
- ▶ $\mathcal{S}^{(n)} \oplus \mathcal{S}^{(n-1,1)}$ is the natural module

Complex representations of symmetric groups

S_n — the **symmetric group** on $\{1, 2, \dots, n\}$

$\mathbf{P}_n := \{\lambda = (\lambda_1, \dots, \lambda_k) : \lambda_1 \geq \dots \geq \lambda_k, \lambda_1 + \dots + \lambda_k = n\}$

$\lambda \in \mathbf{P}_n$ is called a **partition** of n , denoted $\lambda \vdash n$

\mathcal{S}^λ — the **Specht module** associated to λ

Theorem. $\{\mathcal{S}^\lambda : \lambda \vdash n\}$ is a **cross-section** of isomorphism classes of simple S_n -modules.

Examples:

- ▶ $\mathcal{S}^{(n)}$ is the trivial module
- ▶ $\mathcal{S}^{(1,1,\dots,1)}$ is the sign module
- ▶ $\mathcal{S}^{(n)} \oplus \mathcal{S}^{(n-1,1)}$ is the natural module

Complex representations of symmetric groups

S_n — the **symmetric group** on $\{1, 2, \dots, n\}$

$\mathbf{P}_n := \{\lambda = (\lambda_1, \dots, \lambda_k) : \lambda_1 \geq \dots \geq \lambda_k, \lambda_1 + \dots + \lambda_k = n\}$

$\lambda \in \mathbf{P}_n$ is called a **partition** of n , denoted $\lambda \vdash n$

\mathcal{S}^λ — the **Specht module** associated to λ

Theorem. $\{\mathcal{S}^\lambda : \lambda \vdash n\}$ is a **cross-section** of isomorphism classes of simple S_n -modules.

Examples:

- ▶ $\mathcal{S}^{(n)}$ is the **trivial module**
- ▶ $\mathcal{S}^{(1,1,\dots,1)}$ is the **sign module**
- ▶ $\mathcal{S}^{(n)} \oplus \mathcal{S}^{(n-1,1)}$ is the **natural module**

Complex representations of symmetric groups

S_n — the **symmetric group** on $\{1, 2, \dots, n\}$

$\mathbf{P}_n := \{\lambda = (\lambda_1, \dots, \lambda_k) : \lambda_1 \geq \dots \geq \lambda_k, \lambda_1 + \dots + \lambda_k = n\}$

$\lambda \in \mathbf{P}_n$ is called a **partition** of n , denoted $\lambda \vdash n$

\mathcal{S}^λ — the **Specht module** associated to λ

Theorem. $\{\mathcal{S}^\lambda : \lambda \vdash n\}$ is a **cross-section** of isomorphism classes of simple S_n -modules.

Examples:

- ▶ $\mathcal{S}^{(n)}$ is the **trivial module**
- ▶ $\mathcal{S}^{(1,1,\dots,1)}$ is the **sign module**
- ▶ $\mathcal{S}^{(n)} \oplus \mathcal{S}^{(n-1,1)}$ is the **natural module**

Complex representations of symmetric groups

S_n — the **symmetric group** on $\{1, 2, \dots, n\}$

$\mathbf{P}_n := \{\lambda = (\lambda_1, \dots, \lambda_k) : \lambda_1 \geq \dots \geq \lambda_k, \lambda_1 + \dots + \lambda_k = n\}$

$\lambda \in \mathbf{P}_n$ is called a **partition** of n , denoted $\lambda \vdash n$

\mathcal{S}^λ — the **Specht module** associated to λ

Theorem. $\{\mathcal{S}^\lambda : \lambda \vdash n\}$ is a **cross-section** of isomorphism classes of simple S_n -modules.

Examples:

- ▶ $\mathcal{S}^{(n)}$ is the **trivial** module
- ▶ $\mathcal{S}^{(1,1,\dots,1)}$ is the **sign** module
- ▶ $\mathcal{S}^{(n)} \oplus \mathcal{S}^{(n-1,1)}$ is the **natural** module

Complex representations of symmetric groups

S_n — the **symmetric group** on $\{1, 2, \dots, n\}$

$\mathbf{P}_n := \{\lambda = (\lambda_1, \dots, \lambda_k) : \lambda_1 \geq \dots \geq \lambda_k, \lambda_1 + \dots + \lambda_k = n\}$

$\lambda \in \mathbf{P}_n$ is called a **partition** of n , denoted $\lambda \vdash n$

\mathcal{S}^λ — the **Specht module** associated to λ

Theorem. $\{\mathcal{S}^\lambda : \lambda \vdash n\}$ is a **cross-section** of isomorphism classes of simple S_n -modules.

Examples:

- ▶ $\mathcal{S}^{(n)}$ is the **trivial** module
- ▶ $\mathcal{S}^{(1,1,\dots,1)}$ is the **sign** module
- ▶ $\mathcal{S}^{(n)} \oplus \mathcal{S}^{(n-1,1)}$ is the **natural** module

Complex representations of symmetric groups

S_n — the **symmetric group** on $\{1, 2, \dots, n\}$

$\mathbf{P}_n := \{\lambda = (\lambda_1, \dots, \lambda_k) : \lambda_1 \geq \dots \geq \lambda_k, \lambda_1 + \dots + \lambda_k = n\}$

$\lambda \in \mathbf{P}_n$ is called a **partition** of n , denoted $\lambda \vdash n$

\mathcal{S}^λ — the **Specht module** associated to λ

Theorem. $\{\mathcal{S}^\lambda : \lambda \vdash n\}$ is a **cross-section** of isomorphism classes of simple S_n -modules.

Examples:

- ▶ $\mathcal{S}^{(n)}$ is the **trivial** module
- ▶ $\mathcal{S}^{(1,1,\dots,1)}$ is the **sign** module
- ▶ $\mathcal{S}^{(n)} \oplus \mathcal{S}^{(n-1,1)}$ is the **natural** module

Complex representations of symmetric groups

S_n — the **symmetric group** on $\{1, 2, \dots, n\}$

$\mathbf{P}_n := \{\lambda = (\lambda_1, \dots, \lambda_k) : \lambda_1 \geq \dots \geq \lambda_k, \lambda_1 + \dots + \lambda_k = n\}$

$\lambda \in \mathbf{P}_n$ is called a **partition** of n , denoted $\lambda \vdash n$

\mathcal{S}^λ — the **Specht module** associated to λ

Theorem. $\{\mathcal{S}^\lambda : \lambda \vdash n\}$ is a **cross-section** of isomorphism classes of simple S_n -modules.

Examples:

- ▶ $\mathcal{S}^{(n)}$ is the **trivial** module
- ▶ $\mathcal{S}^{(1,1,\dots,1)}$ is the **sign** module
- ▶ $\mathcal{S}^{(n)} \oplus \mathcal{S}^{(n-1,1)}$ is the **natural** module

Principal block of \mathcal{O} for \mathfrak{sl}_n

$\mathfrak{g} := \mathfrak{sl}_n(\mathbb{C})$

\mathcal{O} — BGG category \mathcal{O}

\mathcal{O}_0 — principal block of \mathcal{O}

S_n — Weyl group of \mathfrak{g}

$M(\mu)$ — Verma module with highest weight μ

$L(\mu)$ — unique simple quotient of $M(\mu)$

Theorem. $\{L(w) := L(w \cdot 0) : w \in S_n\}$ is a cross-section of isomorphism classes of simple objects in \mathcal{O}_0

Principal block of \mathcal{O} for \mathfrak{sl}_n

$$\mathfrak{g} := \mathfrak{sl}_n(\mathbb{C})$$

\mathcal{O} — BGG category \mathcal{O}

\mathcal{O}_0 — principal block of \mathcal{O}

S_n — Weyl group of \mathfrak{g}

$M(\mu)$ — Verma module with highest weight μ

$L(\mu)$ — unique simple quotient of $M(\mu)$

Theorem. $\{L(w) := L(w \cdot 0) : w \in S_n\}$ is a cross-section of isomorphism classes of simple objects in \mathcal{O}_0

Principal block of \mathcal{O} for \mathfrak{sl}_n

$$\mathfrak{g} := \mathfrak{sl}_n(\mathbb{C})$$

\mathcal{O} — BGG category \mathcal{O}

\mathcal{O}_0 — principal block of \mathcal{O}

S_n — Weyl group of \mathfrak{g}

$M(\mu)$ — Verma module with highest weight μ

$L(\mu)$ — unique simple quotient of $M(\mu)$

Theorem. $\{L(w) := L(w \cdot 0) : w \in S_n\}$ is a cross-section of isomorphism classes of simple objects in \mathcal{O}_0

Principal block of \mathcal{O} for \mathfrak{sl}_n

$$\mathfrak{g} := \mathfrak{sl}_n(\mathbb{C})$$

\mathcal{O} — BGG category \mathcal{O}

\mathcal{O}_0 — principal block of \mathcal{O}

S_n — Weyl group of \mathfrak{g}

$M(\mu)$ — Verma module with highest weight μ

$L(\mu)$ — unique simple quotient of $M(\mu)$

Theorem. $\{L(w) := L(w \cdot 0) : w \in S_n\}$ is a cross-section of isomorphism classes of simple objects in \mathcal{O}_0

Principal block of \mathcal{O} for \mathfrak{sl}_n

$\mathfrak{g} := \mathfrak{sl}_n(\mathbb{C})$

\mathcal{O} — BGG category \mathcal{O}

\mathcal{O}_0 — principal block of \mathcal{O}

S_n — Weyl group of \mathfrak{g}

$M(\mu)$ — Verma module with highest weight μ

$L(\mu)$ — unique simple quotient of $M(\mu)$

Theorem. $\{L(w) := L(w \cdot 0) : w \in S_n\}$ is a cross-section of isomorphism classes of simple objects in \mathcal{O}_0

Principal block of \mathcal{O} for \mathfrak{sl}_n

$$\mathfrak{g} := \mathfrak{sl}_n(\mathbb{C})$$

\mathcal{O} — BGG category \mathcal{O}

\mathcal{O}_0 — principal block of \mathcal{O}

S_n — Weyl group of \mathfrak{g}

$M(\mu)$ — Verma module with highest weight μ

$L(\mu)$ — unique simple quotient of $M(\mu)$

Theorem. $\{L(w) := L(w \cdot 0) : w \in S_n\}$ is a cross-section of isomorphism classes of simple objects in \mathcal{O}_0

Principal block of \mathcal{O} for \mathfrak{sl}_n

$$\mathfrak{g} := \mathfrak{sl}_n(\mathbb{C})$$

\mathcal{O} — BGG category \mathcal{O}

\mathcal{O}_0 — principal block of \mathcal{O}

S_n — Weyl group of \mathfrak{g}

$M(\mu)$ — Verma module with highest weight μ

$L(\mu)$ — unique simple quotient of $M(\mu)$

Theorem. $\{L(w) := L(w \cdot 0) : w \in S_n\}$ is a cross-section of isomorphism classes of simple objects in \mathcal{O}_0

Principal block of \mathcal{O} for \mathfrak{sl}_n

$$\mathfrak{g} := \mathfrak{sl}_n(\mathbb{C})$$

\mathcal{O} — BGG category \mathcal{O}

\mathcal{O}_0 — principal block of \mathcal{O}

S_n — Weyl group of \mathfrak{g}

$M(\mu)$ — Verma module with highest weight μ

$L(\mu)$ — unique simple quotient of $M(\mu)$

Theorem. $\{L(w) := L(w \cdot 0) : w \in S_n\}$ is a cross-section of isomorphism classes of simple objects in \mathcal{O}_0

Principal block of \mathcal{O} for \mathfrak{sl}_n

$$\mathfrak{g} := \mathfrak{sl}_n(\mathbb{C})$$

\mathcal{O} — BGG category \mathcal{O}

\mathcal{O}_0 — principal block of \mathcal{O}

S_n — Weyl group of \mathfrak{g}

$M(\mu)$ — Verma module with highest weight μ

$L(\mu)$ — unique simple quotient of $M(\mu)$

Theorem. $\{L(w) := L(w \cdot 0) : w \in S_n\}$ is a cross-section of isomorphism classes of simple objects in \mathcal{O}_0

Grothendieck group and different bases

Corollary. $\text{Gr}(\mathcal{O}_0) \cong \mathbb{Z}[S_n]$.

Note. $\{[L(w)] : w \in S_n\}$ is the natural basis in $\text{Gr}(\mathcal{O}_0)$.

$$\Delta(w) := M(w \cdot 0)$$

Fact. $\{[\Delta(w)] : w \in S_n\}$ is the standard basis in $\text{Gr}(\mathcal{O}_0)$.

Reason: $[\Delta(x) : L(y)] \neq 0$ implies $x \leq y$ and $[\Delta(x) : L(x)] = 1$.

Fact. \mathcal{O}_0 has finite global dimension.

$P(w)$ — the indecomposable projective cover of $L(w)$

Corollary. $\{[P(w)] : w \in S_n\}$ is a basis in $\text{Gr}(\mathcal{O}_0)$.

Grothendieck group and different bases

Corollary. $\text{Gr}(\mathcal{O}_0) \cong \mathbb{Z}[S_n]$.

Note. $\{[L(w)] : w \in S_n\}$ is the natural basis in $\text{Gr}(\mathcal{O}_0)$.

$$\Delta(w) := M(w \cdot 0)$$

Fact. $\{[\Delta(w)] : w \in S_n\}$ is the standard basis in $\text{Gr}(\mathcal{O}_0)$.

Reason: $[\Delta(x) : L(y)] \neq 0$ implies $x \leq y$ and $[\Delta(x) : L(x)] = 1$.

Fact. \mathcal{O}_0 has finite global dimension.

$P(w)$ — the indecomposable projective cover of $L(w)$

Corollary. $\{[P(w)] : w \in S_n\}$ is a basis in $\text{Gr}(\mathcal{O}_0)$.

Grothendieck group and different bases

Corollary. $\mathrm{Gr}(\mathcal{O}_0) \cong \mathbb{Z}[S_n]$.

Note. $\{[L(w)] : w \in S_n\}$ is the **natural** basis in $\mathrm{Gr}(\mathcal{O}_0)$.

$$\Delta(w) := M(w \cdot 0)$$

Fact. $\{[\Delta(w)] : w \in S_n\}$ is the standard basis in $\mathrm{Gr}(\mathcal{O}_0)$.

Reason: $[\Delta(x) : L(y)] \neq 0$ implies $x \leq y$ and $[\Delta(x) : L(x)] = 1$.

Fact. \mathcal{O}_0 has finite global dimension.

$P(w)$ — the indecomposable projective cover of $L(w)$

Corollary. $\{[P(w)] : w \in S_n\}$ is a basis in $\mathrm{Gr}(\mathcal{O}_0)$.

Grothendieck group and different bases

Corollary. $\text{Gr}(\mathcal{O}_0) \cong \mathbb{Z}[S_n]$.

Note. $\{[L(w)] : w \in S_n\}$ is the **natural** basis in $\text{Gr}(\mathcal{O}_0)$.

$$\Delta(w) := M(w \cdot 0)$$

Fact. $\{[\Delta(w)] : w \in S_n\}$ is the standard basis in $\text{Gr}(\mathcal{O}_0)$.

Reason: $[\Delta(x) : L(y)] \neq 0$ implies $x \leq y$ and $[\Delta(x) : L(x)] = 1$.

Fact. \mathcal{O}_0 has finite global dimension.

$P(w)$ — the indecomposable projective cover of $L(w)$

Corollary. $\{[P(w)] : w \in S_n\}$ is a basis in $\text{Gr}(\mathcal{O}_0)$.

Grothendieck group and different bases

Corollary. $\text{Gr}(\mathcal{O}_0) \cong \mathbb{Z}[S_n]$.

Note. $\{[L(w)] : w \in S_n\}$ is the **natural** basis in $\text{Gr}(\mathcal{O}_0)$.

$$\Delta(w) := M(w \cdot 0)$$

Fact. $\{[\Delta(w)] : w \in S_n\}$ is the **standard** basis in $\text{Gr}(\mathcal{O}_0)$.

Reason: $[\Delta(x) : L(y)] \neq 0$ implies $x \leq y$ and $[\Delta(x) : L(x)] = 1$.

Fact. \mathcal{O}_0 has finite global dimension.

$P(w)$ — the indecomposable projective cover of $L(w)$

Corollary. $\{[P(w)] : w \in S_n\}$ is a basis in $\text{Gr}(\mathcal{O}_0)$.

Grothendieck group and different bases

Corollary. $\mathrm{Gr}(\mathcal{O}_0) \cong \mathbb{Z}[S_n]$.

Note. $\{[L(w)] : w \in S_n\}$ is the **natural** basis in $\mathrm{Gr}(\mathcal{O}_0)$.

$$\Delta(w) := M(w \cdot 0)$$

Fact. $\{[\Delta(w)] : w \in S_n\}$ is the **standard** basis in $\mathrm{Gr}(\mathcal{O}_0)$.

Reason: $[\Delta(x) : L(y)] \neq 0$ implies $x \leq y$ and $[\Delta(x) : L(x)] = 1$.

Fact. \mathcal{O}_0 has finite global dimension.

$P(w)$ — the indecomposable **projective cover** of $L(w)$

Corollary. $\{[P(w)] : w \in S_n\}$ is a basis in $\mathrm{Gr}(\mathcal{O}_0)$.

Grothendieck group and different bases

Corollary. $\mathrm{Gr}(\mathcal{O}_0) \cong \mathbb{Z}[S_n]$.

Note. $\{[L(w)] : w \in S_n\}$ is the **natural** basis in $\mathrm{Gr}(\mathcal{O}_0)$.

$$\Delta(w) := M(w \cdot 0)$$

Fact. $\{[\Delta(w)] : w \in S_n\}$ is the **standard** basis in $\mathrm{Gr}(\mathcal{O}_0)$.

Reason: $[\Delta(x) : L(y)] \neq 0$ implies $x \leq y$ and $[\Delta(x) : L(x)] = 1$.

Fact. \mathcal{O}_0 has **finite** global dimension.

$P(w)$ — the indecomposable **projective cover** of $L(w)$

Corollary. $\{[P(w)] : w \in S_n\}$ is a basis in $\mathrm{Gr}(\mathcal{O}_0)$.

Grothendieck group and different bases

Corollary. $\text{Gr}(\mathcal{O}_0) \cong \mathbb{Z}[S_n]$.

Note. $\{[L(w)] : w \in S_n\}$ is the **natural** basis in $\text{Gr}(\mathcal{O}_0)$.

$$\Delta(w) := M(w \cdot 0)$$

Fact. $\{[\Delta(w)] : w \in S_n\}$ is the **standard** basis in $\text{Gr}(\mathcal{O}_0)$.

Reason: $[\Delta(x) : L(y)] \neq 0$ implies $x \leq y$ and $[\Delta(x) : L(x)] = 1$.

Fact. \mathcal{O}_0 has **finite** global dimension.

$P(w)$ — the indecomposable **projective cover** of $L(w)$

Corollary. $\{[P(w)] : w \in S_n\}$ is a basis in $\text{Gr}(\mathcal{O}_0)$.

Grothendieck group and different bases

Corollary. $\mathrm{Gr}(\mathcal{O}_0) \cong \mathbb{Z}[S_n]$.

Note. $\{[L(w)] : w \in S_n\}$ is the **natural** basis in $\mathrm{Gr}(\mathcal{O}_0)$.

$$\Delta(w) := M(w \cdot 0)$$

Fact. $\{[\Delta(w)] : w \in S_n\}$ is the **standard** basis in $\mathrm{Gr}(\mathcal{O}_0)$.

Reason: $[\Delta(x) : L(y)] \neq 0$ implies $x \leq y$ and $[\Delta(x) : L(x)] = 1$.

Fact. \mathcal{O}_0 has **finite** global dimension.

$P(w)$ — the indecomposable **projective cover** of $L(w)$

Corollary. $\{[P(w)] : w \in S_n\}$ is a basis in $\mathrm{Gr}(\mathcal{O}_0)$.

Grothendieck group and different bases

Corollary. $\text{Gr}(\mathcal{O}_0) \cong \mathbb{Z}[S_n]$.

Note. $\{[L(w)] : w \in S_n\}$ is the **natural** basis in $\text{Gr}(\mathcal{O}_0)$.

$$\Delta(w) := M(w \cdot 0)$$

Fact. $\{[\Delta(w)] : w \in S_n\}$ is the **standard** basis in $\text{Gr}(\mathcal{O}_0)$.

Reason: $[\Delta(x) : L(y)] \neq 0$ implies $x \leq y$ and $[\Delta(x) : L(x)] = 1$.

Fact. \mathcal{O}_0 has **finite** global dimension.

$P(w)$ — the indecomposable **projective cover** of $L(w)$

Corollary. $\{[P(w)] : w \in S_n\}$ is a basis in $\text{Gr}(\mathcal{O}_0)$.

Tilting modules

Theorem. [Collingwood-Irving, Ringel] For $w \in S_n$ there is a unique indecomposable module $T(w)$ such that

- ▶ $\Delta(w) \subset T(w)$ and the cokernel has a Verma flag;
- ▶ $T(w)$ is self-dual.

$T(w)$ — **tilting module**

Fact. $\{[T(w)] : w \in S_n\}$ is a basis in $\text{Gr}(\mathcal{O}_0)$.

Reason: Extensions between Vermas are directed.

Question. Which bases in $\mathbb{Z}[S_n]$ correspond to:

- ▶ $\{[L(w)] : w \in S_n\}$?
- ▶ $\{[\Delta(w)] : w \in S_n\}$?
- ▶ $\{[P(w)] : w \in S_n\}$?
- ▶ $\{[T(w)] : w \in S_n\}$?

Tilting modules

Theorem. [Collingwood-Irving, Ringel] For $w \in S_n$ there is a unique indecomposable module $T(w)$ such that

- ▶ $\Delta(w) \subset T(w)$ and the cokernel has a Verma flag;
- ▶ $T(w)$ is self-dual.

$T(w)$ — tilting module

Fact. $\{[T(w)] : w \in S_n\}$ is a basis in $\text{Gr}(\mathcal{O}_0)$.

Reason: Extensions between Vermas are directed.

Question. Which bases in $\mathbb{Z}[S_n]$ correspond to:

- ▶ $\{[L(w)] : w \in S_n\}$?
- ▶ $\{[\Delta(w)] : w \in S_n\}$?
- ▶ $\{[P(w)] : w \in S_n\}$?
- ▶ $\{[T(w)] : w \in S_n\}$?

Tilting modules

Theorem. [Collingwood-Irving, Ringel] For $w \in S_n$ there is a unique indecomposable module $T(w)$ such that

- ▶ $\Delta(w) \subset T(w)$ and the cokernel has a Verma flag;
- ▶ $T(w)$ is self-dual.

$T(w)$ — tilting module

Fact. $\{[T(w)] : w \in S_n\}$ is a basis in $\text{Gr}(\mathcal{O}_0)$.

Reason: Extensions between Vermas are directed.

Question. Which bases in $\mathbb{Z}[S_n]$ correspond to:

- ▶ $\{[L(w)] : w \in S_n\}$?
- ▶ $\{[\Delta(w)] : w \in S_n\}$?
- ▶ $\{[P(w)] : w \in S_n\}$?
- ▶ $\{[T(w)] : w \in S_n\}$?

Tilting modules

Theorem. [Collingwood-Irving, Ringel] For $w \in S_n$ there is a unique indecomposable module $T(w)$ such that

- ▶ $\Delta(w) \subset T(w)$ and the cokernel has a Verma flag;
- ▶ $T(w)$ is self-dual.

$T(w)$ — tilting module

Fact. $\{[T(w)] : w \in S_n\}$ is a basis in $\text{Gr}(\mathcal{O}_0)$.

Reason: Extensions between Vermas are directed.

Question. Which bases in $\mathbb{Z}[S_n]$ correspond to:

- ▶ $\{[L(w)] : w \in S_n\}$?
- ▶ $\{[\Delta(w)] : w \in S_n\}$?
- ▶ $\{[P(w)] : w \in S_n\}$?
- ▶ $\{[T(w)] : w \in S_n\}$?

Tilting modules

Theorem. [Collingwood-Irving, Ringel] For $w \in S_n$ there is a unique indecomposable module $T(w)$ such that

- ▶ $\Delta(w) \subset T(w)$ and the cokernel has a Verma flag;
- ▶ $T(w)$ is self-dual.

$T(w)$ — **tilting module**

Fact. $\{[T(w)] : w \in S_n\}$ is a basis in $\text{Gr}(\mathcal{O}_0)$.

Reason: Extensions between Vermas are directed.

Question. Which bases in $\mathbb{Z}[S_n]$ correspond to:

- ▶ $\{[L(w)] : w \in S_n\}$?
- ▶ $\{[\Delta(w)] : w \in S_n\}$?
- ▶ $\{[P(w)] : w \in S_n\}$?
- ▶ $\{[T(w)] : w \in S_n\}$?

Tilting modules

Theorem. [Collingwood-Irving, Ringel] For $w \in S_n$ there is a unique indecomposable module $T(w)$ such that

- ▶ $\Delta(w) \subset T(w)$ and the cokernel has a Verma flag;
- ▶ $T(w)$ is self-dual.

$T(w)$ — **tilting module**

Fact. $\{[T(w)] : w \in S_n\}$ is a basis in $\text{Gr}(\mathcal{O}_0)$.

Reason: Extensions between Vermas are directed.

Question. Which bases in $\mathbb{Z}[S_n]$ correspond to:

- ▶ $\{[L(w)] : w \in S_n\}$?
- ▶ $\{[\Delta(w)] : w \in S_n\}$?
- ▶ $\{[P(w)] : w \in S_n\}$?
- ▶ $\{[T(w)] : w \in S_n\}$?

Tilting modules

Theorem. [Collingwood-Irving, Ringel] For $w \in S_n$ there is a unique indecomposable module $T(w)$ such that

- ▶ $\Delta(w) \subset T(w)$ and the cokernel has a Verma flag;
- ▶ $T(w)$ is self-dual.

$T(w)$ — **tilting module**

Fact. $\{[T(w)] : w \in S_n\}$ is a basis in $\text{Gr}(\mathcal{O}_0)$.

Reason: Extensions between Vermas are directed.

Question. Which bases in $\mathbb{Z}[S_n]$ correspond to:

- ▶ $\{[L(w)] : w \in S_n\}$?
- ▶ $\{[\Delta(w)] : w \in S_n\}$?
- ▶ $\{[P(w)] : w \in S_n\}$?
- ▶ $\{[T(w)] : w \in S_n\}$?

Tilting modules

Theorem. [Collingwood-Irving, Ringel] For $w \in S_n$ there is a unique indecomposable module $T(w)$ such that

- ▶ $\Delta(w) \subset T(w)$ and the cokernel has a Verma flag;
- ▶ $T(w)$ is self-dual.

$T(w)$ — **tilting module**

Fact. $\{[T(w)] : w \in S_n\}$ is a basis in $\text{Gr}(\mathcal{O}_0)$.

Reason: Extensions between Vermas are directed.

Question. Which bases in $\mathbb{Z}[S_n]$ correspond to:

- ▶ $\{[L(w)] : w \in S_n\}$?
- ▶ $\{[\Delta(w)] : w \in S_n\}$?
- ▶ $\{[P(w)] : w \in S_n\}$?
- ▶ $\{[T(w)] : w \in S_n\}$?

Tilting modules

Theorem. [Collingwood-Irving, Ringel] For $w \in S_n$ there is a unique indecomposable module $T(w)$ such that

- ▶ $\Delta(w) \subset T(w)$ and the cokernel has a Verma flag;
- ▶ $T(w)$ is self-dual.

$T(w)$ — **tilting module**

Fact. $\{[T(w)] : w \in S_n\}$ is a basis in $\text{Gr}(\mathcal{O}_0)$.

Reason: Extensions between Vermas are directed.

Question. Which bases in $\mathbb{Z}[S_n]$ correspond to:

- ▶ $\{[L(w)] : w \in S_n\}$?
- ▶ $\{[\Delta(w)] : w \in S_n\}$?
- ▶ $\{[P(w)] : w \in S_n\}$?
- ▶ $\{[T(w)] : w \in S_n\}$?

Tilting modules

Theorem. [Collingwood-Irving, Ringel] For $w \in S_n$ there is a unique indecomposable module $T(w)$ such that

- ▶ $\Delta(w) \subset T(w)$ and the cokernel has a Verma flag;
- ▶ $T(w)$ is self-dual.

$T(w)$ — **tilting module**

Fact. $\{[T(w)] : w \in S_n\}$ is a basis in $\text{Gr}(\mathcal{O}_0)$.

Reason: Extensions between Vermas are directed.

Question. Which bases in $\mathbb{Z}[S_n]$ correspond to:

- ▶ $\{[L(w)] : w \in S_n\}$?
- ▶ $\{[\Delta(w)] : w \in S_n\}$?
- ▶ $\{[P(w)] : w \in S_n\}$?
- ▶ $\{[T(w)] : w \in S_n\}$?

Tilting modules

Theorem. [Collingwood-Irving, Ringel] For $w \in S_n$ there is a unique indecomposable module $T(w)$ such that

- ▶ $\Delta(w) \subset T(w)$ and the cokernel has a Verma flag;
- ▶ $T(w)$ is self-dual.

$T(w)$ — **tilting module**

Fact. $\{[T(w)] : w \in S_n\}$ is a basis in $\text{Gr}(\mathcal{O}_0)$.

Reason: Extensions between Vermas are directed.

Question. Which bases in $\mathbb{Z}[S_n]$ correspond to:

- ▶ $\{[L(w)] : w \in S_n\}$?
- ▶ $\{[\Delta(w)] : w \in S_n\}$?
- ▶ $\{[P(w)] : w \in S_n\}$?
- ▶ $\{[T(w)] : w \in S_n\}$?

Tilting modules

Theorem. [Collingwood-Irving, Ringel] For $w \in S_n$ there is a unique indecomposable module $T(w)$ such that

- ▶ $\Delta(w) \subset T(w)$ and the cokernel has a Verma flag;
- ▶ $T(w)$ is self-dual.

$T(w)$ — **tilting module**

Fact. $\{[T(w)] : w \in S_n\}$ is a basis in $\text{Gr}(\mathcal{O}_0)$.

Reason: Extensions between Vermas are directed.

Question. Which bases in $\mathbb{Z}[S_n]$ correspond to:

- ▶ $\{[L(w)] : w \in S_n\}$?
- ▶ $\{[\Delta(w)] : w \in S_n\}$?
- ▶ $\{[P(w)] : w \in S_n\}$?
- ▶ $\{[T(w)] : w \in S_n\}$?

Tilting modules

Theorem. [Collingwood-Irving, Ringel] For $w \in S_n$ there is a unique indecomposable module $T(w)$ such that

- ▶ $\Delta(w) \subset T(w)$ and the cokernel has a Verma flag;
- ▶ $T(w)$ is self-dual.

$T(w)$ — **tilting module**

Fact. $\{[T(w)] : w \in S_n\}$ is a basis in $\text{Gr}(\mathcal{O}_0)$.

Reason: Extensions between Vermas are directed.

Question. Which bases in $\mathbb{Z}[S_n]$ correspond to:

- ▶ $\{[L(w)] : w \in S_n\}$?
- ▶ $\{[\Delta(w)] : w \in S_n\}$?
- ▶ $\{[P(w)] : w \in S_n\}$?
- ▶ $\{[T(w)] : w \in S_n\}$?

Projective functors

Recall: A **projective functor** $\theta : \mathcal{O}_0 \rightarrow \mathcal{O}_0$ is a direct summand of $V \otimes_{\mathbb{C}} -$, where V is finite dimensional

Theorem. [Bernstein-S. Gelfand]

(a) There is a unique (up to isomorphism) indecomposable projective functor θ_w such that $\theta_w P(e) \cong P(w)$.

(b) $\{\theta_w : w \in S_n\}$ is a cross-section of isomorphism classes of indecomposable projective endofunctors of \mathcal{O}_0 .

Definition. \mathcal{P} — the category of projective functors.

Note: \mathcal{P} is additive, idempotent split with finitely many indecomposables up to isomorphism, it has the structure of a tensor category.

Projective functors

Recall: A **projective functor** $\theta : \mathcal{O}_0 \rightarrow \mathcal{O}_0$ is a direct summand of $V \otimes_{\mathbb{C}} -$, where V is finite dimensional

Theorem. [Bernstein-S. Gelfand]

(a) There is a unique (up to isomorphism) indecomposable projective functor θ_w such that $\theta_w P(e) \cong P(w)$.

(b) $\{\theta_w : w \in S_n\}$ is a cross-section of isomorphism classes of indecomposable projective endofunctors of \mathcal{O}_0 .

Definition. \mathcal{P} — the category of projective functors.

Note: \mathcal{P} is additive, idempotent split with finitely many indecomposables up to isomorphism, it has the structure of a tensor category.

Projective functors

Recall: A **projective functor** $\theta : \mathcal{O}_0 \rightarrow \mathcal{O}_0$ is a direct summand of $V \otimes_{\mathbb{C}} -$, where V is finite dimensional

Theorem. [Bernstein-S. Gelfand]

(a) There is a unique (up to isomorphism) indecomposable projective functor θ_w such that $\theta_w P(e) \cong P(w)$.

(b) $\{\theta_w : w \in S_n\}$ is a cross-section of isomorphism classes of indecomposable projective endofunctors of \mathcal{O}_0 .

Definition. \mathcal{P} — the category of projective functors.

Note: \mathcal{P} is additive, idempotent split with finitely many indecomposables up to isomorphism, it has the structure of a tensor category.

Projective functors

Recall: A **projective functor** $\theta : \mathcal{O}_0 \rightarrow \mathcal{O}_0$ is a direct summand of $V \otimes_{\mathbb{C}} -$, where V is finite dimensional

Theorem. [Bernstein-S. Gelfand]

(a) There is a unique (up to isomorphism) indecomposable projective functor θ_w such that $\theta_w P(e) \cong P(w)$.

(b) $\{\theta_w : w \in S_n\}$ is a cross-section of isomorphism classes of indecomposable projective endofunctors of \mathcal{O}_0 .

Definition. \mathcal{P} — the category of projective functors.

Note: \mathcal{P} is additive, idempotent split with finitely many indecomposables up to isomorphism, it has the structure of a tensor category.

Projective functors

Recall: A **projective functor** $\theta : \mathcal{O}_0 \rightarrow \mathcal{O}_0$ is a direct summand of $V \otimes_{\mathbb{C}} -$, where V is finite dimensional

Theorem. [Bernstein-S. Gelfand]

(a) There is a unique (up to isomorphism) indecomposable projective functor θ_w such that $\theta_w P(e) \cong P(w)$.

(b) $\{\theta_w : w \in S_n\}$ is a cross-section of isomorphism classes of indecomposable projective endofunctors of \mathcal{O}_0 .

Definition. \mathcal{P} — the category of projective functors.

Note: \mathcal{P} is additive, idempotent split with finitely many indecomposables up to isomorphism, it has the structure of a tensor category.

Projective functors

Recall: A **projective functor** $\theta : \mathcal{O}_0 \rightarrow \mathcal{O}_0$ is a direct summand of $V \otimes_{\mathbb{C}} -$, where V is finite dimensional

Theorem. [Bernstein-S. Gelfand]

(a) There is a unique (up to isomorphism) indecomposable projective functor θ_w such that $\theta_w P(e) \cong P(w)$.

(b) $\{\theta_w : w \in S_n\}$ is a cross-section of isomorphism classes of indecomposable projective endofunctors of \mathcal{O}_0 .

Definition. \mathcal{P} — the category of projective functors.

Note: \mathcal{P} is additive, idempotent split with finitely many indecomposables up to isomorphism, it has the structure of a tensor category.

Projective functors

Recall: A **projective functor** $\theta : \mathcal{O}_0 \rightarrow \mathcal{O}_0$ is a direct summand of $V \otimes_{\mathbb{C}} -$, where V is finite dimensional

Theorem. [Bernstein-S. Gelfand]

(a) There is a unique (up to isomorphism) indecomposable projective functor θ_w such that $\theta_w P(e) \cong P(w)$.

(b) $\{\theta_w : w \in S_n\}$ is a cross-section of isomorphism classes of indecomposable projective endofunctors of \mathcal{O}_0 .

Definition. \mathcal{P} — the category of projective functors.

Note: \mathcal{P} is additive, idempotent split with finitely many indecomposables up to isomorphism, it has the structure of a tensor category.

Projective functors

Recall: A **projective functor** $\theta : \mathcal{O}_0 \rightarrow \mathcal{O}_0$ is a direct summand of $V \otimes_{\mathbb{C}} -$, where V is finite dimensional

Theorem. [Bernstein-S. Gelfand]

(a) There is a unique (up to isomorphism) indecomposable projective functor θ_w such that $\theta_w P(e) \cong P(w)$.

(b) $\{\theta_w : w \in S_n\}$ is a cross-section of isomorphism classes of indecomposable projective endofunctors of \mathcal{O}_0 .

Definition. \mathcal{P} — the category of projective functors.

Note: \mathcal{P} is additive, idempotent split with finitely many indecomposables up to isomorphism, it has the structure of a tensor category.

Combinatorics of projective functors

Observation. For s simple reflection and $w \in S_n$ there are s.e.s.

$$\Delta(ws) \hookrightarrow \theta_s \Delta(w) \rightarrow \Delta(w) \text{ if } ws > w,$$

$$\Delta(w) \hookrightarrow \theta_s \Delta(w) \rightarrow \Delta(ws) \text{ if } ws < w.$$

Fact. \mathcal{P} is generated by θ_s , s simple reflection, as a tensor category.

Corollary. $\text{Gr}_{\mathbb{Q}}(\mathcal{P}) \cong \mathbb{Z}[S_n]$

Question. Which basis of $\mathbb{Z}[S_n]$ is $\{[\theta_w], w \in S_n\}$?

“Answer”: The same as $\{[P(w)], w \in S_n\}$.

Combinatorics of projective functors

Observation. For s simple reflection and $w \in S_n$ there are s.e.s.

$$\Delta(ws) \hookrightarrow \theta_s \Delta(w) \rightarrow \Delta(w) \text{ if } ws > w,$$

$$\Delta(w) \hookrightarrow \theta_s \Delta(w) \rightarrow \Delta(ws) \text{ if } ws < w.$$

Fact. \mathcal{P} is generated by θ_s , s simple reflection, as a tensor category.

Corollary. $\text{Gr}_{\mathbb{Q}}(\mathcal{P}) \cong \mathbb{Z}[S_n]$

Question. Which basis of $\mathbb{Z}[S_n]$ is $\{[\theta_w], w \in S_n\}$?

“Answer”: The same as $\{[P(w)], w \in S_n\}$.

Combinatorics of projective functors

Observation. For s simple reflection and $w \in S_n$ there are s.e.s.

$$\Delta(ws) \hookrightarrow \theta_s \Delta(w) \twoheadrightarrow \Delta(w) \text{ if } ws > w,$$

$$\Delta(w) \hookrightarrow \theta_s \Delta(w) \twoheadrightarrow \Delta(ws) \text{ if } ws < w.$$

Fact. \mathcal{P} is generated by θ_s , s simple reflection, as a tensor category.

Corollary. $\text{Gr}_{\mathbb{Q}}(\mathcal{P}) \cong \mathbb{Z}[S_n]$

Question. Which basis of $\mathbb{Z}[S_n]$ is $\{[\theta_w], w \in S_n\}$?

“Answer”: The same as $\{[P(w)], w \in S_n\}$.

Combinatorics of projective functors

Observation. For s simple reflection and $w \in S_n$ there are s.e.s.

$$\Delta(ws) \hookrightarrow \theta_s \Delta(w) \twoheadrightarrow \Delta(w) \text{ if } ws > w,$$

$$\Delta(w) \hookrightarrow \theta_s \Delta(w) \twoheadrightarrow \Delta(ws) \text{ if } ws < w.$$

Fact. \mathcal{P} is generated by θ_s , s simple reflection, as a tensor category.

Corollary. $\text{Gr}_{\oplus}(\mathcal{P}) \cong \mathbb{Z}[S_n]$

Question. Which basis of $\mathbb{Z}[S_n]$ is $\{[\theta_w], w \in S_n\}$?

“Answer”: The same as $\{[P(w)], w \in S_n\}$.

Combinatorics of projective functors

Observation. For s simple reflection and $w \in S_n$ there are s.e.s.

$$\Delta(ws) \hookrightarrow \theta_s \Delta(w) \twoheadrightarrow \Delta(w) \text{ if } ws > w,$$

$$\Delta(w) \hookrightarrow \theta_s \Delta(w) \twoheadrightarrow \Delta(ws) \text{ if } ws < w.$$

Fact. \mathcal{P} is generated by θ_s , s simple reflection, as a tensor category.

Corollary. $\text{Gr}_{\oplus}(\mathcal{P}) \cong \mathbb{Z}[S_n]$

Question. Which basis of $\mathbb{Z}[S_n]$ is $\{[\theta_w], w \in S_n\}$?

“Answer”: The same as $\{[P(w)], w \in S_n\}$.

Combinatorics of projective functors

Observation. For s simple reflection and $w \in S_n$ there are s.e.s.

$$\Delta(ws) \hookrightarrow \theta_s \Delta(w) \twoheadrightarrow \Delta(w) \text{ if } ws > w,$$

$$\Delta(w) \hookrightarrow \theta_s \Delta(w) \twoheadrightarrow \Delta(ws) \text{ if } ws < w.$$

Fact. \mathcal{P} is generated by θ_s , s simple reflection, as a tensor category.

Corollary. $\text{Gr}_{\oplus}(\mathcal{P}) \cong \mathbb{Z}[S_n]$

Question. Which basis of $\mathbb{Z}[S_n]$ is $\{[\theta_w], w \in S_n\}$?

“Answer”: The same as $\{[P(w)], w \in S_n\}$.

Combinatorics of projective functors

Observation. For s simple reflection and $w \in S_n$ there are s.e.s.

$$\Delta(ws) \hookrightarrow \theta_s \Delta(w) \twoheadrightarrow \Delta(w) \text{ if } ws > w,$$

$$\Delta(w) \hookrightarrow \theta_s \Delta(w) \twoheadrightarrow \Delta(ws) \text{ if } ws < w.$$

Fact. \mathcal{P} is generated by θ_s , s simple reflection, as a tensor category.

Corollary. $\text{Gr}_{\oplus}(\mathcal{P}) \cong \mathbb{Z}[S_n]$

Question. Which basis of $\mathbb{Z}[S_n]$ is $\{[\theta_w], w \in S_n\}$?

“Answer”: The same as $\{[P(w)], w \in S_n\}$.

Combinatorics of projective functors

Observation. For s simple reflection and $w \in S_n$ there are s.e.s.

$$\Delta(ws) \hookrightarrow \theta_s \Delta(w) \twoheadrightarrow \Delta(w) \text{ if } ws > w,$$

$$\Delta(w) \hookrightarrow \theta_s \Delta(w) \twoheadrightarrow \Delta(ws) \text{ if } ws < w.$$

Fact. \mathcal{P} is generated by θ_s , s simple reflection, as a tensor category.

Corollary. $\text{Gr}_{\oplus}(\mathcal{P}) \cong \mathbb{Z}[S_n]$

Question. Which basis of $\mathbb{Z}[S_n]$ is $\{[\theta_w], w \in S_n\}$?

“Answer”: The **same** as $\{[P(w)], w \in S_n\}$.

Combinatorics of projective functors

Observation. For s simple reflection and $w \in S_n$ there are s.e.s.

$$\Delta(ws) \hookrightarrow \theta_s \Delta(w) \twoheadrightarrow \Delta(w) \text{ if } ws > w,$$

$$\Delta(w) \hookrightarrow \theta_s \Delta(w) \twoheadrightarrow \Delta(ws) \text{ if } ws < w.$$

Fact. \mathcal{P} is generated by θ_s , s simple reflection, as a tensor category.

Corollary. $\text{Gr}_{\oplus}(\mathcal{P}) \cong \mathbb{Z}[S_n]$

Question. Which basis of $\mathbb{Z}[S_n]$ is $\{[\theta_w], w \in S_n\}$?

“Answer”: The **same** as $\{[P(w)], w \in S_n\}$.

Categorification of the right regular $\mathbb{Z}[S_n]$ -module

Note. Projective functors are exact.

Consequence. Each $[\theta_w]$ is an endomorphism of $\text{Gr}(\mathcal{O}_0)$

Identify: $\text{Gr}(\mathcal{O}_0)$ with $\mathbb{Z}[S_n]$ via $[\Delta(w)] \mapsto w$.

Identify: $\text{Gr}_{\oplus}(\mathcal{P})$ with $\mathbb{Z}[S_n]$ via $[\theta_s] \mapsto (e + s)$.

Theorem The action of \mathcal{P} on \mathcal{O}_0 is a **categorification** of the right regular $\mathbb{Z}[S_n]$ -module.

Diagrammatically:

$$\mathcal{O}_0 \begin{array}{c} \curvearrowright \\ \mathcal{P} \end{array} \xrightarrow{\text{Gr}} \mathbb{Z}[S_n] \begin{array}{c} \curvearrowright \\ \mathbb{Z}[S_n] \end{array}$$

Categorification of the right regular $\mathbb{Z}[S_n]$ -module

Note. Projective functors are exact.

Consequence. Each $[\theta_w]$ is an endomorphism of $\text{Gr}(\mathcal{O}_0)$

Identify: $\text{Gr}(\mathcal{O}_0)$ with $\mathbb{Z}[S_n]$ via $[\Delta(w)] \mapsto w$.

Identify: $\text{Gr}_{\oplus}(\mathcal{P})$ with $\mathbb{Z}[S_n]$ via $[\theta_s] \mapsto (e + s)$.

Theorem The action of \mathcal{P} on \mathcal{O}_0 is a **categorification** of the right regular $\mathbb{Z}[S_n]$ -module.

Diagrammatically:

$$\mathcal{O}_0 \begin{array}{c} \curvearrowright \\ \mathcal{P} \end{array} \xrightarrow{\text{Gr}} \mathbb{Z}[S_n] \begin{array}{c} \curvearrowright \\ \mathbb{Z}[S_n] \end{array}$$

Categorification of the right regular $\mathbb{Z}[S_n]$ -module

Note. Projective functors are exact.

Consequence. Each $[\theta_w]$ is an endomorphism of $\text{Gr}(\mathcal{O}_0)$

Identify: $\text{Gr}(\mathcal{O}_0)$ with $\mathbb{Z}[S_n]$ via $[\Delta(w)] \mapsto w$.

Identify: $\text{Gr}_{\oplus}(\mathcal{P})$ with $\mathbb{Z}[S_n]$ via $[\theta_s] \mapsto (e + s)$.

Theorem The action of \mathcal{P} on \mathcal{O}_0 is a **categorification** of the right regular $\mathbb{Z}[S_n]$ -module.

Diagrammatically:

$$\mathcal{O}_0 \begin{array}{c} \curvearrowright \\ \mathcal{P} \end{array} \xrightarrow{\text{Gr}} \mathbb{Z}[S_n] \begin{array}{c} \curvearrowright \\ \mathbb{Z}[S_n] \end{array}$$

Categorification of the right regular $\mathbb{Z}[S_n]$ -module

Note. Projective functors are exact.

Consequence. Each $[\theta_w]$ is an endomorphism of $\text{Gr}(\mathcal{O}_0)$

Identify: $\text{Gr}(\mathcal{O}_0)$ with $\mathbb{Z}[S_n]$ via $[\Delta(w)] \mapsto w$.

Identify: $\text{Gr}_{\oplus}(\mathcal{P})$ with $\mathbb{Z}[S_n]$ via $[\theta_s] \mapsto (e + s)$.

Theorem The action of \mathcal{P} on \mathcal{O}_0 is a **categorification** of the right regular $\mathbb{Z}[S_n]$ -module.

Diagrammatically:

$$\mathcal{O}_0 \begin{array}{c} \curvearrowright \\ \mathcal{P} \end{array} \xrightarrow{\text{Gr}} \mathbb{Z}[S_n] \begin{array}{c} \curvearrowright \\ \mathbb{Z}[S_n] \end{array}$$

Categorification of the right regular $\mathbb{Z}[S_n]$ -module

Note. Projective functors are exact.

Consequence. Each $[\theta_w]$ is an endomorphism of $\text{Gr}(\mathcal{O}_0)$

Identify: $\text{Gr}(\mathcal{O}_0)$ with $\mathbb{Z}[S_n]$ via $[\Delta(w)] \mapsto w$.

Identify: $\text{Gr}_{\oplus}(\mathcal{P})$ with $\mathbb{Z}[S_n]$ via $[\theta_s] \mapsto (e + s)$.

Theorem The action of \mathcal{P} on \mathcal{O}_0 is a **categorification** of the right regular $\mathbb{Z}[S_n]$ -module.

Diagrammatically:

$$\mathcal{O}_0 \begin{array}{c} \curvearrowright \\ \mathcal{P} \end{array} \xrightarrow{\text{Gr}} \mathbb{Z}[S_n] \begin{array}{c} \curvearrowright \\ \mathbb{Z}[S_n] \end{array}$$

Categorification of the right regular $\mathbb{Z}[S_n]$ -module

Note. Projective functors are exact.

Consequence. Each $[\theta_w]$ is an endomorphism of $\text{Gr}(\mathcal{O}_0)$

Identify: $\text{Gr}(\mathcal{O}_0)$ with $\mathbb{Z}[S_n]$ via $[\Delta(w)] \mapsto w$.

Identify: $\text{Gr}_{\oplus}(\mathcal{P})$ with $\mathbb{Z}[S_n]$ via $[\theta_s] \mapsto (e + s)$.

Theorem The action of \mathcal{P} on \mathcal{O}_0 is a **categorification** of the **right regular** $\mathbb{Z}[S_n]$ -module.

Diagrammatically:

$$\mathcal{O}_0 \xleftarrow{\mathcal{P}} \text{Gr} \xrightarrow{\text{Gr}} \mathbb{Z}[S_n] \xleftarrow{\mathbb{Z}[S_n]} \mathbb{Z}[S_n]$$

Categorification of the right regular $\mathbb{Z}[S_n]$ -module

Note. Projective functors are exact.

Consequence. Each $[\theta_w]$ is an endomorphism of $\text{Gr}(\mathcal{O}_0)$

Identify: $\text{Gr}(\mathcal{O}_0)$ with $\mathbb{Z}[S_n]$ via $[\Delta(w)] \mapsto w$.

Identify: $\text{Gr}_{\oplus}(\mathcal{P})$ with $\mathbb{Z}[S_n]$ via $[\theta_s] \mapsto (e + s)$.

Theorem The action of \mathcal{P} on \mathcal{O}_0 is a **categorification** of the **right regular** $\mathbb{Z}[S_n]$ -module.

Diagrammatically:

$$\mathcal{O}_0 \xleftarrow{\mathcal{P}} \text{Gr} \xrightarrow{\text{Gr}} \mathbb{Z}[S_n] \xleftarrow{\mathbb{Z}[S_n]} \mathbb{Z}[S_n]$$

Categorification of the right regular $\mathbb{Z}[S_n]$ -module

Note. Projective functors are exact.

Consequence. Each $[\theta_w]$ is an endomorphism of $\text{Gr}(\mathcal{O}_0)$

Identify: $\text{Gr}(\mathcal{O}_0)$ with $\mathbb{Z}[S_n]$ via $[\Delta(w)] \mapsto w$.

Identify: $\text{Gr}_{\oplus}(\mathcal{P})$ with $\mathbb{Z}[S_n]$ via $[\theta_s] \mapsto (e + s)$.

Theorem The action of \mathcal{P} on \mathcal{O}_0 is a **categorification** of the **right regular** $\mathbb{Z}[S_n]$ -module.

Diagrammatically:

$$\mathcal{O}_0 \begin{array}{c} \longleftarrow \\ \curvearrowright_{\mathcal{P}} \end{array} \xrightarrow{\text{Gr}} \mathbb{Z}[S_n] \begin{array}{c} \longleftarrow \\ \curvearrowright_{\mathbb{Z}[S_n]} \end{array}$$

Categorification of the right regular $\mathbb{Z}[S_n]$ -module

Note. Projective functors are exact.

Consequence. Each $[\theta_w]$ is an endomorphism of $\text{Gr}(\mathcal{O}_0)$

Identify: $\text{Gr}(\mathcal{O}_0)$ with $\mathbb{Z}[S_n]$ via $[\Delta(w)] \mapsto w$.

Identify: $\text{Gr}_{\oplus}(\mathcal{P})$ with $\mathbb{Z}[S_n]$ via $[\theta_s] \mapsto (e + s)$.

Theorem The action of \mathcal{P} on \mathcal{O}_0 is a **categorification** of the **right regular** $\mathbb{Z}[S_n]$ -module.

Diagrammatically:

$$\mathcal{O}_0 \begin{array}{c} \longleftarrow \\ \curvearrowright_{\mathcal{P}} \end{array} \xrightarrow{\text{Gr}} \mathbb{Z}[S_n] \begin{array}{c} \longleftarrow \\ \curvearrowright_{\mathbb{Z}[S_n]} \end{array}$$

Kazhdan-Lusztig basis

Note. The action of \mathcal{P} categorifies $\mathbb{Z}[S_n]$ and **not** S_n .

Question. What is $\{[\theta_w], w \in S_n\}$?

Answer. This is the **Kazhdan-Lusztig** basis.

Remark. This is equivalent to Kazhdan-Lusztig conjecture (=theorem).

Remark. Recent algebraic proof by **Elias-Williamson**.

Remark. To define Kazhdan-Lusztig basis one needs to deform $\mathbb{Z}[S_n]$ to the **Hecke algebra**.

Categorically this means to introduce a grading on \mathcal{O}_0 .

Kazhdan-Lusztig basis

Note. The action of \mathcal{P} categorifies $\mathbb{Z}[S_n]$ and **not** S_n .

Question. What is $\{[\theta_w], w \in S_n\}$?

Answer. This is the **Kazhdan-Lusztig** basis.

Remark. This is equivalent to Kazhdan-Lusztig conjecture (=theorem).

Remark. Recent algebraic proof by **Elias-Williamson**.

Remark. To define Kazhdan-Lusztig basis one needs to deform $\mathbb{Z}[S_n]$ to the **Hecke algebra**.

Categorically this means to introduce a grading on \mathcal{O}_0 .

Kazhdan-Lusztig basis

Note. The action of \mathcal{P} categorifies $\mathbb{Z}[S_n]$ and **not** S_n .

Question. What is $\{[\theta_w], w \in S_n\}$?

Answer. This is the **Kazhdan-Lusztig** basis.

Remark. This is equivalent to Kazhdan-Lusztig conjecture (=theorem).

Remark. Recent algebraic proof by **Elias-Williamson**.

Remark. To define Kazhdan-Lusztig basis one needs to deform $\mathbb{Z}[S_n]$ to the **Hecke algebra**.

Categorically this means to introduce a grading on \mathcal{O}_0 .

Kazhdan-Lusztig basis

Note. The action of \mathcal{P} categorifies $\mathbb{Z}[S_n]$ and **not** S_n .

Question. What is $\{[\theta_w], w \in S_n\}$?

Answer. This is the **Kazhdan-Lusztig** basis.

Remark. This is equivalent to Kazhdan-Lusztig conjecture (=theorem).

Remark. Recent algebraic proof by **Elias-Williamson**.

Remark. To define Kazhdan-Lusztig basis one needs to deform $\mathbb{Z}[S_n]$ to the **Hecke algebra**.

Categorically this means to introduce a grading on \mathcal{O}_0 .

Kazhdan-Lusztig basis

Note. The action of \mathcal{P} categorifies $\mathbb{Z}[S_n]$ and **not** S_n .

Question. What is $\{[\theta_w], w \in S_n\}$?

Answer. This is the **Kazhdan-Lusztig** basis.

Remark. This is equivalent to Kazhdan-Lusztig conjecture (=theorem).

Remark. Recent algebraic proof by **Elias-Williamson**.

Remark. To define Kazhdan-Lusztig basis one needs to deform $\mathbb{Z}[S_n]$ to the **Hecke algebra**.

Categorically this means to introduce a grading on \mathcal{O}_0 .

Kazhdan-Lusztig basis

Note. The action of \mathcal{P} categorifies $\mathbb{Z}[S_n]$ and **not** S_n .

Question. What is $\{[\theta_w], w \in S_n\}$?

Answer. This is the **Kazhdan-Lusztig** basis.

Remark. This is equivalent to Kazhdan-Lusztig conjecture (=theorem).

Remark. Recent algebraic proof by **Elias-Williamson**.

Remark. To define Kazhdan-Lusztig basis one needs to deform $\mathbb{Z}[S_n]$ to the **Hecke algebra**.

Categorically this means to introduce a **grading** on \mathcal{O}_0 .

Kazhdan-Lusztig basis

Note. The action of \mathcal{P} categorifies $\mathbb{Z}[S_n]$ and **not** S_n .

Question. What is $\{[\theta_w], w \in S_n\}$?

Answer. This is the **Kazhdan-Lusztig** basis.

Remark. This is equivalent to Kazhdan-Lusztig conjecture (=theorem).

Remark. Recent algebraic proof by **Elias-Williamson**.

Remark. To define Kazhdan-Lusztig basis one needs to **deform** $\mathbb{Z}[S_n]$ to the **Hecke algebra**.

Categorically this means to introduce a grading on \mathcal{O}_0 .

Kazhdan-Lusztig basis

Note. The action of \mathcal{P} categorifies $\mathbb{Z}[S_n]$ and **not** S_n .

Question. What is $\{[\theta_w], w \in S_n\}$?

Answer. This is the **Kazhdan-Lusztig** basis.

Remark. This is equivalent to Kazhdan-Lusztig conjecture (=theorem).

Remark. Recent algebraic proof by **Elias-Williamson**.

Remark. To define Kazhdan-Lusztig basis one needs to **deform** $\mathbb{Z}[S_n]$ to the **Hecke algebra**.

Categorically this means to introduce a **grading** on \mathcal{O}_0 .

Kazhdan-Lusztig basis

Note. The action of \mathcal{P} categorifies $\mathbb{Z}[S_n]$ and **not** S_n .

Question. What is $\{[\theta_w], w \in S_n\}$?

Answer. This is the **Kazhdan-Lusztig** basis.

Remark. This is equivalent to Kazhdan-Lusztig conjecture (=theorem).

Remark. Recent algebraic proof by **Elias-Williamson**.

Remark. To define Kazhdan-Lusztig basis one needs to **deform** $\mathbb{Z}[S_n]$ to the **Hecke algebra**.

Categorically this means to introduce a **grading** on \mathcal{O}_0 .

Alternative approach: coinvariants

$\mathbb{C}[x_1, x_2, \dots, x_n]$ — polynomial algebra

grading: $\deg(x_i) = 2$

S_n acts on $\mathbb{C}[x_1, x_2, \dots, x_n]$ by permuting indices

$\mathbb{C}[x_1, x_2, \dots, x_n]_i^{S_n}$ — invariant homogeneous polynomials of degree i

$$\mathbb{C}[x_1, x_2, \dots, x_n]_+^{S_n} = \bigoplus_{i>0} \mathbb{C}[x_1, x_2, \dots, x_n]_i^{S_n}$$

Definition. The **coinvariant** algebra is

$$\mathbf{C} := \mathbb{C}[x_1, x_2, \dots, x_n] / (\mathbb{C}[x_1, x_2, \dots, x_n]_+^{S_n}).$$

Fact. $\mathbf{C} \cong \mathbb{C}[S_n]_{S_n}$ as an S_n -module, in particular, $\dim \mathbf{C} = |S_n|$.

Alternative approach: coinvariants

$\mathbb{C}[x_1, x_2, \dots, x_n]$ — polynomial algebra

grading: $\deg(x_i) = 2$

S_n acts on $\mathbb{C}[x_1, x_2, \dots, x_n]$ by permuting indices

$\mathbb{C}[x_1, x_2, \dots, x_n]_i^{S_n}$ — invariant homogeneous polynomials of degree i

$$\mathbb{C}[x_1, x_2, \dots, x_n]_+^{S_n} = \bigoplus_{i>0} \mathbb{C}[x_1, x_2, \dots, x_n]_i^{S_n}$$

Definition. The **coinvariant** algebra is

$$\mathbf{C} := \mathbb{C}[x_1, x_2, \dots, x_n] / (\mathbb{C}[x_1, x_2, \dots, x_n]_+^{S_n}).$$

Fact. $\mathbf{C} \cong \mathbb{C}[S_n]_{S_n}$ as an S_n -module, in particular, $\dim \mathbf{C} = |S_n|$.

Alternative approach: coinvariants

$\mathbb{C}[x_1, x_2, \dots, x_n]$ — polynomial algebra

grading: $\deg(x_i) = 2$

S_n acts on $\mathbb{C}[x_1, x_2, \dots, x_n]$ by permuting indices

$\mathbb{C}[x_1, x_2, \dots, x_n]_i^{S_n}$ — invariant homogeneous polynomials of degree i

$$\mathbb{C}[x_1, x_2, \dots, x_n]_+^{S_n} = \bigoplus_{i \geq 0} \mathbb{C}[x_1, x_2, \dots, x_n]_i^{S_n}$$

Definition. The coinvariant algebra is

$$\mathbf{C} := \mathbb{C}[x_1, x_2, \dots, x_n] / (\mathbb{C}[x_1, x_2, \dots, x_n]_+^{S_n}).$$

Fact. $\mathbf{C} \cong \mathbb{C}[S_n]_{S_n}$ as an S_n -module, in particular, $\dim \mathbf{C} = |S_n|$.

Alternative approach: coinvariants

$\mathbb{C}[x_1, x_2, \dots, x_n]$ — polynomial algebra

grading: $\deg(x_i) = 2$

S_n acts on $\mathbb{C}[x_1, x_2, \dots, x_n]$ by **permuting indices**

$\mathbb{C}[x_1, x_2, \dots, x_n]_i^{S_n}$ — invariant homogeneous polynomials of degree i

$$\mathbb{C}[x_1, x_2, \dots, x_n]_+^{S_n} = \bigoplus_{i \geq 0} \mathbb{C}[x_1, x_2, \dots, x_n]_i^{S_n}$$

Definition. The **coinvariant** algebra is

$$\mathbb{C} := \mathbb{C}[x_1, x_2, \dots, x_n] / (\mathbb{C}[x_1, x_2, \dots, x_n]_+^{S_n}).$$

Fact. $\mathbb{C} \cong \mathbb{C}[S_n]_{S_n}$ as an S_n -module, in particular, $\dim \mathbb{C} = |S_n|$.

Alternative approach: coinvariants

$\mathbb{C}[x_1, x_2, \dots, x_n]$ — polynomial algebra

grading: $\deg(x_i) = 2$

S_n acts on $\mathbb{C}[x_1, x_2, \dots, x_n]$ by **permuting indices**

$\mathbb{C}[x_1, x_2, \dots, x_n]_i^{S_n}$ — **invariant** homogeneous polynomials of degree i

$$\mathbb{C}[x_1, x_2, \dots, x_n]_+^{S_n} = \bigoplus_{i \geq 0} \mathbb{C}[x_1, x_2, \dots, x_n]_i^{S_n}$$

Definition. The **coinvariant** algebra is

$$\mathbf{C} := \mathbb{C}[x_1, x_2, \dots, x_n] / (\mathbb{C}[x_1, x_2, \dots, x_n]_+^{S_n}).$$

Fact. $\mathbf{C} \cong \mathbb{C}[S_n]_{S_n}$ as an S_n -module, in particular, $\dim \mathbf{C} = |S_n|$.

Alternative approach: coinvariants

$\mathbb{C}[x_1, x_2, \dots, x_n]$ — polynomial algebra

grading: $\deg(x_i) = 2$

S_n acts on $\mathbb{C}[x_1, x_2, \dots, x_n]$ by **permuting indices**

$\mathbb{C}[x_1, x_2, \dots, x_n]_i^{S_n}$ — **invariant** homogeneous polynomials of degree i

$$\mathbb{C}[x_1, x_2, \dots, x_n]_+^{S_n} = \bigoplus_{i \geq 0} \mathbb{C}[x_1, x_2, \dots, x_n]_i^{S_n}$$

Definition. The **coinvariant** algebra is

$$\mathbf{C} := \mathbb{C}[x_1, x_2, \dots, x_n] / (\mathbb{C}[x_1, x_2, \dots, x_n]_+^{S_n}).$$

Fact. $\mathbf{C} \cong \mathbb{C}[S_n]_{S_n}$ as an S_n -module, in particular, $\dim \mathbf{C} = |S_n|$.

Alternative approach: coinvariants

$\mathbb{C}[x_1, x_2, \dots, x_n]$ — polynomial algebra

grading: $\deg(x_i) = 2$

S_n acts on $\mathbb{C}[x_1, x_2, \dots, x_n]$ by **permuting indices**

$\mathbb{C}[x_1, x_2, \dots, x_n]_i^{S_n}$ — **invariant** homogeneous polynomials of degree i

$$\mathbb{C}[x_1, x_2, \dots, x_n]_+^{S_n} = \bigoplus_{i>0} \mathbb{C}[x_1, x_2, \dots, x_n]_i^{S_n}$$

Definition. The **coinvariant** algebra is

$$\mathbf{C} := \mathbb{C}[x_1, x_2, \dots, x_n] / (\mathbb{C}[x_1, x_2, \dots, x_n]_+^{S_n}).$$

Fact. $\mathbf{C} \cong \mathbb{C}[S_n]_{S_n}$ as an S_n -module, in particular, $\dim \mathbf{C} = |S_n|$.

Alternative approach: coinvariants

$\mathbb{C}[x_1, x_2, \dots, x_n]$ — polynomial algebra

grading: $\deg(x_i) = 2$

S_n acts on $\mathbb{C}[x_1, x_2, \dots, x_n]$ by **permuting indices**

$\mathbb{C}[x_1, x_2, \dots, x_n]_i^{S_n}$ — **invariant** homogeneous polynomials of degree i

$$\mathbb{C}[x_1, x_2, \dots, x_n]_+^{S_n} = \bigoplus_{i>0} \mathbb{C}[x_1, x_2, \dots, x_n]_i^{S_n}$$

Definition. The **coinvariant** algebra is

$$\mathbf{C} := \mathbb{C}[x_1, x_2, \dots, x_n] / (\mathbb{C}[x_1, x_2, \dots, x_n]_+^{S_n}).$$

Fact. $\mathbf{C} \cong \mathbb{C}[S_n]_{S_n}$ as an S_n -module, in particular, $\dim \mathbf{C} = |S_n|$.

Alternative approach: coinvariants

$\mathbb{C}[x_1, x_2, \dots, x_n]$ — polynomial algebra

grading: $\deg(x_i) = 2$

S_n acts on $\mathbb{C}[x_1, x_2, \dots, x_n]$ by **permuting indices**

$\mathbb{C}[x_1, x_2, \dots, x_n]_i^{S_n}$ — **invariant** homogeneous polynomials of degree i

$$\mathbb{C}[x_1, x_2, \dots, x_n]_+^{S_n} = \bigoplus_{i>0} \mathbb{C}[x_1, x_2, \dots, x_n]_i^{S_n}$$

Definition. The **coinvariant** algebra is

$$\mathbf{C} := \mathbb{C}[x_1, x_2, \dots, x_n] / (\mathbb{C}[x_1, x_2, \dots, x_n]_+^{S_n}).$$

Fact. $\mathbf{C} \cong \mathbb{C}[S_n]_{S_n}$ as an S_n -module, in particular, $\dim \mathbf{C} = |S_n|$.

Alternative approach: Soergel bimodules

$s_i = (i, i + 1)$ — simple reflection in S_n for $i = 1, 2, \dots, n - 1$

Fact. S_n is a Coxeter group with generators s_i

$l : S_n \rightarrow \mathbb{Z}$ — the length function

\mathbb{C}^{s_i} — the algebra of s_i -invariants in \mathbb{C}

Definition. [Soergel] Define inductively \mathbb{C} - \mathbb{C} -bimodules B_w , $w \in S_n$ as follows:

- ▶ $B_e := \mathbb{C}$
- ▶ For $w = s_{i_1} s_{i_2} \dots s_{i_k}$ reduced decomposition, the bimodule B_w is the unique indecomposable direct summand of

$$\mathbb{C} \otimes_{\mathbb{C}^{s_{i_1}}} \mathbb{C} \otimes_{\mathbb{C}^{s_{i_2}}} \dots \otimes_{\mathbb{C}^{s_{i_k}}} \mathbb{C}$$

which is not isomorphic to B_x , where $l(x) < l(w)$.

Alternative approach: Soergel bimodules

$s_i = (i, i + 1)$ — simple reflection in S_n for $i = 1, 2, \dots, n - 1$

Fact. S_n is a Coxeter group with generators s_i

$l : S_n \rightarrow \mathbb{Z}$ — the length function

\mathbb{C}^{s_i} — the algebra of s_i -invariants in \mathbb{C}

Definition. [Soergel] Define inductively \mathbb{C} - \mathbb{C} -bimodules B_w , $w \in S_n$ as follows:

- ▶ $B_e := \mathbb{C}$
- ▶ For $w = s_{i_1} s_{i_2} \dots s_{i_k}$ reduced decomposition, the bimodule B_w is the unique indecomposable direct summand of

$$\mathbb{C} \otimes_{\mathbb{C}^{s_{i_1}}} \mathbb{C} \otimes_{\mathbb{C}^{s_{i_2}}} \dots \otimes_{\mathbb{C}^{s_{i_k}}} \mathbb{C}$$

which is not isomorphic to B_x , where $l(x) < l(w)$.

Alternative approach: Soergel bimodules

$s_i = (i, i + 1)$ — simple reflection in S_n for $i = 1, 2, \dots, n - 1$

Fact. S_n is a **Coxeter group** with generators s_i

$l : S_n \rightarrow \mathbb{Z}$ — the **length** function

\mathbb{C}^{s_i} — the algebra of s_i -invariants in \mathbb{C}

Definition. [Soergel] Define inductively \mathbb{C} - \mathbb{C} -bimodules B_w , $w \in S_n$ as follows:

- ▶ $B_e := \mathbb{C}$
- ▶ For $w = s_{i_1} s_{i_2} \dots s_{i_k}$ reduced decomposition, the bimodule B_w is the unique indecomposable direct summand of

$$\mathbb{C} \otimes_{\mathbb{C}^{s_{i_1}}} \mathbb{C} \otimes_{\mathbb{C}^{s_{i_2}}} \dots \otimes_{\mathbb{C}^{s_{i_k}}} \mathbb{C}$$

which is not isomorphic to B_x , where $l(x) < l(w)$.

Alternative approach: Soergel bimodules

$s_i = (i, i + 1)$ — simple reflection in S_n for $i = 1, 2, \dots, n - 1$

Fact. S_n is a **Coxeter group** with generators s_i

$l : S_n \rightarrow \mathbb{Z}$ — the **length** function

\mathbb{C}^{s_i} — the algebra of s_i -invariants in \mathbb{C}

Definition. [Soergel] Define inductively **\mathbb{C} - \mathbb{C} -bimodules** B_w , $w \in S_n$ as follows:

- ▶ $B_e := \mathbb{C}$
- ▶ For $w = s_{i_1} s_{i_2} \dots s_{i_k}$ **reduced decomposition**, the bimodule B_w is the **unique indecomposable direct summand** of

$$\mathbb{C} \otimes_{\mathbb{C}^{s_{i_1}}} \mathbb{C} \otimes_{\mathbb{C}^{s_{i_2}}} \dots \otimes_{\mathbb{C}^{s_{i_k}}} \mathbb{C}$$

which is not isomorphic to B_x , where $l(x) < l(w)$.

Alternative approach: Soergel bimodules

$s_i = (i, i + 1)$ — simple reflection in S_n for $i = 1, 2, \dots, n - 1$

Fact. S_n is a Coxeter group with generators s_i

$l : S_n \rightarrow \mathbb{Z}$ — the length function

\mathbf{C}^{s_i} — the algebra of s_i -invariants in \mathbf{C}

Definition. [Soergel] Define inductively \mathbf{C} - \mathbf{C} -bimodules B_w , $w \in S_n$ as follows:

- ▶ $B_e := \mathbf{C}$
- ▶ For $w = s_{i_1} s_{i_2} \dots s_{i_k}$ reduced decomposition, the bimodule B_w is the unique indecomposable direct summand of

$$\mathbf{C} \otimes_{\mathbf{C}^{s_{i_1}}} \mathbf{C} \otimes_{\mathbf{C}^{s_{i_2}}} \dots \otimes_{\mathbf{C}^{s_{i_k}}} \mathbf{C}$$

which is not isomorphic to B_x , where $l(x) < l(w)$.

Alternative approach: Soergel bimodules

$s_i = (i, i + 1)$ — simple reflection in S_n for $i = 1, 2, \dots, n - 1$

Fact. S_n is a **Coxeter group** with generators s_i

$l : S_n \rightarrow \mathbb{Z}$ — the **length** function

\mathbf{C}^{s_i} — the algebra of s_i -invariants in \mathbf{C}

Definition. [Soergel] Define inductively **C-C**-bimodules B_w , $w \in S_n$ as follows:

- ▶ $B_e := \mathbf{C}$
- ▶ For $w = s_{i_1} s_{i_2} \dots s_{i_k}$ **reduced decomposition**, the bimodule B_w is the **unique indecomposable direct summand** of

$$\mathbf{C} \otimes_{\mathbf{C}^{s_{i_1}}} \mathbf{C} \otimes_{\mathbf{C}^{s_{i_2}}} \dots \otimes_{\mathbf{C}^{s_{i_k}}} \mathbf{C}$$

which is not isomorphic to B_x , where $l(x) < l(w)$.

Alternative approach: Soergel bimodules

$s_i = (i, i + 1)$ — simple reflection in S_n for $i = 1, 2, \dots, n - 1$

Fact. S_n is a **Coxeter group** with generators s_i

$l : S_n \rightarrow \mathbb{Z}$ — the **length** function

\mathbf{C}^{s_i} — the algebra of s_i -invariants in \mathbf{C}

Definition. [Soergel] Define inductively **C-C**-bimodules B_w , $w \in S_n$ as follows:

- ▶ $B_e := \mathbf{C}$
- ▶ For $w = s_{i_1} s_{i_2} \dots s_{i_k}$ **reduced decomposition**, the bimodule B_w is the **unique indecomposable direct summand** of

$$\mathbf{C} \otimes_{\mathbf{C}^{s_{i_1}}} \mathbf{C} \otimes_{\mathbf{C}^{s_{i_2}}} \dots \otimes_{\mathbf{C}^{s_{i_k}}} \mathbf{C}$$

which is not isomorphic to B_x , where $l(x) < l(w)$.

Alternative approach: Soergel bimodules

$s_i = (i, i + 1)$ — simple reflection in S_n for $i = 1, 2, \dots, n - 1$

Fact. S_n is a **Coxeter group** with generators s_i

$l : S_n \rightarrow \mathbb{Z}$ — the **length** function

\mathbf{C}^{s_i} — the algebra of s_i -invariants in \mathbf{C}

Definition. [Soergel] Define inductively **C-C**-bimodules B_w , $w \in S_n$ as follows:

- ▶ $B_e := \mathbf{C}$
- ▶ For $w = s_{i_1} s_{i_2} \dots s_{i_k}$ **reduced decomposition**, the bimodule B_w is the **unique** indecomposable direct summand of

$$\mathbf{C} \otimes_{\mathbf{C}^{s_{i_1}}} \mathbf{C} \otimes_{\mathbf{C}^{s_{i_2}}} \dots \otimes_{\mathbf{C}^{s_{i_k}}} \mathbf{C}$$

which is not isomorphic to B_x , where $l(x) < l(w)$.

Alternative approach: Soergel bimodules

$s_i = (i, i + 1)$ — simple reflection in S_n for $i = 1, 2, \dots, n - 1$

Fact. S_n is a **Coxeter group** with generators s_i

$l : S_n \rightarrow \mathbb{Z}$ — the **length** function

\mathbf{C}^{s_i} — the algebra of s_i -invariants in \mathbf{C}

Definition. [Soergel] Define inductively **C-C**-bimodules B_w , $w \in S_n$ as follows:

- ▶ $B_e := \mathbf{C}$
- ▶ For $w = s_{i_1} s_{i_2} \dots s_{i_k}$ **reduced decomposition**, the bimodule B_w is the **unique** indecomposable direct summand of

$$\mathbf{C} \otimes_{\mathbf{C}^{s_{i_1}}} \mathbf{C} \otimes_{\mathbf{C}^{s_{i_2}}} \dots \otimes_{\mathbf{C}^{s_{i_k}}} \mathbf{C}$$

which is not isomorphic to B_x , where $l(x) < l(w)$.

Alternative approach: combinatorial description of \mathcal{P}

B_w , $w \in S_n$ — Soergel bimodules

Note: For s simple reflection, $B_s \otimes_{\mathbb{C}} B_s \cong B_s \oplus B_s$.

Fact. For $x, y \in S_n$, each direct summand of the \mathbb{C} - \mathbb{C} -bimodule $B_x \otimes_{\mathbb{C}} B_y$ is isomorphic to B_z for some $z \in S_n$.

Definition. \mathcal{S} is the (additive) tensor category of Soergel bimodules.

Theorem. [Soergel's combinatorial description]

The categories \mathcal{P} and \mathcal{S} are equivalent as tensor categories.

Corollary. $\text{Gr}_{\oplus}[\mathcal{S}] \cong \mathbb{Z}[S_n]$.

Alternative approach: combinatorial description of \mathcal{P}

B_w , $w \in S_n$ — Soergel bimodules

Note: For s simple reflection, $B_s \otimes_{\mathbb{C}} B_s \cong B_s \oplus B_s$.

Fact. For $x, y \in S_n$, each direct summand of the \mathbb{C} - \mathbb{C} -bimodule $B_x \otimes_{\mathbb{C}} B_y$ is isomorphic to B_z for some $z \in S_n$.

Definition. \mathcal{S} is the (additive) tensor category of Soergel bimodules.

Theorem. [Soergel's combinatorial description]
The categories \mathcal{P} and \mathcal{S} are equivalent as tensor categories.

Corollary. $\text{Gr}_{\oplus}[\mathcal{S}] \cong \mathbb{Z}[S_n]$.

Alternative approach: combinatorial description of \mathcal{P}

B_w , $w \in S_n$ — Soergel bimodules

Note: For s simple reflection, $B_s \otimes_{\mathbb{C}} B_s \cong B_s \oplus B_s$.

Fact. For $x, y \in S_n$, each direct summand of the \mathbb{C} - \mathbb{C} -bimodule $B_x \otimes_{\mathbb{C}} B_y$ is isomorphic to B_z for some $z \in S_n$.

Definition. \mathcal{S} is the (additive) tensor category of Soergel bimodules.

Theorem. [Soergel's combinatorial description]
The categories \mathcal{P} and \mathcal{S} are equivalent as tensor categories.

Corollary. $\text{Gr}_{\oplus}[\mathcal{S}] \cong \mathbb{Z}[S_n]$.

Alternative approach: combinatorial description of \mathcal{P}

B_w , $w \in S_n$ — Soergel bimodules

Note: For s simple reflection, $B_s \otimes_{\mathbf{C}} B_s \cong B_s \oplus B_s$.

Fact. For $x, y \in S_n$, each direct summand of the \mathbf{C} - \mathbf{C} -bimodule $B_x \otimes_{\mathbf{C}} B_y$ is isomorphic to B_z for some $z \in S_n$.

Definition. \mathcal{S} is the (additive) tensor category of Soergel bimodules.

Theorem. [Soergel's combinatorial description]
The categories \mathcal{P} and \mathcal{S} are equivalent as tensor categories.

Corollary. $\text{Gr}_{\oplus}[\mathcal{S}] \cong \mathbb{Z}[S_n]$.

Alternative approach: combinatorial description of \mathcal{P}

B_w , $w \in S_n$ — Soergel bimodules

Note: For s simple reflection, $B_s \otimes_{\mathbf{C}} B_s \cong B_s \oplus B_s$.

Fact. For $x, y \in S_n$, each direct summand of the \mathbf{C} - \mathbf{C} -bimodule $B_x \otimes_{\mathbf{C}} B_y$ is isomorphic to B_z for some $z \in S_n$.

Definition. \mathcal{S} is the (additive) tensor category of Soergel bimodules.

Theorem. [Soergel's combinatorial description]
The categories \mathcal{P} and \mathcal{S} are equivalent as tensor categories.

Corollary. $\text{Gr}_{\oplus}[\mathcal{S}] \cong \mathbb{Z}[S_n]$.

Alternative approach: combinatorial description of \mathcal{P}

B_w , $w \in S_n$ — Soergel bimodules

Note: For s simple reflection, $B_s \otimes_{\mathbf{C}} B_s \cong B_s \oplus B_s$.

Fact. For $x, y \in S_n$, each direct summand of the \mathbf{C} - \mathbf{C} -bimodule $B_x \otimes_{\mathbf{C}} B_y$ is isomorphic to B_z for some $z \in S_n$.

Definition. \mathcal{S} is the (additive) tensor category of Soergel bimodules.

Theorem. [Soergel's combinatorial description]

The categories \mathcal{P} and \mathcal{S} are equivalent as tensor categories.

Corollary. $\text{Gr}_{\oplus}[\mathcal{S}] \cong \mathbb{Z}[S_n]$.

Alternative approach: combinatorial description of \mathcal{P}

B_w , $w \in S_n$ — Soergel bimodules

Note: For s simple reflection, $B_s \otimes_{\mathbf{C}} B_s \cong B_s \oplus B_s$.

Fact. For $x, y \in S_n$, each direct summand of the \mathbf{C} - \mathbf{C} -bimodule $B_x \otimes_{\mathbf{C}} B_y$ is isomorphic to B_z for some $z \in S_n$.

Definition. \mathcal{S} is the (additive) tensor category of Soergel bimodules.

Theorem. [Soergel's combinatorial description]

The categories \mathcal{P} and \mathcal{S} are equivalent as tensor categories.

Corollary. $\text{Gr}_{\oplus}[\mathcal{S}] \cong \mathbb{Z}[S_n]$.

Alternative approach: combinatorial description of \mathcal{P}

B_w , $w \in S_n$ — Soergel bimodules

Note: For s simple reflection, $B_s \otimes_{\mathbf{C}} B_s \cong B_s \oplus B_s$.

Fact. For $x, y \in S_n$, each direct summand of the \mathbf{C} - \mathbf{C} -bimodule $B_x \otimes_{\mathbf{C}} B_y$ is isomorphic to B_z for some $z \in S_n$.

Definition. \mathcal{S} is the (additive) tensor category of Soergel bimodules.

Theorem. [Soergel's combinatorial description]

The categories \mathcal{P} and \mathcal{S} are equivalent as tensor categories.

Corollary. $\text{Gr}_{\oplus}[\mathcal{S}] \cong \mathbb{Z}[S_n]$.

Categorification of permutation modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — composition of n

$\mathfrak{g}_\lambda \subset \mathfrak{g}$ — corresponding parabolic subalgebra.

W_λ — corresponding Young subgroup of S_n

${}_\lambda\text{Long}$ — longest representatives in $W_\lambda \setminus W$

\mathcal{X}_λ — Serre subcategory of \mathcal{O}_0 generated by $L(w)$, $w \notin {}_\lambda\text{Long}$

Fact: \mathcal{P} preserves \mathcal{X}_λ

Theorem. [M.-Stroppel] The induced action of \mathcal{P} on $\mathcal{O}_0/\mathcal{X}_\lambda$ categorifies the permutation module $\text{Ind}_{W_\lambda}^W \text{triv}$.

Categorification of permutation modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — composition of n

$\mathfrak{g}_\lambda \subset \mathfrak{g}$ — corresponding parabolic subalgebra.

W_λ — corresponding Young subgroup of S_n

${}_\lambda\text{Long}$ — longest representatives in $W_\lambda \setminus W$

\mathcal{X}_λ — Serre subcategory of \mathcal{O}_0 generated by $L(w)$, $w \notin {}_\lambda\text{Long}$

Fact: \mathcal{P} preserves \mathcal{X}_λ

Theorem. [M.-Stroppel] The induced action of \mathcal{P} on $\mathcal{O}_0/\mathcal{X}_\lambda$ categorifies the permutation module $\text{Ind}_{W_\lambda}^W \text{triv}$.

Categorification of permutation modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — composition of n

$\mathfrak{g}_\lambda \subset \mathfrak{g}$ — corresponding parabolic subalgebra.

W_λ — corresponding Young subgroup of S_n

${}_\lambda\text{Long}$ — longest representatives in $W_\lambda \setminus W$

\mathcal{X}_λ — Serre subcategory of \mathcal{O}_0 generated by $L(w)$, $w \notin {}_\lambda\text{Long}$

Fact: \mathcal{P} preserves \mathcal{X}_λ

Theorem. [M.-Stroppel] The induced action of \mathcal{P} on $\mathcal{O}_0/\mathcal{X}_\lambda$ categorifies the permutation module $\text{Ind}_{W_\lambda}^W \text{triv}$.

Categorification of permutation modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — composition of n

$\mathfrak{g}_\lambda \subset \mathfrak{g}$ — corresponding parabolic subalgebra.

W_λ — corresponding Young subgroup of S_n

${}_\lambda \text{Long}$ — longest representatives in $W_\lambda \backslash W$

\mathcal{X}_λ — Serre subcategory of \mathcal{O}_0 generated by $L(w)$, $w \notin {}_\lambda \text{Long}$

Fact: \mathcal{P} preserves \mathcal{X}_λ

Theorem. [M.-Stroppel] The induced action of \mathcal{P} on $\mathcal{O}_0/\mathcal{X}_\lambda$ categorifies the permutation module $\text{Ind}_{W_\lambda}^W \text{triv}$.

Categorification of permutation modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — composition of n

$\mathfrak{g}_\lambda \subset \mathfrak{g}$ — corresponding parabolic subalgebra.

W_λ — corresponding Young subgroup of S_n

${}_\lambda\text{Long}$ — longest representatives in $W_\lambda \backslash W$

\mathcal{X}_λ — Serre subcategory of \mathcal{O}_0 generated by $L(w)$, $w \notin {}_\lambda\text{Long}$

Fact: \mathcal{P} preserves \mathcal{X}_λ

Theorem. [M.-Stroppel] The induced action of \mathcal{P} on $\mathcal{O}_0/\mathcal{X}_\lambda$ categorifies the permutation module $\text{Ind}_{W_\lambda}^W \text{triv}$.

Categorification of permutation modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — composition of n

$\mathfrak{g}_\lambda \subset \mathfrak{g}$ — corresponding parabolic subalgebra.

W_λ — corresponding Young subgroup of S_n

${}_\lambda\text{Long}$ — longest representatives in $W_\lambda \setminus W$

\mathcal{X}_λ — Serre subcategory of \mathcal{O}_0 generated by $L(w)$, $w \notin {}_\lambda\text{Long}$

Fact: \mathcal{P} preserves \mathcal{X}_λ

Theorem. [M.-Stroppel] The induced action of \mathcal{P} on $\mathcal{O}_0/\mathcal{X}_\lambda$ categorifies the permutation module $\text{Ind}_{W_\lambda}^W \text{triv}$.

Categorification of permutation modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — composition of n

$\mathfrak{g}_\lambda \subset \mathfrak{g}$ — corresponding parabolic subalgebra.

W_λ — corresponding Young subgroup of S_n

${}_\lambda\text{Long}$ — longest representatives in $W_\lambda \backslash W$

\mathcal{X}_λ — Serre subcategory of \mathcal{O}_0 generated by $L(w)$, $w \notin {}_\lambda\text{Long}$

Fact: \mathcal{P} preserves \mathcal{X}_λ

Theorem. [M.-Stroppel] The induced action of \mathcal{P} on $\mathcal{O}_0/\mathcal{X}_\lambda$ categorifies the permutation module $\text{Ind}_{W_\lambda}^W \text{triv}$.

Categorification of permutation modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — composition of n

$\mathfrak{g}_\lambda \subset \mathfrak{g}$ — corresponding parabolic subalgebra.

W_λ — corresponding Young subgroup of S_n

${}_\lambda \text{Long}$ — longest representatives in $W_\lambda \setminus W$

\mathcal{X}_λ — Serre subcategory of \mathcal{O}_0 generated by $L(w)$, $w \notin {}_\lambda \text{Long}$

Fact: \mathcal{P} preserves \mathcal{X}_λ

Theorem. [M.-Stroppel] The induced action of \mathcal{P} on $\mathcal{O}_0/\mathcal{X}_\lambda$ categorifies the permutation module $\text{Ind}_{W_\lambda}^W \text{triv.}$

Categorification of permutation modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — composition of n

$\mathfrak{g}_\lambda \subset \mathfrak{g}$ — corresponding parabolic subalgebra.

W_λ — corresponding Young subgroup of S_n

${}_\lambda \text{Long}$ — longest representatives in $W_\lambda \setminus W$

\mathcal{X}_λ — Serre subcategory of \mathcal{O}_0 generated by $L(w)$, $w \notin {}_\lambda \text{Long}$

Fact: \mathcal{P} preserves \mathcal{X}_λ

Theorem. [M.-Stroppel] The induced action of \mathcal{P} on $\mathcal{O}_0/\mathcal{X}_\lambda$ categorifies the permutation module $\text{Ind}_{W_\lambda}^W \text{triv.}$

Categorification of induced sign modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — composition of n

$\mathfrak{g}_\lambda \subset \mathfrak{g}$ — corresponding parabolic subalgebra.

W_λ — corresponding Young subgroup of S_n

Short_λ — shortest representatives in W/W_λ

\mathcal{Y}_λ — Serre subcategory of \mathcal{O}_0 generated by $L(w)$, $w \in \text{Short}_\lambda$,
(Rocha-Caridi's parabolic category \mathcal{O})

Fact: \mathcal{P} preserves \mathcal{Y}_λ

Theorem. [Soergel]

The action of \mathcal{P} on \mathcal{Y}_λ categorifies the induced sign module $\text{Ind}_{W_\lambda}^W \text{sign}$.

Categorification of induced sign modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **composition** of n

$\mathfrak{g}_\lambda \subset \mathfrak{g}$ — corresponding parabolic subalgebra.

W_λ — corresponding Young subgroup of S_n

Short_λ — shortest representatives in W/W_λ

\mathcal{Y}_λ — Serre subcategory of \mathcal{O}_0 generated by $L(w)$, $w \in \text{Short}_\lambda$,
(Rocha-Caridi's parabolic category \mathcal{O})

Fact: \mathcal{P} preserves \mathcal{Y}_λ

Theorem. [Soergel]

The action of \mathcal{P} on \mathcal{Y}_λ categorifies the induced sign module $\text{Ind}_{W_\lambda}^W \text{sign}$.

Categorification of induced sign modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **composition** of n

$\mathfrak{g}_\lambda \subset \mathfrak{g}$ — corresponding **parabolic** subalgebra.

W_λ — corresponding Young subgroup of S_n

Short_λ — shortest representatives in W/W_λ

\mathcal{Y}_λ — Serre subcategory of \mathcal{O}_0 generated by $L(w)$, $w \in \text{Short}_\lambda$,
(Rocha-Caridi's parabolic category \mathcal{O})

Fact: \mathcal{P} preserves \mathcal{Y}_λ

Theorem. [Soergel]

The action of \mathcal{P} on \mathcal{Y}_λ categorifies the induced sign module $\text{Ind}_{W_\lambda}^W \text{sign}$.

Categorification of induced sign modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **composition** of n

$\mathfrak{g}_\lambda \subset \mathfrak{g}$ — corresponding **parabolic** subalgebra.

W_λ — corresponding **Young subgroup** of S_n

Short_λ — **shortest** representatives in W/W_λ

\mathcal{Y}_λ — **Serre subcategory** of \mathcal{O}_0 generated by $L(w)$, $w \in \text{Short}_\lambda$,
(Rocha-Caridi's **parabolic category** \mathcal{O})

Fact: \mathcal{P} preserves \mathcal{Y}_λ

Theorem. [Soergel]

The action of \mathcal{P} on \mathcal{Y}_λ categorifies the induced sign module $\text{Ind}_{W_\lambda}^W \text{sign}$.

Categorification of induced sign modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **composition** of n

$\mathfrak{g}_\lambda \subset \mathfrak{g}$ — corresponding **parabolic** subalgebra.

W_λ — corresponding **Young subgroup** of S_n

Short_λ — **shortest representatives** in W/W_λ

\mathcal{Y}_λ — Serre subcategory of \mathcal{O}_0 generated by $L(w)$, $w \in \text{Short}_\lambda$,
(Rocha-Caridi's parabolic category \mathcal{O})

Fact: \mathcal{P} preserves \mathcal{Y}_λ

Theorem. [Soergel]

The action of \mathcal{P} on \mathcal{Y}_λ categorifies the induced sign module $\text{Ind}_{W_\lambda}^W \text{sign}$.

Categorification of induced sign modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **composition** of n

$\mathfrak{g}_\lambda \subset \mathfrak{g}$ — corresponding **parabolic** subalgebra.

W_λ — corresponding **Young subgroup** of S_n

Short_λ — **shortest representatives** in W/W_λ

\mathcal{Y}_λ — **Serre subcategory** of \mathcal{O}_0 generated by $L(w)$, $w \in \text{Short}_\lambda$,
(Rocha-Caridi's **parabolic** category \mathcal{O})

Fact: \mathcal{P} preserves \mathcal{Y}_λ

Theorem. [Soergel]

The action of \mathcal{P} on \mathcal{Y}_λ categorifies the induced sign module $\text{Ind}_{W_\lambda}^W \text{sign}$.

Categorification of induced sign modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **composition** of n

$\mathfrak{g}_\lambda \subset \mathfrak{g}$ — corresponding **parabolic** subalgebra.

W_λ — corresponding **Young subgroup** of S_n

Short_λ — **shortest representatives** in W/W_λ

\mathcal{Y}_λ — **Serre subcategory** of \mathcal{O}_0 generated by $L(w)$, $w \in \text{Short}_\lambda$,
(Rocha-Caridi's **parabolic** category \mathcal{O})

Fact: \mathcal{P} preserves \mathcal{Y}_λ

Theorem. [Soergel]

The action of \mathcal{P} on \mathcal{Y}_λ categorifies the induced sign module $\text{Ind}_{W_\lambda}^W \text{sign}$.

Categorification of induced sign modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **composition** of n

$\mathfrak{g}_\lambda \subset \mathfrak{g}$ — corresponding **parabolic** subalgebra.

W_λ — corresponding **Young subgroup** of S_n

Short_λ — **shortest representatives** in W/W_λ

\mathcal{Y}_λ — **Serre subcategory** of \mathcal{O}_0 generated by $L(w)$, $w \in \text{Short}_\lambda$,
(Rocha-Caridi's **parabolic** category \mathcal{O})

Fact: \mathcal{P} preserves \mathcal{Y}_λ

Theorem. [Soergel]

The action of \mathcal{P} on \mathcal{Y}_λ categorifies the **induced sign module** $\text{Ind}_{W_\lambda}^W \text{sign}$.

Categorification of induced sign modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **composition** of n

$\mathfrak{g}_\lambda \subset \mathfrak{g}$ — corresponding **parabolic** subalgebra.

W_λ — corresponding **Young subgroup** of S_n

Short_λ — **shortest representatives** in W/W_λ

\mathcal{Y}_λ — **Serre subcategory** of \mathcal{O}_0 generated by $L(w)$, $w \in \text{Short}_\lambda$,
(Rocha-Caridi's **parabolic** category \mathcal{O})

Fact: \mathcal{P} preserves \mathcal{Y}_λ

Theorem. [Soergel]

The action of \mathcal{P} on \mathcal{Y}_λ categorifies the **induced sign module** $\text{Ind}_{W_\lambda}^W \text{sign}$.

Categorification of Specht modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — partition of n

w_0 — longest element in W

w_0^λ — longest element in W_λ

\mathcal{Y}'_λ — Serre subcategory of \mathcal{Y}_λ generated by $L(w)$, $w \in \text{Short}_\lambda$, such that $\text{GKdim}(L(w)) < \text{GKdim}(L(w_0^\lambda w_0))$

Fact: \mathcal{P} preserves \mathcal{Y}'_λ

Theorem. [Khovanov-M.-Stroppel]

The induced action of \mathcal{P} on $\mathcal{Y}_\lambda/\mathcal{Y}'_\lambda$ categorifies \mathcal{S}^{λ^t} .

Categorification of Specht modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **partition** of n

w_0 — **longest** element in W

w_0^λ — **longest** element in W_λ

\mathcal{Y}'_λ — Serre subcategory of \mathcal{Y}_λ generated by $L(w)$, $w \in \text{Short}_\lambda$, such that $\text{GKdim}(L(w)) < \text{GKdim}(L(w_0^\lambda w_0))$

Fact: \mathcal{P} preserves \mathcal{Y}'_λ

Theorem. [Khovanov-M.-Stroppel]

The induced action of \mathcal{P} on $\mathcal{Y}_\lambda/\mathcal{Y}'_\lambda$ categorifies \mathcal{S}^{λ^*} .

Categorification of Specht modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **partition** of n

w_0 — **longest** element in W

w_0^λ — **longest** element in W_λ

\mathcal{Y}'_λ — Serre subcategory of \mathcal{Y}_λ generated by $L(w)$, $w \in \text{Short}_\lambda$, such that $\text{GKdim}(L(w)) < \text{GKdim}(L(w_0^\lambda w_0))$

Fact: \mathcal{P} preserves \mathcal{Y}'_λ

Theorem. [Khovanov-M.-Stroppel]

The induced action of \mathcal{P} on $\mathcal{Y}_\lambda/\mathcal{Y}'_\lambda$ categorifies \mathcal{S}^{λ^*} .

Categorification of Specht modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **partition** of n

w_0 — **longest** element in W

w_0^λ — **longest** element in W_λ

\mathcal{Y}'_λ — Serre subcategory of \mathcal{Y}_λ generated by $L(w)$, $w \in \text{Short}_\lambda$, such that $\text{GKdim}(L(w)) < \text{GKdim}(L(w_0^\lambda w_0))$

Fact: \mathcal{P} preserves \mathcal{Y}'_λ

Theorem. [Khovanov-M.-Stroppel]

The induced action of \mathcal{P} on $\mathcal{Y}_\lambda/\mathcal{Y}'_\lambda$ categorifies \mathcal{S}^{λ^*} .

Categorification of Specht modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **partition** of n

w_0 — **longest** element in W

w_0^λ — **longest** element in W_λ

\mathcal{Y}'_λ — **Serre subcategory** of \mathcal{Y}_λ generated by $L(w)$, $w \in \text{Short}_\lambda$, such that $\text{GKdim}(L(w)) < \text{GKdim}(L(w_0^\lambda w_0))$

Fact: \mathcal{P} preserves \mathcal{Y}'_λ

Theorem. [Khovanov-M.-Stroppel]

The induced action of \mathcal{P} on $\mathcal{Y}_\lambda/\mathcal{Y}'_\lambda$ categorifies \mathcal{S}^{λ^t} .

Categorification of Specht modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **partition** of n

w_0 — **longest** element in W

w_0^λ — **longest** element in W_λ

\mathcal{Y}'_λ — **Serre subcategory** of \mathcal{Y}_λ generated by $L(w)$, $w \in \text{Short}_\lambda$, such that $\text{GKdim}(L(w)) < \text{GKdim}(L(w_0^\lambda w_0))$

Fact: \mathcal{P} preserves \mathcal{Y}'_λ

Theorem. [Khovanov-M.-Stroppel]

The induced action of \mathcal{P} on $\mathcal{Y}_\lambda/\mathcal{Y}'_\lambda$ categorifies \mathcal{S}^{λ^t} .

Categorification of Specht modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **partition** of n

w_0 — **longest** element in W

w_0^λ — **longest** element in W_λ

\mathcal{Y}'_λ — **Serre subcategory** of \mathcal{Y}_λ generated by $L(w)$, $w \in \text{Short}_\lambda$, such that $\text{GKdim}(L(w)) < \text{GKdim}(L(w_0^\lambda w_0))$

Fact: \mathcal{P} preserves \mathcal{Y}'_λ

Theorem. [Khovanov-M.-Stroppel]

The induced action of \mathcal{P} on $\mathcal{Y}_\lambda/\mathcal{Y}'_\lambda$ categorifies \mathcal{S}^{λ^t} .

Categorification of Specht modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **partition** of n

w_0 — **longest** element in W

w_0^λ — **longest** element in W_λ

\mathcal{Y}'_λ — **Serre subcategory** of \mathcal{Y}_λ generated by $L(w)$, $w \in \text{Short}_\lambda$, such that $\text{GKdim}(L(w)) < \text{GKdim}(L(w_0^\lambda w_0))$

Fact: \mathcal{P} preserves \mathcal{Y}'_λ

Theorem. [Khovanov-M.-Stroppel]

The induced action of \mathcal{P} on $\mathcal{Y}_\lambda/\mathcal{Y}'_\lambda$ categorifies \mathcal{S}^{λ^t} .

Alternative categorification of Specht modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — partition of n

w_0^λ — longest element in W_λ

\mathcal{X}'_λ — Serre subcategory of $\mathcal{O}_0/\mathcal{X}_\lambda$ generated by $L(w)$, $w \in {}_\lambda\text{Long}$, such that $\text{GKdim}(L(w)) = \text{GKdim}(L(w_0^\lambda))$

Fact: \mathcal{P} preserves \mathcal{X}'_λ

Theorem. [M.-Stroppel]

The action of \mathcal{P} on \mathcal{X}'_λ categorifies S^λ .

Alternative categorification of Specht modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **partition** of n

w_0^λ — longest element in W_λ

\mathcal{X}'_λ — Serre subcategory of $\mathcal{O}_0/\mathcal{X}_\lambda$ generated by $L(w)$, $w \in {}_\lambda\text{Long}$, such that $\text{GKdim}(L(w)) = \text{GKdim}(L(w_0^\lambda))$

Fact: \mathcal{P} preserves \mathcal{X}'_λ

Theorem. [M.-Stroppel]

The action of \mathcal{P} on \mathcal{X}'_λ categorifies S^λ .

Alternative categorification of Specht modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **partition** of n

w_0^λ — **longest** element in W_λ

\mathcal{X}'_λ — Serre subcategory of $\mathcal{O}_0/\mathcal{X}_\lambda$ generated by $L(w)$, $w \in {}_\lambda\text{Long}$, such that $\text{GKdim}(L(w)) = \text{GKdim}(L(w_0^\lambda))$

Fact: \mathcal{P} preserves \mathcal{X}'_λ

Theorem. [M.-Stroppel]

The action of \mathcal{P} on \mathcal{X}'_λ categorifies S^λ .

Alternative categorification of Specht modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **partition** of n

w_0^λ — **longest** element in W_λ

\mathcal{X}'_λ — **Serre subcategory** of $\mathcal{O}_0/\mathcal{X}_\lambda$ generated by $L(w)$, $w \in {}_\lambda\text{Long}$, such that $\text{GKdim}(L(w)) = \text{GKdim}(L(w_0^\lambda))$

Fact: \mathcal{P} preserves \mathcal{X}'_λ

Theorem. [M.-Stroppel]

The action of \mathcal{P} on \mathcal{X}'_λ categorifies \mathcal{S}^λ .

Alternative categorification of Specht modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **partition** of n

w_0^λ — **longest** element in W_λ

\mathcal{X}'_λ — **Serre subcategory** of $\mathcal{O}_0/\mathcal{X}_\lambda$ generated by $L(w)$, $w \in {}_\lambda\text{Long}$, such that $\text{GKdim}(L(w)) = \text{GKdim}(L(w_0^\lambda))$

Fact: \mathcal{P} preserves \mathcal{X}'_λ

Theorem. [M.-Stroppel]

The action of \mathcal{P} on \mathcal{X}'_λ categorifies \mathcal{S}^λ .

Alternative categorification of Specht modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **partition** of n

w_0^λ — **longest** element in W_λ

\mathcal{X}'_λ — **Serre subcategory** of $\mathcal{O}_0/\mathcal{X}_\lambda$ generated by $L(w)$, $w \in {}_\lambda\text{Long}$, such that $\text{GKdim}(L(w)) = \text{GKdim}(L(w_0^\lambda))$

Fact: \mathcal{P} preserves \mathcal{X}'_λ

Theorem. [M.-Stroppel]

The action of \mathcal{P} on \mathcal{X}'_λ categorifies \mathcal{S}^λ .

Alternative categorification of Specht modules

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — partition of n

w_0^λ — longest element in W_λ

\mathcal{X}'_λ — Serre subcategory of $\mathcal{O}_0/\mathcal{X}_\lambda$ generated by $L(w)$, $w \in {}_\lambda\text{Long}$, such that $\text{GKdim}(L(w)) = \text{GKdim}(L(w_0^\lambda))$

Fact: \mathcal{P} preserves \mathcal{X}'_λ

Theorem. [M.-Stroppel]

The action of \mathcal{P} on \mathcal{X}'_λ categorifies \mathcal{S}^λ .

Comparison

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — partition of n

Question. Are categorifications of \mathcal{S}^λ via \mathcal{X}'_λ and $\mathcal{Y}_{\lambda^*}/\mathcal{Y}'_{\lambda^*}$ equivalent?

Need: An equivalence between these two categories which naturally commutes with the action of \mathcal{P}

Theorem. [M.-Stroppel]

These two categorifications are indeed equivalent (using derived completion functors).

Theorem. [M.-Miemietz]

Simple transitive categorification of a Specht module (using \mathcal{P}) is **unique** up to equivalence.

Note. The latter works only in type A .

Comparison

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **partition** of n

Question. Are categorifications of \mathcal{S}^λ via \mathcal{X}'_λ and $\mathcal{Y}_{\lambda^*}/\mathcal{Y}'_{\lambda^*}$ equivalent?

Need: An equivalence between these two categories which **naturally** commutes with the action of \mathcal{P}

Theorem. [M.-Stroppel]

These two categorifications are indeed equivalent (using derived completion functors).

Theorem. [M.-Miemietz]

Simple transitive categorification of a Specht module (using \mathcal{P}) is **unique** up to equivalence.

Note. The latter works only in type A .

Comparison

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **partition** of n

Question. Are categorifications of \mathcal{S}^λ via \mathcal{X}'_λ and $\mathcal{Y}_{\lambda^\dagger}/\mathcal{Y}'_{\lambda^\dagger}$ **equivalent**?

Need: An equivalence between these two categories which **naturally** commutes with the action of \mathcal{P}

Theorem. [M.-Stroppel]

These two categorifications are indeed equivalent (using derived completion functors).

Theorem. [M.-Miemietz]

Simple transitive categorification of a Specht module (using \mathcal{P}) is **unique** up to equivalence.

Note. The latter works only in type A .

Comparison

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **partition** of n

Question. Are categorifications of \mathcal{S}^λ via \mathcal{X}'_λ and $\mathcal{Y}_{\lambda^\dagger}/\mathcal{Y}'_{\lambda^\dagger}$ **equivalent**?

Need: An equivalence between these two categories which **naturally** commutes with the action of \mathcal{P}

Theorem. [M.-Stroppel]

These two categorifications are indeed equivalent (using derived completion functors).

Theorem. [M.-Miemietz]

Simple transitive categorification of a Specht module (using \mathcal{P}) is **unique** up to equivalence.

Note. The latter works only in type A .

Comparison

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **partition** of n

Question. Are categorifications of \mathcal{S}^λ via \mathcal{X}'_λ and $\mathcal{Y}_{\lambda^\dagger}/\mathcal{Y}'_{\lambda^\dagger}$ **equivalent**?

Need: An equivalence between these two categories which **naturally** commutes with the action of \mathcal{P}

Theorem. [M.-Stroppel]

These two categorifications are indeed equivalent (using derived completion functors).

Theorem. [M.-Miemietz]

Simple transitive categorification of a Specht module (using \mathcal{P}) is **unique** up to equivalence.

Note. The latter works only in type A .

Comparison

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **partition** of n

Question. Are categorifications of \mathcal{S}^λ via \mathcal{X}'_λ and $\mathcal{Y}_{\lambda^\dagger}/\mathcal{Y}'_{\lambda^\dagger}$ **equivalent**?

Need: An equivalence between these two categories which **naturally** commutes with the action of \mathcal{P}

Theorem. [M.-Stroppel]

These two categorifications are indeed equivalent (using derived completion functors).

Theorem. [M.-Miemietz]

Simple transitive categorification of a Specht module (using \mathcal{P}) is **unique** up to equivalence.

Note. The latter works only in type A .

Comparison

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **partition** of n

Question. Are categorifications of \mathcal{S}^λ via \mathcal{X}'_λ and $\mathcal{Y}_{\lambda^\dagger}/\mathcal{Y}'_{\lambda^\dagger}$ **equivalent**?

Need: An equivalence between these two categories which **naturally** commutes with the action of \mathcal{P}

Theorem. [M.-Stroppel]

These two categorifications are indeed equivalent (using derived completion functors).

Theorem. [M.-Miemietz]

Simple transitive categorification of a Specht module (using \mathcal{P}) is **unique** up to equivalence.

Note. The latter works only in type A .

Comparison

$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ — **partition** of n

Question. Are categorifications of \mathcal{S}^λ via \mathcal{X}'_λ and $\mathcal{Y}_{\lambda^\dagger}/\mathcal{Y}'_{\lambda^\dagger}$ **equivalent**?

Need: An equivalence between these two categories which **naturally** commutes with the action of \mathcal{P}

Theorem. [M.-Stroppel]

These two categorifications are indeed equivalent (using derived completion functors).

Theorem. [M.-Miemietz]

Simple transitive categorification of a Specht module (using \mathcal{P}) is **unique** up to equivalence.

Note. The latter works only in type A .

Further categorifications using \mathcal{P}

Kazhdan-Lusztig's cell modules for S_n

Note. This requires a generalization of parabolic category \mathcal{O}

Induced cell modules $\text{Ind}_{W_\lambda}^W \text{cell}$

Note. Uses combinatorially defined subquotients of \mathcal{O}_0

Note. Permutation and induced sign modules are special cases

Wedderburn's basis for $\mathbb{Q}[S_n]$

Schur-Weyl duality

Further categorifications using \mathcal{P}

Kazhdan-Lusztig's **cell modules** for S_n

Note. This requires a generalization of parabolic category \mathcal{O}

Induced cell modules $\text{Ind}_{W_\lambda}^W \text{cell}$

Note. Uses combinatorially defined subquotients of \mathcal{O}_0

Note. Permutation and induced sign modules are special cases

Wedderburn's basis for $\mathbb{Q}[S_n]$

Schur-Weyl duality

Further categorifications using \mathcal{P}

Kazhdan-Lusztig's **cell modules** for S_n

Note. This requires a generalization of **parabolic** category \mathcal{O}

Induced cell modules $\text{Ind}_{W_\lambda}^W \text{cell}$

Note. Uses combinatorially defined **subquotients** of \mathcal{O}_0

Note. Permutation and induced sign modules are **special cases**

Wedderburn's basis for $\mathbb{Q}[S_n]$

Schur-Weyl duality

Further categorifications using \mathcal{P}

Kazhdan-Lusztig's **cell modules** for S_n

Note. This requires a generalization of **parabolic** category \mathcal{O}

Induced cell modules $\text{Ind}_{W_\lambda}^W \text{cell}$

Note. Uses combinatorially defined **subquotients** of \mathcal{O}_0

Note. Permutation and induced sign modules are **special cases**

Wedderburn's basis for $\mathbb{Q}[S_n]$

Schur-Weyl duality

Further categorifications using \mathcal{P}

Kazhdan-Lusztig's **cell modules** for S_n

Note. This requires a generalization of **parabolic** category \mathcal{O}

Induced cell modules $\text{Ind}_{W_\lambda}^W \text{cell}$

Note. Uses combinatorially defined **subquotients** of \mathcal{O}_0

Note. Permutation and induced sign modules are **special cases**

Wedderburn's basis for $\mathbb{Q}[S_n]$

Schur-Weyl duality

Further categorifications using \mathcal{P}

Kazhdan-Lusztig's **cell modules** for S_n

Note. This requires a generalization of **parabolic** category \mathcal{O}

Induced cell modules $\text{Ind}_{W_\lambda}^W \text{cell}$

Note. Uses combinatorially defined **subquotients** of \mathcal{O}_0

Note. Permutation and induced sign modules are **special cases**

Wedderburn's basis for $\mathbb{Q}[S_n]$

Schur-Weyl duality

Further categorifications using \mathcal{P}

Kazhdan-Lusztig's **cell modules** for S_n

Note. This requires a generalization of **parabolic** category \mathcal{O}

Induced cell modules $\text{Ind}_{W_\lambda}^W \text{cell}$

Note. Uses combinatorially defined **subquotients** of \mathcal{O}_0

Note. Permutation and induced sign modules are **special cases**

Wedderburn's basis for $\mathbb{Q}[S_n]$

Schur-Weyl duality

Further categorifications using \mathcal{P}

Kazhdan-Lusztig's **cell modules** for S_n

Note. This requires a generalization of **parabolic** category \mathcal{O}

Induced cell modules $\text{Ind}_{W_\lambda}^W \text{cell}$

Note. Uses combinatorially defined **subquotients** of \mathcal{O}_0

Note. Permutation and induced sign modules are **special cases**

Wedderburn's basis for $\mathbb{Q}[S_n]$

Schur-Weyl duality

Further categorifications using \mathcal{P}

Kazhdan-Lusztig's **cell modules** for S_n

Note. This requires a generalization of **parabolic** category \mathcal{O}

Induced cell modules $\text{Ind}_{W_\lambda}^W \text{cell}$

Note. Uses combinatorially defined **subquotients** of \mathcal{O}_0

Note. Permutation and induced sign modules are **special cases**

Wedderburn's basis for $\mathbb{Q}[S_n]$

Schur-Weyl duality

Summary of bonuses provided by categorification

Various bases in $\mathbb{Z}[S_n]$ and in other categorified modules

Different submodules in $\mathbb{Z}[S_n]$

Filtration of $\mathbb{Z}[S_n]$ using Gelfand-Kirillov dimension of simples in \mathcal{O}

Uniqueness of categorification allows to compare different categories of \mathfrak{g} -modules

Explicit knowledge of categorification provides explicit (partial) information on the underlying category (e.g. quiver and relations for blocks of \mathcal{O})

Summary of bonuses provided by categorification

Various **bases** in $\mathbb{Z}[S_n]$ and in other categorified modules

Different **submodules** in $\mathbb{Z}[S_n]$

Filtration of $\mathbb{Z}[S_n]$ using Gelfand-Kirillov dimension of simples in \mathcal{O}

Uniqueness of categorification allows to **compare** different categories of \mathfrak{g} -modules

Explicit knowledge of categorification provides **explicit** (partial) information on the underlying category (e.g. quiver and relations for blocks of \mathcal{O})

Summary of bonuses provided by categorification

Various **bases** in $\mathbb{Z}[S_n]$ and in other categorified modules

Different **submodules** in $\mathbb{Z}[S_n]$

Filtration of $\mathbb{Z}[S_n]$ using Gelfand-Kirillov dimension of simples in \mathcal{O}

Uniqueness of categorification allows to **compare** different categories of \mathfrak{g} -modules

Explicit knowledge of categorification provides **explicit** (partial) information on the underlying category (e.g. quiver and relations for blocks of \mathcal{O})

Summary of bonuses provided by categorification

Various **bases** in $\mathbb{Z}[S_n]$ and in other categorified modules

Different **submodules** in $\mathbb{Z}[S_n]$

Filtration of $\mathbb{Z}[S_n]$ using Gelfand-Kirillov dimension of simples in \mathcal{O}

Uniqueness of categorification allows to **compare** different categories of \mathfrak{g} -modules

Explicit knowledge of categorification provides **explicit** (partial) information on the underlying category (e.g. quiver and relations for blocks of \mathcal{O})

Summary of bonuses provided by categorification

Various **bases** in $\mathbb{Z}[S_n]$ and in other categorified modules

Different **submodules** in $\mathbb{Z}[S_n]$

Filtration of $\mathbb{Z}[S_n]$ using Gelfand-Kirillov dimension of simples in \mathcal{O}

Uniqueness of categorification allows to **compare** different categories of \mathfrak{g} -modules

Explicit knowledge of categorification provides **explicit** (partial) information on the underlying category (e.g. quiver and relations for blocks of \mathcal{O})

Summary of bonuses provided by categorification

Various **bases** in $\mathbb{Z}[S_n]$ and in other categorified modules

Different **submodules** in $\mathbb{Z}[S_n]$

Filtration of $\mathbb{Z}[S_n]$ using Gelfand-Kirillov dimension of simples in \mathcal{O}

Uniqueness of categorification allows to **compare** different categories of \mathfrak{g} -modules

Explicit knowledge of categorification provides **explicit** (partial) information on the underlying category (e.g. quiver and relations for blocks of \mathcal{O})

Summary of bonuses provided by categorification

Various **bases** in $\mathbb{Z}[S_n]$ and in other categorified modules

Different **submodules** in $\mathbb{Z}[S_n]$

Filtration of $\mathbb{Z}[S_n]$ using Gelfand-Kirillov dimension of simples in \mathcal{O}

Uniqueness of categorification allows to **compare** different categories of \mathfrak{g} -modules

Explicit knowledge of categorification provides **explicit** (partial) information on the underlying category (e.g. quiver and relations for blocks of \mathcal{O})

THANK YOU!!!