

Algebraic categorification and its applications, I

Volodymyr Mazorchuk
(Uppsala University)

Winter School “Geometry and physics”
January 17-24, 2015, Srní, Czech Republic

Categorification in short

“Definition”

Roughly speaking, **categorification** means an “upgrade” from set theory to category theory, in particular:

sets are upgraded to **categories**

functions are upgraded to **functors**

equalities are upgraded to **isomorphisms**

Categorification in short

“Definition”

Roughly speaking, **categorification** means an “upgrade” from set theory to category theory, in particular:

sets are upgraded to **categories**

functions are upgraded to **functors**

equalities are upgraded to **isomorphisms**

Categorification in short

“Definition”

Roughly speaking, **categorification** means an “upgrade” from set theory to category theory, in particular:

sets are upgraded to **categories**

functions are upgraded to **functors**

equalities are upgraded to **isomorphisms**

Categorification in short

“Definition”

Roughly speaking, **categorification** means an “upgrade” from set theory to category theory, in particular:

sets are upgraded to **categories**

functions are upgraded to **functors**

equalities are upgraded to **isomorphisms**

Categorification in short

“Definition”

Roughly speaking, **categorification** means an “upgrade” from set theory to category theory, in particular:

sets are upgraded to **categories**

functions are upgraded to **functors**

equalities are upgraded to **isomorphisms**

Categorification in short

“Definition”

Roughly speaking, **categorification** means an “upgrade” from set theory to category theory, in particular:

sets are upgraded to **categories**

functions are upgraded to **functors**

equalities are upgraded to **isomorphisms**

Categorification in short

“Definition”

Roughly speaking, **categorification** means an “upgrade” from set theory to category theory, in particular:

sets are upgraded to **categories**

functions are upgraded to **functors**

equalities are upgraded to **isomorphisms**

Motivation

Question: Why do we need categorification?

Answer: Categories have more **structure** than sets.

This can be used to get new useful information about objects we study.

Motivation

Question: Why do we need categorification?

Answer: Categories have more **structure** than sets.

This can be used to get new useful information about objects we study.

Motivation

Question: Why do we need categorification?

Answer: Categories have more **structure** than sets.

This can be used to get new useful information about objects we study.

Motivation

Question: Why do we need categorification?

Answer: Categories have more **structure** than sets.

This can be used to get new useful information about objects we study.

Motivation

Question: Why do we need categorification?

Answer: Categories have more **structure** than sets.

This can be used to get new useful information about objects we study.

Example: Khovanov homology — links and crossings

L — diagram of an oriented link

n_+ — number of right crossings

n_- — number of left crossings



right crossing



left crossing

Example: Khovanov homology — links and crossings

L — diagram of an oriented link

n_+ — number of right crossings

n_- — number of left crossings



Example: Khovanov homology — links and crossings

L — diagram of an oriented link

n_+ — number of right crossings

n_- — number of left crossings



Example: Khovanov homology — links and crossings

L — diagram of an oriented link

n_+ — number of right crossings

n_- — number of left crossings



Example: Khovanov homology — links and crossings

L — diagram of an oriented link

n_+ — number of right crossings

n_- — number of left crossings



right crossing



left crossing

Example: Khovanov homology — links and crossings

L — diagram of an oriented link

n_+ — number of right crossings

n_- — number of left crossings



right crossing



left crossing

Example: Khovanov homology — Kauffman bracket

Definition. The **Kauffman bracket** $\{L\} \in \mathbb{Z}[v, v^{-1}]$ of L is defined via the following rule:

$$\left\{ \begin{array}{c} \diagdown \\ \diagup \end{array} \right\} = \left\{ \begin{array}{c} \frown \\ \smile \end{array} \right\} - v \left\{ \begin{array}{c} \text{)} \\ \text{(} \end{array} \right\}$$

together with $\{\bigcirc L\} = (v + v^{-1})\{L\}$

and normalized by the conditions $\{\emptyset\} = 1$.

Example: Khovanov homology — Kauffman bracket

Definition. The **Kauffman bracket** $\{L\} \in \mathbb{Z}[v, v^{-1}]$ of L is defined via the following rule:

$$\left\{ \begin{array}{c} \diagdown \\ \diagup \end{array} \right\} = \left\{ \begin{array}{c} \text{---} \\ \text{---} \end{array} \right\} - v \left\{ \begin{array}{c} \text{---} \\ \text{---} \end{array} \right\}$$

together with $\{\bigcirc L\} = (v + v^{-1})\{L\}$

and normalized by the conditions $\{\emptyset\} = 1$.

Example: Khovanov homology — Kauffman bracket

Definition. The **Kauffman bracket** $\{L\} \in \mathbb{Z}[v, v^{-1}]$ of L is defined via the following rule:

$$\left\{ \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \right\} = \left\{ \begin{array}{c} \frown \\ \smile \end{array} \right\} - v \left\{ \begin{array}{c}) \\ (\end{array} \right\}$$

together with $\{\bigcirc L\} = (v + v^{-1})\{L\}$

and normalized by the conditions $\{\emptyset\} = 1$.

Example: Khovanov homology — Kauffman bracket

Definition. The **Kauffman bracket** $\{L\} \in \mathbb{Z}[v, v^{-1}]$ of L is defined via the following rule:

$$\left\{ \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \right\} = \left\{ \begin{array}{c} \frown \\ \smile \end{array} \right\} - v \left\{ \begin{array}{c}) \\ (\end{array} \right\}$$

together with $\{\bigcirc L\} = (v + v^{-1})\{L\}$

and normalized by the conditions $\{\emptyset\} = 1$.

Example: Khovanov homology — Kauffman bracket

Definition. The **Kauffman bracket** $\{L\} \in \mathbb{Z}[v, v^{-1}]$ of L is defined via the following rule:

$$\left\{ \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \right\} = \left\{ \begin{array}{c} \frown \\ \smile \end{array} \right\} - v \left\{ \begin{array}{c}) \\ (\end{array} \right\}$$

together with $\{\bigcirc L\} = (v + v^{-1})\{L\}$

and normalized by the conditions $\{\emptyset\} = 1$.

Example: Khovanov homology — Kauffman bracket

Definition. The **Kauffman bracket** $\{L\} \in \mathbb{Z}[v, v^{-1}]$ of L is defined via the following rule:

$$\left\{ \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \right\} = \left\{ \begin{array}{c} \frown \\ \smile \end{array} \right\} - v \left\{ \begin{array}{c} \text{)} \\ \text{(} \end{array} \right\}$$

together with $\{\bigcirc L\} = (v + v^{-1})\{L\}$

and normalized by the conditions $\{\emptyset\} = 1$.

Example: Khovanov homology — Jones polynomial

Definition. The **unnormalized Jones polynomial** $\hat{J}(L)$ of L is defined by

$$\hat{J}(L) := (-1)^{n-} v^{n+ - 2n-} \{L\} \in \mathbb{Z}[v, v^{-1}]$$

Definition. The (usual) **Jones polynomial** $J(L)$ is defined via $(v + v^{-1})J(L) = \hat{J}(L)$.

Theorem. [Jones] $J(L)$ is an invariant of an oriented link.

Example. For the Hopf link



we have $\hat{J} = (v + v^{-1})(v + v^5)$ and $J(H) = v + v^5$.

Example: Khovanov homology — Jones polynomial

Definition. The **unnormalized Jones polynomial** $\hat{J}(L)$ of L is defined by

$$\hat{J}(L) := (-1)^{n-} v^{n+ - 2n-} \{L\} \in \mathbb{Z}[v, v^{-1}]$$

Definition. The (usual) **Jones polynomial** $J(L)$ is defined via $(v + v^{-1})J(L) = \hat{J}(L)$.

Theorem. [Jones] $J(L)$ is an invariant of an oriented link.

Example. For the Hopf link



we have $\hat{J} = (v + v^{-1})(v + v^5)$ and $J(H) = v + v^5$.

Example: Khovanov homology — Jones polynomial

Definition. The **unnormalized Jones polynomial** $\hat{J}(L)$ of L is defined by

$$\hat{J}(L) := (-1)^{n-} v^{n+ - 2n-} \{L\} \in \mathbb{Z}[v, v^{-1}]$$

Definition. The (usual) **Jones polynomial** $J(L)$ is defined via $(v + v^{-1})J(L) = \hat{J}(L)$.

Theorem. [Jones] $J(L)$ is an invariant of an oriented link.

Example. For the Hopf link



we have $\hat{J} = (v + v^{-1})(v + v^5)$ and $J(H) = v + v^5$.

Example: Khovanov homology — Jones polynomial

Definition. The **unnormalized Jones polynomial** $\hat{J}(L)$ of L is defined by

$$\hat{J}(L) := (-1)^{n-} v^{n+ - 2n-} \{L\} \in \mathbb{Z}[v, v^{-1}]$$

Definition. The (usual) **Jones polynomial** $J(L)$ is defined via $(v + v^{-1})J(L) = \hat{J}(L)$.

Theorem. [Jones] $J(L)$ is an invariant of an oriented link.

Example. For the Hopf link



we have $\hat{J} = (v + v^{-1})(v + v^5)$ and $J(H) = v + v^5$.

Example: Khovanov homology — Jones polynomial

Definition. The **unnormalized Jones polynomial** $\hat{J}(L)$ of L is defined by

$$\hat{J}(L) := (-1)^{n-} v^{n+ - 2n-} \{L\} \in \mathbb{Z}[v, v^{-1}]$$

Definition. The (usual) **Jones polynomial** $J(L)$ is defined via $(v + v^{-1})J(L) = \hat{J}(L)$.

Theorem. [Jones] $J(L)$ is an invariant of an oriented link.

Example. For the **Hopf link**



we have $\hat{J} = (v + v^{-1})(v + v^5)$ and $J(H) = v + v^5$.

Example: Khovanov homology — Jones polynomial

Definition. The **unnormalized Jones polynomial** $\hat{J}(L)$ of L is defined by

$$\hat{J}(L) := (-1)^{n-} v^{n+ - 2n-} \{L\} \in \mathbb{Z}[v, v^{-1}]$$

Definition. The (usual) **Jones polynomial** $J(L)$ is defined via $(v + v^{-1})J(L) = \hat{J}(L)$.

Theorem. [Jones] $J(L)$ is an invariant of an oriented link.

Example. For the **Hopf link**



we have $\hat{J} = (v + v^{-1})(v + v^5)$ and $J(H) = v + v^5$.

Example: Khovanov homology — Jones polynomial

Definition. The **unnormalized Jones polynomial** $\hat{J}(L)$ of L is defined by

$$\hat{J}(L) := (-1)^{n-} v^{n+ - 2n-} \{L\} \in \mathbb{Z}[v, v^{-1}]$$

Definition. The (usual) **Jones polynomial** $J(L)$ is defined via $(v + v^{-1})J(L) = \hat{J}(L)$.

Theorem. [Jones] $J(L)$ is an invariant of an oriented link.

Example. For the **Hopf link**



we have $\hat{J} = (v + v^{-1})(v + v^5)$ and $J(H) = v + v^5$.

Example: Khovanov homology — characterization of J

Theorem. The Jones polynomial is uniquely determined by the property $J(\bigcirc) = 1$

and the **skain relation**

$$v^2 J \left(\begin{array}{c} \nearrow \\ \searrow \\ \swarrow \\ \searrow \end{array} \right) - v^{-2} J \left(\begin{array}{c} \nearrow \\ \swarrow \\ \searrow \\ \searrow \end{array} \right) = (v - v^{-1}) J \left(\begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} \right)$$

Example: Khovanov homology — characterization of J

Theorem. The Jones polynomial is uniquely determined by the property $J(\bigcirc) = 1$

and the **skein relation**

$$v^2 J \left(\begin{array}{c} \nearrow \\ \searrow \\ \swarrow \\ \searrow \end{array} \right) - v^{-2} J \left(\begin{array}{c} \nwarrow \\ \swarrow \\ \nearrow \\ \swarrow \end{array} \right) = (v - v^{-1}) J \left(\begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} \right)$$

Example: Khovanov homology — characterization of J

Theorem. The Jones polynomial is uniquely determined by the property $J(\bigcirc) = 1$

and the **skain relation**

$$v^2 J \left(\begin{array}{c} \nearrow \\ \searrow \\ \swarrow \\ \searrow \end{array} \right) - v^{-2} J \left(\begin{array}{c} \nwarrow \\ \swarrow \\ \nearrow \\ \nwarrow \end{array} \right) = (v - v^{-1}) J \left(\begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} \right)$$

Example: Khovanov homology — characterization of J

Theorem. The Jones polynomial is uniquely determined by the property $J(\bigcirc) = 1$

and the **skain relation**

$$v^2 J \left(\begin{array}{c} \nearrow \\ \nwarrow \\ \swarrow \\ \searrow \end{array} \right) - v^{-2} J \left(\begin{array}{c} \nwarrow \\ \nearrow \\ \swarrow \\ \searrow \end{array} \right) = (v - v^{-1}) J \left(\begin{array}{c} \uparrow \\ \uparrow \end{array} \right)$$

Example: Khovanov homology — characterization of J

Theorem. The Jones polynomial is uniquely determined by the property $J(\bigcirc) = 1$

and the **skain relation**

$$v^2 J \left(\begin{array}{c} \nearrow \\ \searrow \\ \swarrow \\ \searrow \end{array} \right) - v^{-2} J \left(\begin{array}{c} \nwarrow \\ \swarrow \\ \nearrow \\ \swarrow \end{array} \right) = (v - v^{-1}) J \left(\begin{array}{c} \uparrow \\ \uparrow \end{array} \right)$$

Example: Khovanov homology — idea and ingredients

Main idea: [Khovanov] Upgrade Kauffman bracket to a new bracket $[[\cdot]]$

$\mathbb{C}\text{-mod}$ — category of finite dimensional \mathbb{C} -vector spaces

$\mathbb{C}\text{-gmod}$ — category of finite dimensional graded \mathbb{C} -vector spaces

$\text{Com}^b(\mathbb{C}\text{-gmod})$ — category of finite complexes over $\mathbb{C}\text{-gmod}$

$[[\cdot]]$ takes values in $\text{Com}^b(\mathbb{C}\text{-mod})$

V — \mathbb{C} in degree 1 \oplus \mathbb{C} in degree -1

Example: Khovanov homology — idea and ingredients

Main idea: [Khovanov] Upgrade Kauffman bracket to a new bracket $[[\cdot]]$

$\mathbb{C}\text{-mod}$ — category of finite dimensional \mathbb{C} -vector spaces

$\mathbb{C}\text{-gmod}$ — category of finite dimensional graded \mathbb{C} -vector spaces

$\text{Com}^b(\mathbb{C}\text{-gmod})$ — category of finite complexes over $\mathbb{C}\text{-gmod}$

$[[\cdot]]$ takes values in $\text{Com}^b(\mathbb{C}\text{-mod})$

V — \mathbb{C} in degree 1 \oplus \mathbb{C} in degree -1

Example: Khovanov homology — idea and ingredients

Main idea: [Khovanov] Upgrade Kauffman bracket to a new bracket $[[\cdot]]$

$\mathbb{C}\text{-mod}$ — category of finite dimensional \mathbb{C} -vector spaces

$\mathbb{C}\text{-gmod}$ — category of finite dimensional graded \mathbb{C} -vector spaces

$\text{Com}^b(\mathbb{C}\text{-gmod})$ — category of finite complexes over $\mathbb{C}\text{-gmod}$

$[[\cdot]]$ takes values in $\text{Com}^b(\mathbb{C}\text{-mod})$

V — \mathbb{C} in degree 1 \oplus \mathbb{C} in degree -1

Example: Khovanov homology — idea and ingredients

Main idea: [Khovanov] Upgrade Kauffman bracket to a new bracket $[[\cdot]]$

$\mathbb{C}\text{-mod}$ — category of finite dimensional \mathbb{C} -vector spaces

$\mathbb{C}\text{-gmod}$ — category of finite dimensional graded \mathbb{C} -vector spaces

$\text{Com}^b(\mathbb{C}\text{-gmod})$ — category of finite complexes over $\mathbb{C}\text{-gmod}$

$[[\cdot]]$ takes values in $\text{Com}^b(\mathbb{C}\text{-mod})$

$V \text{ — } \mathbb{C} \text{ in degree } 1 \oplus \mathbb{C} \text{ in degree } -1$

Example: Khovanov homology — idea and ingredients

Main idea: [Khovanov] Upgrade Kauffman bracket to a new bracket $[[\cdot]]$

$\mathbb{C}\text{-mod}$ — category of finite dimensional \mathbb{C} -vector spaces

$\mathbb{C}\text{-gmod}$ — category of finite dimensional graded \mathbb{C} -vector spaces

$\text{Com}^b(\mathbb{C}\text{-gmod})$ — category of finite complexes over $\mathbb{C}\text{-gmod}$

$[[\cdot]]$ takes values in $\text{Com}^b(\mathbb{C}\text{-mod})$

$V \text{ — } \mathbb{C} \text{ in degree } 1 \oplus \mathbb{C} \text{ in degree } -1$

Example: Khovanov homology — idea and ingredients

Main idea: [Khovanov] Upgrade Kauffman bracket to a new bracket $[[\cdot]]$

$\mathbb{C}\text{-mod}$ — category of finite dimensional \mathbb{C} -vector spaces

$\mathbb{C}\text{-gmod}$ — category of finite dimensional graded \mathbb{C} -vector spaces

$\text{Com}^b(\mathbb{C}\text{-gmod})$ — category of finite complexes over $\mathbb{C}\text{-gmod}$

$[[\cdot]]$ takes values in $\text{Com}^b(\mathbb{C}\text{-mod})$

$V \text{ — } \mathbb{C} \text{ in degree } 1 \oplus \mathbb{C} \text{ in degree } -1$

Example: Khovanov homology — idea and ingredients

Main idea: [Khovanov] Upgrade Kauffman bracket to a new bracket $[[\cdot]]$

$\mathbb{C}\text{-mod}$ — category of finite dimensional \mathbb{C} -vector spaces

$\mathbb{C}\text{-gmod}$ — category of finite dimensional graded \mathbb{C} -vector spaces

$\text{Com}^b(\mathbb{C}\text{-gmod})$ — category of finite complexes over $\mathbb{C}\text{-gmod}$

$[[\cdot]]$ takes values in $\text{Com}^b(\mathbb{C}\text{-mod})$

V — \mathbb{C} in degree 1 \oplus \mathbb{C} in degree -1

Example: Khovanov homology — idea and ingredients

Main idea: [Khovanov] Upgrade Kauffman bracket to a new bracket $[[\cdot]]$

$\mathbb{C}\text{-mod}$ — category of finite dimensional \mathbb{C} -vector spaces

$\mathbb{C}\text{-gmod}$ — category of finite dimensional graded \mathbb{C} -vector spaces

$\text{Com}^b(\mathbb{C}\text{-gmod})$ — category of finite complexes over $\mathbb{C}\text{-gmod}$

$[[\cdot]]$ takes values in $\text{Com}^b(\mathbb{C}\text{-mod})$

V — \mathbb{C} in degree 1 \oplus \mathbb{C} in degree -1

Example: Khovanov homology — categorification

Categorification of normalization conditions:

$$[[\emptyset]] = 0 \rightarrow \mathbb{C} \rightarrow 0$$

$$[[\bigcirc L]] = V \otimes [L]$$

Categorification of the Kauffman bracket:

$$\left[\begin{array}{c} \diagdown \\ \diagup \end{array} \right] = \text{Total} \left(0 \rightarrow \left[\begin{array}{c} \frown \\ \smile \end{array} \right] \xrightarrow{d} \left[\begin{array}{c} \smile \\ \frown \end{array} \right] \langle -1 \rangle \rightarrow 0 \right)$$

Main difficulty: Definition of d .

Example: Khovanov homology — categorification

Categorification of normalization conditions:

$$[\emptyset] = 0 \rightarrow \mathbb{C} \rightarrow 0$$

$$[\bigcirc L] = V \otimes [L]$$

Categorification of the Kauffman bracket:

$$\left[\begin{array}{c} \diagdown \\ \diagup \end{array} \right] = \text{Total} \left(0 \rightarrow \left[\begin{array}{c} \frown \\ \smile \end{array} \right] \xrightarrow{d} \left[\begin{array}{c} \smile \\ \frown \end{array} \right] \langle -1 \rangle \rightarrow 0 \right)$$

Main difficulty: Definition of d .

Example: Khovanov homology — categorification

Categorification of normalization conditions:

$$[[\emptyset]] = 0 \rightarrow \mathbb{C} \rightarrow 0$$

$$[[\bigcirc L]] = V \otimes [L]$$

Categorification of the Kauffman bracket:

$$\left[\begin{array}{c} \diagup \\ \diagdown \end{array} \right] = \text{Total} \left(0 \rightarrow \left[\begin{array}{c} \text{---} \\ \text{---} \end{array} \right] \xrightarrow{d} \left[\begin{array}{c} \text{---} \\ \text{---} \end{array} \right] \langle -1 \rangle \rightarrow 0 \right)$$

Main difficulty: Definition of d .

Example: Khovanov homology — categorification

Categorification of normalization conditions:

$$[[\emptyset]] = 0 \rightarrow \mathbb{C} \rightarrow 0$$

$$[[\bigcirc L]] = V \otimes [[L]]$$

Categorification of the Kauffman bracket:

$$\left[\begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \right] = \text{Total} \left(0 \rightarrow \left[\begin{array}{c} \text{---} \\ \text{---} \end{array} \right] \xrightarrow{d} \left[\begin{array}{c} \text{---} \\ \text{---} \end{array} \right] \langle -1 \rangle \rightarrow 0 \right)$$

Main difficulty: Definition of d .

Example: Khovanov homology — categorification

Categorification of normalization conditions:

$$[[\emptyset]] = 0 \rightarrow \mathbb{C} \rightarrow 0$$

$$[[\bigcirc L]] = V \otimes [[L]]$$

Categorification of the Kauffman bracket:

$$\left[\begin{array}{c} \diagdown \\ \diagup \end{array} \right] = \text{Total} \left(0 \rightarrow \left[\begin{array}{c} \text{---} \\ \text{---} \end{array} \right] \xrightarrow{d} \left[\begin{array}{c} \text{---} \\ \text{---} \end{array} \right] \left(\begin{array}{c} \text{---} \\ \text{---} \end{array} \right) \langle -1 \rangle \rightarrow 0 \right)$$

Main difficulty: Definition of d .

Example: Khovanov homology — categorification

Categorification of normalization conditions:

$$[[\emptyset]] = 0 \rightarrow \mathbb{C} \rightarrow 0$$

$$[[\bigcirc L]] = V \otimes [[L]]$$

Categorification of the Kauffman bracket:

$$\left[\left[\begin{array}{c} \diagdown \\ \diagup \end{array} \right] \right] = \text{Total} \left(0 \rightarrow \left[\left[\begin{array}{c} \frown \\ \smile \end{array} \right] \right] \xrightarrow{d} \left[\left[\right] \right] \left(\left[\left[\langle -1 \rangle \right] \right] \rightarrow 0 \right) \right)$$

Main difficulty: Definition of d .

Example: Khovanov homology — categorification

Categorification of normalization conditions:

$$[[\emptyset]] = 0 \rightarrow \mathbb{C} \rightarrow 0$$

$$[[\bigcirc L]] = V \otimes [[L]]$$

Categorification of the Kauffman bracket:

$$\left[\left[\begin{array}{c} \diagdown \\ \diagup \end{array} \right] \right] = \text{Total} \left(0 \rightarrow \left[\begin{array}{c} \frown \\ \smile \end{array} \right] \xrightarrow{d} \left[\begin{array}{c} \text{) } \\ \text{ (} \end{array} \right] \left(\left[\langle -1 \rangle \right] \rightarrow 0 \right) \right)$$

Main difficulty: Definition of d .

Example: Khovanov homology — the result

$[\cdot]$ — shift in homological position

$\langle \cdot \rangle$ — shift in grading

Theorem. [Khovanov]

Homology of $[[\cdot]][n_-]\langle n_+ - 2n_- \rangle$ is an invariant of an oriented link.

Note: $[[\cdot]][n_-]\langle n_+ - 2n_- \rangle$ is not an invariant of an oriented link.

Decategorification theorem. [Khovanov]

Graded Euler characteristic of $[[L]][n_-]\langle n_+ - 2n_- \rangle$ equals $\hat{J}(L)$.

Benefit: Khovanov homology is a strictly stronger invariant than Jones polynomial. For example:

Theorem. [Kronheimer-Mrowka] Khovanov homology detects the unknot.

Example: Khovanov homology — the result

$[\cdot]$ — shift in homological position

$\langle \cdot \rangle$ — shift in grading

Theorem. [Khovanov]

Homology of $[[\cdot]][n_-]\langle n_+ - 2n_- \rangle$ is an invariant of an oriented link.

Note: $[[\cdot]][n_-]\langle n_+ - 2n_- \rangle$ is not an invariant of an oriented link.

Decategorification theorem. [Khovanov]

Graded Euler characteristic of $[[L]][n_-]\langle n_+ - 2n_- \rangle$ equals $\hat{J}(L)$.

Benefit: Khovanov homology is a strictly stronger invariant than Jones polynomial. For example:

Theorem. [Kronheimer-Mrowka] Khovanov homology detects the unknot.

Example: Khovanov homology — the result

$[\cdot]$ — shift in homological position

$\langle \cdot \rangle$ — shift in grading

Theorem. [Khovanov]

Homology of $[[\cdot]][n_-]\langle n_+ - 2n_- \rangle$ is an invariant of an oriented link.

Note: $[[\cdot]][n_-]\langle n_+ - 2n_- \rangle$ is not an invariant of an oriented link.

Decategorification theorem. [Khovanov]

Graded Euler characteristic of $[[L]][n_-]\langle n_+ - 2n_- \rangle$ equals $\hat{J}(L)$.

Benefit: Khovanov homology is a strictly stronger invariant than Jones polynomial. For example:

Theorem. [Kronheimer-Mrowka] Khovanov homology detects the unknot.

Example: Khovanov homology — the result

$[\cdot]$ — shift in homological position

$\langle \cdot \rangle$ — shift in grading

Theorem. [Khovanov]

Homology of $[[\cdot]][n_-]\langle n_+ - 2n_- \rangle$ is an invariant of an oriented link.

Note: $[[\cdot]][n_-]\langle n_+ - 2n_- \rangle$ is not an invariant of an oriented link.

Decategorification theorem. [Khovanov]

Graded Euler characteristic of $[[L]][n_-]\langle n_+ - 2n_- \rangle$ equals $\hat{J}(L)$.

Benefit: Khovanov homology is a strictly stronger invariant than Jones polynomial. For example:

Theorem. [Kronheimer-Mrowka] Khovanov homology detects the unknot.

Example: Khovanov homology — the result

$[\cdot]$ — shift in homological position

$\langle \cdot \rangle$ — shift in grading

Theorem. [Khovanov]

Homology of $[[\cdot]][n_-]\langle n_+ - 2n_- \rangle$ is an invariant of an oriented link.

Note: $[[\cdot]][n_-]\langle n_+ - 2n_- \rangle$ is not an invariant of an oriented link.

Decategorification theorem. [Khovanov]

Graded Euler characteristic of $[[L]][n_-]\langle n_+ - 2n_- \rangle$ equals $\hat{J}(L)$.

Benefit: Khovanov homology is a strictly stronger invariant than Jones polynomial. For example:

Theorem. [Kronheimer-Mrowka] Khovanov homology detects the unknot.

Example: Khovanov homology — the result

$[\cdot]$ — shift in homological position

$\langle \cdot \rangle$ — shift in grading

Theorem. [Khovanov]

Homology of $[[\cdot]][n_-]\langle n_+ - 2n_- \rangle$ is an invariant of an oriented link.

Note: $[[\cdot]][n_-]\langle n_+ - 2n_- \rangle$ is not an invariant of an oriented link.

Decategorification theorem. [Khovanov]

Graded Euler characteristic of $[[L]][n_-]\langle n_+ - 2n_- \rangle$ equals $\hat{J}(L)$.

Benefit: Khovanov homology is a strictly stronger invariant than Jones polynomial. For example:

Theorem. [Kronheimer-Mrowka] Khovanov homology detects the unknot.

Example: Khovanov homology — the result

$[\cdot]$ — shift in homological position

$\langle \cdot \rangle$ — shift in grading

Theorem. [Khovanov]

Homology of $[[\cdot]][n_-]\langle n_+ - 2n_- \rangle$ is an invariant of an oriented link.

Note: $[[\cdot]][n_-]\langle n_+ - 2n_- \rangle$ is not an invariant of an oriented link.

Decategorification theorem. [Khovanov]

Graded Euler characteristic of $[[L]][n_-]\langle n_+ - 2n_- \rangle$ equals $\hat{J}(L)$.

Benefit: Khovanov homology is a strictly stronger invariant than Jones polynomial. For example:

Theorem. [Kronheimer-Mrowka] Khovanov homology detects the unknot.

Example: Khovanov homology — the result

$[\cdot]$ — shift in homological position

$\langle \cdot \rangle$ — shift in grading

Theorem. [Khovanov]

Homology of $[[\cdot]][n_-]\langle n_+ - 2n_- \rangle$ is an invariant of an oriented link.

Note: $[[\cdot]][n_-]\langle n_+ - 2n_- \rangle$ is not an invariant of an oriented link.

Decategorification theorem. [Khovanov]

Graded Euler characteristic of $[[L]][n_-]\langle n_+ - 2n_- \rangle$ equals $\hat{J}(L)$.

Benefit: Khovanov homology is a strictly stronger invariant than Jones polynomial. For example:

Theorem. [Kronheimer-Mrowka] Khovanov homology detects the unknot.

Example: Khovanov homology — the result

$[\cdot]$ — shift in homological position

$\langle \cdot \rangle$ — shift in grading

Theorem. [Khovanov]

Homology of $[[\cdot]][n_-]\langle n_+ - 2n_- \rangle$ is an invariant of an oriented link.

Note: $[[\cdot]][n_-]\langle n_+ - 2n_- \rangle$ is not an invariant of an oriented link.

Decategorification theorem. [Khovanov]

Graded Euler characteristic of $[[L]][n_-]\langle n_+ - 2n_- \rangle$ equals $\hat{J}(L)$.

Benefit: Khovanov homology is a strictly stronger invariant than Jones polynomial. For example:

Theorem. [Kronheimer-Mrowka] Khovanov homology detects the unknot.

Alternative approach — Alexander theorem

Theorem. [Alexander] Every link is a closure of a braid

Elementary diagrams:



the cup diagram



the cap diagram



right crossing



left crossing

Corollary. Every oriented link is a composition of elementary diagrams.

Alternative approach — Alexander theorem

Theorem. [Alexander] Every link is a closure of a braid

Elementary diagrams:



the cup diagram



the cap diagram



right crossing



left crossing

Corollary. Every oriented link is a composition of elementary diagrams.

Alternative approach — Alexander theorem

Theorem. [Alexander] Every link is a closure of a braid

Elementary diagrams:



the cup diagram



the cap diagram



right crossing



left crossing

Corollary. Every oriented link is a composition of elementary diagrams.

Alternative approach — Alexander theorem

Theorem. [Alexander] Every link is a closure of a braid

Elementary diagrams:



the cup diagram



the cap diagram



right crossing



left crossing

Corollary. Every oriented link is a composition of elementary diagrams.

Alternative approach — Alexander theorem

Theorem. [Alexander] Every link is a closure of a braid

Elementary diagrams:



the cup diagram



the cap diagram



right crossing



left crossing

Corollary. Every oriented link is a composition of elementary diagrams.

Alternative approach — Alexander theorem

Theorem. [Alexander] Every link is a closure of a braid

Elementary diagrams:



the cup diagram



the cap diagram



right crossing



left crossing

Corollary. Every oriented link is a composition of elementary diagrams.

Alternative approach — the Hopf link example

Example. For the Hopf link we could take:




Alternative approach — the Hopf link example

Example. For the Hopf link we could take:




Alternative approach — the Hopf link example

Example. For the Hopf link we could take:



Alternative approach — the Hopf link example

Example. For the Hopf link we could take:



Alternative approach — quantum groups

\mathfrak{g} — simple finite dimensional Lie algebra

$U(\mathfrak{g})$ — the universal enveloping algebra of \mathfrak{g}

Fact. $U(\mathfrak{g})$ is a cocommutative Hopf algebra.

Consequence. The isomorphism $V \otimes W \cong W \otimes V$ is involutive.

$U_v(\mathfrak{g})$ — the quantum enveloping algebra of \mathfrak{g}

Fact. $U_v(\mathfrak{g})$ is a Hopf algebra, not cocommutative.

Consequence. The isomorphism $V \otimes W \cong W \otimes V$ is not involutive.

Alternative approach — quantum groups

\mathfrak{g} — simple finite dimensional Lie algebra

$U(\mathfrak{g})$ — the universal enveloping algebra of \mathfrak{g}

Fact. $U(\mathfrak{g})$ is a cocommutative Hopf algebra.

Consequence. The isomorphism $V \otimes W \cong W \otimes V$ is involutive.

$U_v(\mathfrak{g})$ — the quantum enveloping algebra of \mathfrak{g}

Fact. $U_v(\mathfrak{g})$ is a Hopf algebra, not cocommutative.

Consequence. The isomorphism $V \otimes W \cong W \otimes V$ is not involutive.

Alternative approach — quantum groups

\mathfrak{g} — simple finite dimensional Lie algebra

$U(\mathfrak{g})$ — the universal enveloping algebra of \mathfrak{g}

Fact. $U(\mathfrak{g})$ is a cocommutative Hopf algebra.

Consequence. The isomorphism $V \otimes W \cong W \otimes V$ is involutive.

$U_v(\mathfrak{g})$ — the quantum enveloping algebra of \mathfrak{g}

Fact. $U_v(\mathfrak{g})$ is a Hopf algebra, not cocommutative.

Consequence. The isomorphism $V \otimes W \cong W \otimes V$ is not involutive.

Alternative approach — quantum groups

\mathfrak{g} — simple finite dimensional Lie algebra

$U(\mathfrak{g})$ — the universal enveloping algebra of \mathfrak{g}

Fact. $U(\mathfrak{g})$ is a cocommutative Hopf algebra.

Consequence. The isomorphism $V \otimes W \cong W \otimes V$ is involutive.

$U_v(\mathfrak{g})$ — the quantum enveloping algebra of \mathfrak{g}

Fact. $U_v(\mathfrak{g})$ is a Hopf algebra, not cocommutative.

Consequence. The isomorphism $V \otimes W \cong W \otimes V$ is not involutive.

Alternative approach — quantum groups

\mathfrak{g} — simple finite dimensional Lie algebra

$U(\mathfrak{g})$ — the universal enveloping algebra of \mathfrak{g}

Fact. $U(\mathfrak{g})$ is a cocommutative Hopf algebra.

Consequence. The isomorphism $V \otimes W \cong W \otimes V$ is involutive.

$U_v(\mathfrak{g})$ — the quantum enveloping algebra of \mathfrak{g}

Fact. $U_v(\mathfrak{g})$ is a Hopf algebra, not cocommutative.

Consequence. The isomorphism $V \otimes W \cong W \otimes V$ is not involutive.

Alternative approach — quantum groups

\mathfrak{g} — simple finite dimensional Lie algebra

$U(\mathfrak{g})$ — the universal enveloping algebra of \mathfrak{g}

Fact. $U(\mathfrak{g})$ is a cocommutative Hopf algebra.

Consequence. The isomorphism $V \otimes W \cong W \otimes V$ is involutive.

$U_v(\mathfrak{g})$ — the quantum enveloping algebra of \mathfrak{g}

Fact. $U_v(\mathfrak{g})$ is a Hopf algebra, not cocommutative.

Consequence. The isomorphism $V \otimes W \cong W \otimes V$ is not involutive.

Alternative approach — quantum groups

\mathfrak{g} — simple finite dimensional Lie algebra

$U(\mathfrak{g})$ — the universal enveloping algebra of \mathfrak{g}

Fact. $U(\mathfrak{g})$ is a cocommutative Hopf algebra.

Consequence. The isomorphism $V \otimes W \cong W \otimes V$ is involutive.

$U_v(\mathfrak{g})$ — the quantum enveloping algebra of \mathfrak{g}

Fact. $U_v(\mathfrak{g})$ is a Hopf algebra, not cocommutative.

Consequence. The isomorphism $V \otimes W \cong W \otimes V$ is not involutive.

Alternative approach — quantum groups

\mathfrak{g} — simple finite dimensional Lie algebra

$U(\mathfrak{g})$ — the universal enveloping algebra of \mathfrak{g}

Fact. $U(\mathfrak{g})$ is a cocommutative Hopf algebra.

Consequence. The isomorphism $V \otimes W \cong W \otimes V$ is involutive.

$U_v(\mathfrak{g})$ — the quantum enveloping algebra of \mathfrak{g}

Fact. $U_v(\mathfrak{g})$ is a Hopf algebra, not cocommutative.

Consequence. The isomorphism $V \otimes W \cong W \otimes V$ is not involutive.

Alternative approach — quantum groups

\mathfrak{g} — simple finite dimensional Lie algebra

$U(\mathfrak{g})$ — the universal enveloping algebra of \mathfrak{g}

Fact. $U(\mathfrak{g})$ is a cocommutative Hopf algebra.

Consequence. The isomorphism $V \otimes W \cong W \otimes V$ is involutive.

$U_v(\mathfrak{g})$ — the quantum enveloping algebra of \mathfrak{g}

Fact. $U_v(\mathfrak{g})$ is a Hopf algebra, not cocommutative.

Consequence. The isomorphism $V \otimes W \cong W \otimes V$ is not involutive.

Alternative approach — tangles

Tang — the category of oriented **tangles**

Objects: Non-negative integers

Informally: $n \in \{0, 1, 2, \dots\}$ should be thought of as a collection of n points.

Morphisms: Oriented diagrams generated by (oriented) elementary diagrams (up to isotopy), connecting the corresponding points, read from bottom to top.

Composition: Concatenation

Example 1: An oriented cup diagram is a morphism from 0 to 2.

Example 2: An oriented cap diagram is a morphism from 2 to 0.

Example 3: An oriented crossing is a morphism from 2 to 2.

Alternative approach — tangles

Tang — the category of oriented **tangles**

Objects: Non-negative integers

Informally: $n \in \{0, 1, 2, \dots\}$ should be thought of as a collection of n points.

Morphisms: Oriented diagrams generated by (oriented) elementary diagrams (up to isotopy), connecting the corresponding points, read from bottom to top.

Composition: Concatenation

Example 1: An oriented cup diagram is a morphism from 0 to 2.

Example 2: An oriented cap diagram is a morphism from 2 to 0.

Example 3: An oriented crossing is a morphism from 2 to 2.

Alternative approach — tangles

Tang — the category of oriented **tangles**

Objects: Non-negative integers

Informally: $n \in \{0, 1, 2, \dots\}$ should be thought of as a collection of n points.

Morphisms: Oriented diagrams generated by (oriented) elementary diagrams (up to isotopy), connecting the corresponding points, read from bottom to top.

Composition: Concatenation

Example 1: An oriented cup diagram is a morphism from 0 to 2.

Example 2: An oriented cap diagram is a morphism from 2 to 0.

Example 3: An oriented crossing is a morphism from 2 to 2.

Alternative approach — tangles

Tang — the category of oriented **tangles**

Objects: Non-negative integers

Informally: $n \in \{0, 1, 2, \dots\}$ should be thought of as a collection of n points.

Morphisms: Oriented diagrams generated by (oriented) elementary diagrams (up to isotopy), connecting the corresponding points, read from bottom to top.

Composition: Concatenation

Example 1: An oriented cup diagram is a morphism from 0 to 2.

Example 2: An oriented cap diagram is a morphism from 2 to 0.

Example 3: An oriented crossing is a morphism from 2 to 2.

Alternative approach — tangles

Tang — the category of oriented **tangles**

Objects: **Non-negative** integers

Informally: $n \in \{0, 1, 2, \dots\}$ should be thought of as a collection of n points.

Morphisms: **Oriented diagrams** generated by (oriented) elementary diagrams (up to isotopy), connecting the corresponding points, read from bottom to top.

Composition: **Concatenation**

Example 1: An oriented cup diagram is a morphism from 0 to 2.

Example 2: An oriented cap diagram is a morphism from 2 to 0.

Example 3: An oriented crossing is a morphism from 2 to 2.

Alternative approach — tangles

Tang — the category of oriented **tangles**

Objects: **Non-negative** integers

Informally: $n \in \{0, 1, 2, \dots\}$ should be thought of as a collection of n points.

Morphisms: **Oriented diagrams** generated by (oriented) elementary diagrams (up to isotopy), connecting the corresponding points, read from bottom to top.

Composition: **Concatenation**

Example 1: An oriented cup diagram is a morphism from 0 to 2.

Example 2: An oriented cap diagram is a morphism from 2 to 0.

Example 3: An oriented crossing is a morphism from 2 to 2.

Alternative approach — tangles

Tang — the category of oriented **tangles**

Objects: **Non-negative** integers

Informally: $n \in \{0, 1, 2, \dots\}$ should be thought of as a collection of n points.

Morphisms: **Oriented diagrams** generated by (oriented) elementary diagrams (up to isotopy), connecting the corresponding points, read from bottom to top.

Composition: **Concatenation**

Example 1: An oriented cup diagram is a morphism from 0 to 2.

Example 2: An oriented cap diagram is a morphism from 2 to 0.

Example 3: An oriented crossing is a morphism from 2 to 2.

Alternative approach — tangles

Tang — the category of oriented **tangles**

Objects: **Non-negative** integers

Informally: $n \in \{0, 1, 2, \dots\}$ should be thought of as a collection of n points.

Morphisms: **Oriented diagrams** generated by (oriented) elementary diagrams (up to isotopy), connecting the corresponding points, read from bottom to top.

Composition: **Concatenation**

Example 1: An oriented cup diagram is a morphism from 0 to 2.

Example 2: An oriented cap diagram is a morphism from 2 to 0.

Example 3: An oriented crossing is a morphism from 2 to 2.

Alternative approach — tangles

Tang — the category of oriented **tangles**

Objects: **Non-negative** integers

Informally: $n \in \{0, 1, 2, \dots\}$ should be thought of as a collection of n points.

Morphisms: **Oriented diagrams** generated by (oriented) elementary diagrams (up to isotopy), connecting the corresponding points, read from bottom to top.

Composition: **Concatenation**

Example 1: An oriented cup diagram is a morphism from 0 to 2.

Example 2: An oriented cap diagram is a morphism from 2 to 0.

Example 3: An oriented crossing is a morphism from 2 to 2.

Alternative approach — tangles

Tang — the category of oriented **tangles**

Objects: **Non-negative** integers

Informally: $n \in \{0, 1, 2, \dots\}$ should be thought of as a collection of n points.

Morphisms: **Oriented diagrams** generated by (oriented) elementary diagrams (up to isotopy), connecting the corresponding points, read from bottom to top.

Composition: **Concatenation**

Example 1: An oriented cup diagram is a morphism from 0 to 2.

Example 2: An oriented cap diagram is a morphism from 2 to 0.

Example 3: An oriented crossing is a morphism from 2 to 2.

Alternative approach — idea of quantum knot invariants

Idea of quantum knot invariants. [Reshetikhin-Turaev]

Consider some $U_{\mathfrak{v}}(\mathfrak{g})$.

V — the “natural” $U_{\mathfrak{v}}(\mathfrak{g})$ -module

Define a functor $F : \mathbf{Tang} \rightarrow U_{\mathfrak{v}}(\mathfrak{g})\text{-mod}$

$F(n) := V^{\otimes n}$, where $F(0) := \mathbb{C}(v)$

$F(\text{elementary diagram}) :=$ certain explicit homomorphisms of $U_{\mathfrak{v}}(\mathfrak{g})$ -modules

oriented link $L \rightarrow$ tangle $T_L \rightarrow$ endom. $F(T_L)$ of $\mathbb{C}(v)$

Consequence: $F(T_L)(1)$ is an invariant of L .

Alternative approach — idea of quantum knot invariants

Idea of quantum knot invariants. [Reshetikhin-Turaev]

Consider some $U_{\nu}(\mathfrak{g})$.

V — the “natural” $U_{\nu}(\mathfrak{g})$ -module

Define a functor $F : \mathbf{Tang} \rightarrow U_{\nu}(\mathfrak{g})\text{-mod}$

$F(n) := V^{\otimes n}$, where $F(0) := \mathbb{C}(v)$

$F(\text{elementary diagram}) :=$ certain explicit homomorphisms of $U_{\nu}(\mathfrak{g})$ -modules

oriented link $L \rightarrow$ tangle $T_L \rightarrow$ endom. $F(T_L)$ of $\mathbb{C}(v)$

Consequence: $F(T_L)(1)$ is an invariant of L .

Alternative approach — idea of quantum knot invariants

Idea of quantum knot invariants. [Reshetikhin-Turaev]

Consider some $U_{\nu}(\mathfrak{g})$.

V — the “natural” $U_{\nu}(\mathfrak{g})$ -module

Define a functor $F : \mathbf{Tang} \rightarrow U_{\nu}(\mathfrak{g})\text{-mod}$

$F(n) := V^{\otimes n}$, where $F(0) := \mathbb{C}(v)$

$F(\text{elementary diagram}) :=$ certain explicit homomorphisms of $U_{\nu}(\mathfrak{g})$ -modules

oriented link $L \rightarrow$ tangle $T_L \rightarrow$ endom. $F(T_L)$ of $\mathbb{C}(v)$

Consequence: $F(T_L)(1)$ is an invariant of L .

Alternative approach — idea of quantum knot invariants

Idea of quantum knot invariants. [Reshetikhin-Turaev]

Consider some $U_{\nu}(\mathfrak{g})$.

V — the “natural” $U_{\nu}(\mathfrak{g})$ -module

Define a functor $F : \mathbf{Tang} \rightarrow U_{\nu}(\mathfrak{g})\text{-mod}$

$F(n) := V^{\otimes n}$, where $F(0) := \mathbb{C}(v)$

$F(\text{elementary diagram}) :=$ certain explicit homomorphisms of $U_{\nu}(\mathfrak{g})$ -modules

oriented link $L \rightarrow$ tangle $T_L \rightarrow$ endom. $F(T_L)$ of $\mathbb{C}(v)$

Consequence: $F(T_L)(1)$ is an invariant of L .

Alternative approach — idea of quantum knot invariants

Idea of quantum knot invariants. [Reshetikhin-Turaev]

Consider some $U_V(\mathfrak{g})$.

V — the “natural” $U_V(\mathfrak{g})$ -module

Define a functor $F : \mathbf{Tang} \rightarrow U_V(\mathfrak{g})\text{-mod}$

$F(n) := V^{\otimes n}$, where $F(0) := \mathbb{C}(v)$

$F(\text{elementary diagram}) :=$ certain explicit homomorphisms of $U_V(\mathfrak{g})$ -modules

oriented link $L \rightarrow$ tangle $T_L \rightarrow$ endom. $F(T_L)$ of $\mathbb{C}(v)$

Consequence: $F(T_L)(1)$ is an invariant of L .

Alternative approach — idea of quantum knot invariants

Idea of quantum knot invariants. [Reshetikhin-Turaev]

Consider some $U_{\mathfrak{v}}(\mathfrak{g})$.

V — the “natural” $U_{\mathfrak{v}}(\mathfrak{g})$ -module

Define a functor $F : \mathbf{Tang} \rightarrow U_{\mathfrak{v}}(\mathfrak{g})\text{-mod}$

$F(n) := V^{\otimes n}$, where $F(0) := \mathbb{C}(v)$

$F(\text{elementary diagram}) :=$ certain explicit homomorphisms of $U_{\mathfrak{v}}(\mathfrak{g})$ -modules

oriented link $L \rightarrow$ tangle $T_L \rightarrow$ endom. $F(T_L)$ of $\mathbb{C}(v)$

Consequence: $F(T_L)(1)$ is an invariant of L .

Alternative approach — idea of quantum knot invariants

Idea of quantum knot invariants. [Reshetikhin-Turaev]

Consider some $U_{\mathfrak{v}}(\mathfrak{g})$.

V — the “natural” $U_{\mathfrak{v}}(\mathfrak{g})$ -module

Define a functor $F : \mathbf{Tang} \rightarrow U_{\mathfrak{v}}(\mathfrak{g})\text{-mod}$

$F(n) := V^{\otimes n}$, where $F(0) := \mathbb{C}(v)$

$F(\text{elementary diagram}) :=$ certain explicit homomorphisms of $U_{\mathfrak{v}}(\mathfrak{g})$ -modules

oriented link $L \rightarrow$ tangle $T_L \rightarrow$ endom. $F(T_L)$ of $\mathbb{C}(v)$

Consequence: $F(T_L)(1)$ is an invariant of L .

Alternative approach — idea of quantum knot invariants

Idea of quantum knot invariants. [Reshetikhin-Turaev]

Consider some $U_{\mathfrak{v}}(\mathfrak{g})$.

V — the “natural” $U_{\mathfrak{v}}(\mathfrak{g})$ -module

Define a functor $F : \mathbf{Tang} \rightarrow U_{\mathfrak{v}}(\mathfrak{g})\text{-mod}$

$F(n) := V^{\otimes n}$, where $F(0) := \mathbb{C}(v)$

$F(\text{elementary diagram}) :=$ certain explicit homomorphisms of $U_{\mathfrak{v}}(\mathfrak{g})$ -modules

oriented link $L \rightarrow$ tangle $T_L \rightarrow$ endom. $F(T_L)$ of $\mathbb{C}(v)$

Consequence: $F(T_L)(1)$ is an invariant of L .

Alternative approach — idea of quantum knot invariants

Idea of quantum knot invariants. [Reshetikhin-Turaev]

Consider some $U_{\mathfrak{v}}(\mathfrak{g})$.

V — the “natural” $U_{\mathfrak{v}}(\mathfrak{g})$ -module

Define a functor $F : \mathbf{Tang} \rightarrow U_{\mathfrak{v}}(\mathfrak{g})\text{-mod}$

$F(n) := V^{\otimes n}$, where $F(0) := \mathbb{C}(v)$

$F(\text{elementary diagram}) :=$ certain explicit homomorphisms of $U_{\mathfrak{v}}(\mathfrak{g})$ -modules

oriented link $L \rightarrow$ tangle $T_L \rightarrow$ endom. $F(T_L)$ of $\mathbb{C}(v)$

Consequence: $F(T_L)(1)$ is an invariant of L .

Alternative approach — idea of quantum knot invariants

Idea of quantum knot invariants. [Reshetikhin-Turaev]

Consider some $U_{\mathfrak{v}}(\mathfrak{g})$.

V — the “natural” $U_{\mathfrak{v}}(\mathfrak{g})$ -module

Define a functor $F : \mathbf{Tang} \rightarrow U_{\mathfrak{v}}(\mathfrak{g})\text{-mod}$

$F(n) := V^{\otimes n}$, where $F(0) := \mathbb{C}(v)$

$F(\text{elementary diagram}) :=$ certain explicit homomorphisms of $U_{\mathfrak{v}}(\mathfrak{g})$ -modules

oriented link $L \rightarrow$ tangle $T_L \rightarrow$ endom. $F(T_L)$ of $\mathbb{C}(v)$

Consequence: $F(T_L)(1)$ is an invariant of L .

Alternative approach — idea of quantum knot invariants

Idea of quantum knot invariants. [Reshetikhin-Turaev]

Consider some $U_{\nu}(\mathfrak{g})$.

V — the “natural” $U_{\nu}(\mathfrak{g})$ -module

Define a functor $F : \mathbf{Tang} \rightarrow U_{\nu}(\mathfrak{g})\text{-mod}$

$F(n) := V^{\otimes n}$, where $F(0) := \mathbb{C}(v)$

$F(\text{elementary diagram}) :=$ certain explicit homomorphisms of $U_{\nu}(\mathfrak{g})$ -modules

oriented link $L \rightarrow$ tangle $T_L \rightarrow$ endom. $F(T_L)$ of $\mathbb{C}(v)$

Consequence: $F(T_L)(1)$ is an invariant of L .

Alternative approach — idea of quantum knot invariants

Idea of quantum knot invariants. [Reshetikhin-Turaev]

Consider some $U_{\mathfrak{g}}(\mathfrak{g})$.

V — the “natural” $U_{\mathfrak{g}}(\mathfrak{g})$ -module

Define a functor $F : \mathbf{Tang} \rightarrow U_{\mathfrak{g}}(\mathfrak{g})\text{-mod}$

$F(n) := V^{\otimes n}$, where $F(0) := \mathbb{C}(v)$

$F(\text{elementary diagram}) :=$ certain explicit homomorphisms of $U_{\mathfrak{g}}(\mathfrak{g})$ -modules

oriented link $L \rightarrow$ tangle $T_L \rightarrow$ endom. $F(T_L)$ of $\mathbb{C}(v)$

Consequence: $F(T_L)(1)$ is an invariant of L .

Quantum invariants — $U_q(\mathfrak{sl}_2)$

Definition: $U_v(\mathfrak{sl}_2)$ has generators E, F, K, K^{-1} and relations

$$KE = v^2 EK, \quad KF = v^{-2} FK, \quad KK^{-1} = K^{-1}K = 1,$$

$$EF - FE = \frac{K - K^{-1}}{v - v^{-1}}.$$

Hopf structure:

$$\Delta(E) = 1 \otimes E + E \otimes K, \quad \Delta(F) = K^{-1} \otimes F + F \otimes 1, \quad \Delta(K^{\pm 1}) = K^{\pm 1} \otimes K^{\pm 1}.$$

Quantum invariants — $U_q(\mathfrak{sl}_2)$

Definition: $U_v(\mathfrak{sl}_2)$ has **generators** E, F, K, K^{-1} and **relations**

$$KE = v^2 EK, \quad KF = v^{-2} FK, \quad KK^{-1} = K^{-1}K = 1,$$

$$EF - FE = \frac{K - K^{-1}}{v - v^{-1}}.$$

Hopf structure:

$$\Delta(E) = 1 \otimes E + E \otimes K, \quad \Delta(F) = K^{-1} \otimes F + F \otimes 1, \quad \Delta(K^{\pm 1}) = K^{\pm 1} \otimes K^{\pm 1}.$$

Quantum invariants — $U_q(\mathfrak{sl}_2)$

Definition: $U_v(\mathfrak{sl}_2)$ has **generators** E, F, K, K^{-1} and **relations**

$$KE = v^2EK, \quad KF = v^{-2}FK, \quad KK^{-1} = K^{-1}K = 1,$$

$$EF - FE = \frac{K - K^{-1}}{v - v^{-1}}.$$

Hopf structure:

$$\Delta(E) = 1 \otimes E + E \otimes K, \quad \Delta(F) = K^{-1} \otimes F + F \otimes 1, \quad \Delta(K^{\pm 1}) = K^{\pm 1} \otimes K^{\pm 1}.$$

Quantum invariants — $U_q(\mathfrak{sl}_2)$

Definition: $U_v(\mathfrak{sl}_2)$ has **generators** E, F, K, K^{-1} and **relations**

$$KE = v^2EK, \quad KF = v^{-2}FK, \quad KK^{-1} = K^{-1}K = 1,$$

$$EF - FE = \frac{K - K^{-1}}{v - v^{-1}}.$$

Hopf structure:

$$\Delta(E) = 1 \otimes E + E \otimes K, \quad \Delta(F) = K^{-1} \otimes F + F \otimes 1, \quad \Delta(K^{\pm 1}) = K^{\pm 1} \otimes K^{\pm 1}.$$

Quantum invariants — $U_q(\mathfrak{sl}_2)$

Definition: $U_v(\mathfrak{sl}_2)$ has **generators** E, F, K, K^{-1} and **relations**

$$KE = v^2EK, \quad KF = v^{-2}FK, \quad KK^{-1} = K^{-1}K = 1,$$

$$EF - FE = \frac{K - K^{-1}}{v - v^{-1}}.$$

Hopf structure:

$$\Delta(E) = 1 \otimes E + E \otimes K, \quad \Delta(F) = K^{-1} \otimes F + F \otimes 1, \quad \Delta(K^{\pm 1}) = K^{\pm 1} \otimes K^{\pm 1}.$$

Quantum invariants — $U_q(\mathfrak{sl}_2)$

Definition: $U_v(\mathfrak{sl}_2)$ has generators E, F, K, K^{-1} and relations

$$KE = v^2EK, \quad KF = v^{-2}FK, \quad KK^{-1} = K^{-1}K = 1,$$

$$EF - FE = \frac{K - K^{-1}}{v - v^{-1}}.$$

Hopf structure:

$$\Delta(E) = 1 \otimes E + E \otimes K, \quad \Delta(F) = K^{-1} \otimes F + F \otimes 1, \quad \Delta(K^{\pm 1}) = K^{\pm 1} \otimes K^{\pm 1}.$$

Quantum invariants — natural $U_v(\mathfrak{sl}_2)$ -module

Quantum numbers: $[a] := \frac{v^a - v^{-a}}{v - v^{-1}}$, $a \in \mathbb{Z}$

V — the “natural” $U_v(\mathfrak{sl}_2)$ -module

Basis: w_0 and w_1

Action:

$$\begin{aligned} Ew_k &= [k+1]w_{k+1}, & Fw_k &= -[n-k+1]w_{k-1}, \\ K^{\pm 1}w_k &= -v^{\pm(2k-n)}w_k \end{aligned}$$

Notation: $w_{i_1} \otimes w_{i_2} \otimes \cdots \otimes w_{i_k}$ denoted by $i_1 i_2 \dots i_k$

Consequence: Basis in $V^{\otimes n}$ consists of 0-1-sequences of length n .

Quantum invariants — natural $U_v(\mathfrak{sl}_2)$ -module

Quantum numbers: $[a] := \frac{v^a - v^{-a}}{v - v^{-1}}$, $a \in \mathbb{Z}$

V — the “natural” $U_v(\mathfrak{sl}_2)$ -module

Basis: w_0 and w_1

Action:

$$\begin{aligned} Ew_k &= [k+1]w_{k+1}, & Fw_k &= -[n-k+1]w_{k-1}, \\ K^{\pm 1}w_k &= -v^{\pm(2k-n)}w_k \end{aligned}$$

Notation: $w_{i_1} \otimes w_{i_2} \otimes \cdots \otimes w_{i_k}$ denoted by $i_1 i_2 \dots i_k$

Consequence: Basis in $V^{\otimes n}$ consists of 0-1-sequences of length n .

Quantum invariants — natural $U_v(\mathfrak{sl}_2)$ -module

Quantum numbers: $[a] := \frac{v^a - v^{-a}}{v - v^{-1}}$, $a \in \mathbb{Z}$

V — the “natural” $U_v(\mathfrak{sl}_2)$ -module

Basis: w_0 and w_1

Action:

$$\begin{aligned} Ew_k &= [k+1]w_{k+1}, & Fw_k &= -[n-k+1]w_{k-1}, \\ K^{\pm 1}w_k &= -v^{\pm(2k-n)}w_k \end{aligned}$$

Notation: $w_{i_1} \otimes w_{i_2} \otimes \cdots \otimes w_{i_k}$ denoted by $i_1 i_2 \dots i_k$

Consequence: Basis in $V^{\otimes n}$ consists of 0-1-sequences of length n .

Quantum invariants — natural $U_v(\mathfrak{sl}_2)$ -module

Quantum numbers: $[a] := \frac{v^a - v^{-a}}{v - v^{-1}}$, $a \in \mathbb{Z}$

V — the “natural” $U_v(\mathfrak{sl}_2)$ -module

Basis: w_0 and w_1

Action:

$$\begin{aligned} Ew_k &= [k+1]w_{k+1}, & Fw_k &= -[n-k+1]w_{k-1}, \\ K^{\pm 1}w_k &= -v^{\pm(2k-n)}w_k \end{aligned}$$

Notation: $w_{i_1} \otimes w_{i_2} \otimes \cdots \otimes w_{i_k}$ denoted by $i_1 i_2 \dots i_k$

Consequence: Basis in $V^{\otimes n}$ consists of 0-1-sequences of length n .

Quantum invariants — natural $U_v(\mathfrak{sl}_2)$ -module

Quantum numbers: $[a] := \frac{v^a - v^{-a}}{v - v^{-1}}$, $a \in \mathbb{Z}$

V — the “natural” $U_v(\mathfrak{sl}_2)$ -module

Basis: w_0 and w_1

Action:

$$\begin{aligned} Ew_k &= [k+1]w_{k+1}, & Fw_k &= -[n-k+1]w_{k-1}, \\ K^{\pm 1}w_k &= -v^{\pm(2k-n)}w_k \end{aligned}$$

Notation: $w_{i_1} \otimes w_{i_2} \otimes \cdots \otimes w_{i_k}$ denoted by $i_1 i_2 \dots i_k$

Consequence: Basis in $V^{\otimes n}$ consists of 0-1-sequences of length n .

Quantum invariants — natural $U_v(\mathfrak{sl}_2)$ -module

Quantum numbers: $[a] := \frac{v^a - v^{-a}}{v - v^{-1}}$, $a \in \mathbb{Z}$

V — the “natural” $U_v(\mathfrak{sl}_2)$ -module

Basis: w_0 and w_1

Action:

$$\begin{aligned} Ew_k &= [k+1]w_{k+1}, & Fw_k &= -[n-k+1]w_{k-1}, \\ K^{\pm 1}w_k &= -v^{\pm(2k-n)}w_k \end{aligned}$$

Notation: $w_{i_1} \otimes w_{i_2} \otimes \cdots \otimes w_{i_k}$ denoted by $i_1 i_2 \dots i_k$

Consequence: Basis in $V^{\otimes n}$ consists of 0-1-sequences of length n .

Quantum invariants — natural $U_v(\mathfrak{sl}_2)$ -module

Quantum numbers: $[a] := \frac{v^a - v^{-a}}{v - v^{-1}}$, $a \in \mathbb{Z}$

V — the “natural” $U_v(\mathfrak{sl}_2)$ -module

Basis: w_0 and w_1

Action:

$$\begin{aligned} Ew_k &= [k+1]w_{k+1}, & Fw_k &= -[n-k+1]w_{k-1}, \\ K^{\pm 1}w_k &= -v^{\pm(2k-n)}w_k \end{aligned}$$

Notation: $w_{i_1} \otimes w_{i_2} \otimes \cdots \otimes w_{i_k}$ denoted by $i_1 i_2 \dots i_k$

Consequence: Basis in $V^{\otimes n}$ consists of 0-1-sequences of length n .

Quantum invariants — natural $U_v(\mathfrak{sl}_2)$ -module

Quantum numbers: $[a] := \frac{v^a - v^{-a}}{v - v^{-1}}$, $a \in \mathbb{Z}$

V — the “natural” $U_v(\mathfrak{sl}_2)$ -module

Basis: w_0 and w_1

Action:

$$\begin{aligned} Ew_k &= [k+1]w_{k+1}, & Fw_k &= -[n-k+1]w_{k-1}, \\ K^{\pm 1}w_k &= -v^{\pm(2k-n)}w_k \end{aligned}$$

Notation: $w_{i_1} \otimes w_{i_2} \otimes \cdots \otimes w_{i_k}$ denoted by $i_1 i_2 \dots i_k$

Consequence: Basis in $V^{\otimes n}$ consists of 0-1-sequences of length n .

Quantum invariants — natural $U_v(\mathfrak{sl}_2)$ -module

Quantum numbers: $[a] := \frac{v^a - v^{-a}}{v - v^{-1}}$, $a \in \mathbb{Z}$

V — the “natural” $U_v(\mathfrak{sl}_2)$ -module

Basis: w_0 and w_1

Action:

$$\begin{aligned} Ew_k &= [k+1]w_{k+1}, & Fw_k &= -[n-k+1]w_{k-1}, \\ K^{\pm 1}w_k &= -v^{\pm(2k-n)}w_k \end{aligned}$$

Notation: $w_{i_1} \otimes w_{i_2} \otimes \cdots \otimes w_{i_k}$ denoted by $i_1 i_2 \dots i_k$

Consequence: Basis in $V^{\otimes n}$ consists of 0-1-sequences of length n .

Quantum invariants — action of tangles

Definition. The functor $F : \mathbf{Tang} \rightarrow U_v(\mathfrak{sl}_2)\text{-mod}$ is given by:

$U : \mathbb{C}(v) \rightarrow \hat{\mathcal{V}}_1^{\otimes 2}$ is given by:

$$1 \mapsto 01 + v10.$$

$\cap : \hat{\mathcal{V}}_1^{\otimes 2} \rightarrow \mathbb{C}(v)$ is given by:

$$00 \mapsto 0, \quad 11 \mapsto 0, \quad 01 \mapsto v^{-1}, \quad 10 \mapsto 1.$$

right crossing: $\hat{\mathcal{V}}_1^{\otimes 2} \rightarrow \hat{\mathcal{V}}_1^{\otimes 2}$ is given by:

$$00 \mapsto -v00, \quad 11 \mapsto -v11, \quad 01 \mapsto 10 + (v^{-1} - v)01, \quad 10 \mapsto 01.$$

left crossing: $\hat{\mathcal{V}}_1^{\otimes 2} \rightarrow \hat{\mathcal{V}}_1^{\otimes 2}$ is given by:

$$00 \mapsto -v^{-1}00, \quad 11 \mapsto -v^{-1}11, \quad 01 \mapsto 10, \quad 10 \mapsto 01 + (v - v^{-1})10.$$

Quantum invariants — action of tangles

Definition. The functor $F : \mathbf{Tang} \rightarrow U_v(\mathfrak{sl}_2)\text{-mod}$ is given by:

$U : \mathbb{C}(v) \rightarrow \hat{\mathcal{V}}_1^{\otimes 2}$ is given by:

$$1 \mapsto 01 + v10.$$

$\cap : \hat{\mathcal{V}}_1^{\otimes 2} \rightarrow \mathbb{C}(v)$ is given by:

$$00 \mapsto 0, \quad 11 \mapsto 0, \quad 01 \mapsto v^{-1}, \quad 10 \mapsto 1.$$

right crossing: $\hat{\mathcal{V}}_1^{\otimes 2} \rightarrow \hat{\mathcal{V}}_1^{\otimes 2}$ is given by:

$$00 \mapsto -v00, \quad 11 \mapsto -v11, \quad 01 \mapsto 10 + (v^{-1} - v)01, \quad 10 \mapsto 01.$$

left crossing: $\hat{\mathcal{V}}_1^{\otimes 2} \rightarrow \hat{\mathcal{V}}_1^{\otimes 2}$ is given by:

$$00 \mapsto -v^{-1}00, \quad 11 \mapsto -v^{-1}11, \quad 01 \mapsto 10, \quad 10 \mapsto 01 + (v - v^{-1})10.$$

Quantum invariants — action of tangles

Definition. The functor $F : \mathbf{Tang} \rightarrow U_v(\mathfrak{sl}_2)\text{-mod}$ is given by:

$U : \mathbb{C}(v) \rightarrow \hat{\mathcal{V}}_1^{\otimes 2}$ is given by:

$$1 \mapsto 01 + v10.$$

$\cap : \hat{\mathcal{V}}_1^{\otimes 2} \rightarrow \mathbb{C}(v)$ is given by:

$$00 \mapsto 0, \quad 11 \mapsto 0, \quad 01 \mapsto v^{-1}, \quad 10 \mapsto 1.$$

right crossing: $\hat{\mathcal{V}}_1^{\otimes 2} \rightarrow \hat{\mathcal{V}}_1^{\otimes 2}$ is given by:

$$00 \mapsto -v00, \quad 11 \mapsto -v11, \quad 01 \mapsto 10 + (v^{-1} - v)01, \quad 10 \mapsto 01.$$

left crossing: $\hat{\mathcal{V}}_1^{\otimes 2} \rightarrow \hat{\mathcal{V}}_1^{\otimes 2}$ is given by:

$$00 \mapsto -v^{-1}00, \quad 11 \mapsto -v^{-1}11, \quad 01 \mapsto 10, \quad 10 \mapsto 01 + (v - v^{-1})10.$$

Quantum invariants — action of tangles

Definition. The functor $F : \mathbf{Tang} \rightarrow U_v(\mathfrak{sl}_2)\text{-mod}$ is given by:

$U : \mathbb{C}(v) \rightarrow \hat{\mathcal{V}}_1^{\otimes 2}$ is given by:

$$1 \mapsto 01 + v10.$$

$\cap : \hat{\mathcal{V}}_1^{\otimes 2} \rightarrow \mathbb{C}(v)$ is given by:

$$00 \mapsto 0, \quad 11 \mapsto 0, \quad 01 \mapsto v^{-1}, \quad 10 \mapsto 1.$$

right crossing: $\hat{\mathcal{V}}_1^{\otimes 2} \rightarrow \hat{\mathcal{V}}_1^{\otimes 2}$ is given by:

$$00 \mapsto -v00, \quad 11 \mapsto -v11, \quad 01 \mapsto 10 + (v^{-1} - v)01, \quad 10 \mapsto 01.$$

left crossing: $\hat{\mathcal{V}}_1^{\otimes 2} \rightarrow \hat{\mathcal{V}}_1^{\otimes 2}$ is given by:

$$00 \mapsto -v^{-1}00, \quad 11 \mapsto -v^{-1}11, \quad 01 \mapsto 10, \quad 10 \mapsto 01 + (v - v^{-1})10.$$

Quantum invariants — action of tangles

Definition. The functor $F : \mathbf{Tang} \rightarrow U_v(\mathfrak{sl}_2)\text{-mod}$ is given by:

$U : \mathbb{C}(v) \rightarrow \hat{\mathcal{V}}_1^{\otimes 2}$ is given by:

$$1 \mapsto 01 + v10.$$

$\cap : \hat{\mathcal{V}}_1^{\otimes 2} \rightarrow \mathbb{C}(v)$ is given by:

$$00 \mapsto 0, \quad 11 \mapsto 0, \quad 01 \mapsto v^{-1}, \quad 10 \mapsto 1.$$

right crossing: $\hat{\mathcal{V}}_1^{\otimes 2} \rightarrow \hat{\mathcal{V}}_1^{\otimes 2}$ is given by:

$$00 \mapsto -v00, \quad 11 \mapsto -v11, \quad 01 \mapsto 10 + (v^{-1} - v)01, \quad 10 \mapsto 01.$$

left crossing: $\hat{\mathcal{V}}_1^{\otimes 2} \rightarrow \hat{\mathcal{V}}_1^{\otimes 2}$ is given by:

$$00 \mapsto -v^{-1}00, \quad 11 \mapsto -v^{-1}11, \quad 01 \mapsto 10, \quad 10 \mapsto 01 + (v - v^{-1})10.$$

Quantum invariants — action of tangles

Definition. The functor $F : \mathbf{Tang} \rightarrow U_v(\mathfrak{sl}_2)\text{-mod}$ is given by:

$U : \mathbb{C}(v) \rightarrow \hat{\mathcal{V}}_1^{\otimes 2}$ is given by:

$$1 \mapsto 01 + v10.$$

$\cap : \hat{\mathcal{V}}_1^{\otimes 2} \rightarrow \mathbb{C}(v)$ is given by:

$$00 \mapsto 0, \quad 11 \mapsto 0, \quad 01 \mapsto v^{-1}, \quad 10 \mapsto 1.$$

right crossing: $\hat{\mathcal{V}}_1^{\otimes 2} \rightarrow \hat{\mathcal{V}}_1^{\otimes 2}$ is given by:

$$00 \mapsto -v00, \quad 11 \mapsto -v11, \quad 01 \mapsto 10 + (v^{-1} - v)01, \quad 10 \mapsto 01.$$

left crossing: $\hat{\mathcal{V}}_1^{\otimes 2} \rightarrow \hat{\mathcal{V}}_1^{\otimes 2}$ is given by:

$$00 \mapsto -v^{-1}00, \quad 11 \mapsto -v^{-1}11, \quad 01 \mapsto 10, \quad 10 \mapsto 01 + (v - v^{-1})10.$$

Quantum invariants — action of tangles

Definition. The functor $F : \mathbf{Tang} \rightarrow U_v(\mathfrak{sl}_2)\text{-mod}$ is given by:

$U : \mathbb{C}(v) \rightarrow \hat{\mathcal{V}}_1^{\otimes 2}$ is given by:

$$1 \mapsto 01 + v10.$$

$\cap : \hat{\mathcal{V}}_1^{\otimes 2} \rightarrow \mathbb{C}(v)$ is given by:

$$00 \mapsto 0, \quad 11 \mapsto 0, \quad 01 \mapsto v^{-1}, \quad 10 \mapsto 1.$$

right crossing: $\hat{\mathcal{V}}_1^{\otimes 2} \rightarrow \hat{\mathcal{V}}_1^{\otimes 2}$ is given by:

$$00 \mapsto -v00, \quad 11 \mapsto -v11, \quad 01 \mapsto 10 + (v^{-1} - v)01, \quad 10 \mapsto 01.$$

left crossing: $\hat{\mathcal{V}}_1^{\otimes 2} \rightarrow \hat{\mathcal{V}}_1^{\otimes 2}$ is given by:

$$00 \mapsto -v^{-1}00, \quad 11 \mapsto -v^{-1}11, \quad 01 \mapsto 10, \quad 10 \mapsto 01 + (v - v^{-1})10.$$

Quantum $U_v(\mathfrak{sl}_2)$ -invariants and Jones polynomial

Theorem. [Reshetikhin-Turaev]

Let L be an oriented link. Then

the polynomials $F(T_L)(1)$ and $\hat{J}(L)$ coincide.

Quantum $U_v(\mathfrak{sl}_2)$ -invariants and Jones polynomial

Theorem. [Reshetikhin-Turaev]

Let L be an oriented link. Then

the polynomials $F(T_L)(1)$ and $\hat{J}(L)$ coincide.

Quantum $U_v(\mathfrak{sl}_2)$ -invariants and Jones polynomial

Theorem. [Reshetikhin-Turaev]

Let L be an oriented link. Then

the polynomials $F(T_L)(1)$ and $\hat{J}(L)$ coincide.

Quantum $U_v(\mathfrak{sl}_2)$ -invariants and Jones polynomial

Theorem. [Reshetikhin-Turaev]

Let L be an oriented link. Then

the polynomials $F(T_L)(1)$ and $\hat{J}(L)$ coincide.

Quantum $U_v(\mathfrak{sl}_2)$ -invariants and Jones polynomial

Theorem. [Reshetikhin-Turaev]

Let L be an oriented link. Then

the polynomials $F(T_L)(1)$ and $\hat{J}(L)$ coincide.

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — the idea

Cat — category of categories

Idea: Construct a functor from **Tang** to **Cat**?

Results in: Khovanov's "functor-valued invariants of tangles"

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — the idea

Cat — category of categories

Idea: Construct a functor from **Tang** to **Cat**?

Results in: Khovanov's “functor-valued invariants of tangles”

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — the idea

Cat — category of categories

Idea: Construct a functor from **Tang** to **Cat**?

Results in: Khovanov's “functor-valued invariants of tangles”

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — the idea

Cat — category of categories

Idea: Construct a functor from **Tang** to **Cat**?

Results in: Khovanov's “functor-valued invariants of tangles”

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — the idea

Cat — category of categories

Idea: Construct a functor from **Tang** to **Cat**?

Results in: Khovanov's “**functor-valued invariants of tangles**”

Approach via category \mathcal{O}

\mathfrak{gl}_n — reductive Lie algebra over \mathbb{C}

$\mathfrak{gl}_n = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$ — standard triangular decomposition

\mathcal{O} — BGG category \mathcal{O}

S_n — the *Weyl group* of \mathfrak{gl}_n

Fact: S_n acts on \mathfrak{h}^* in the natural way

$M(\lambda)$ — *Verma* module with highest weight $\lambda - \rho$

Approach via category \mathcal{O}

\mathfrak{gl}_n — reductive Lie algebra over \mathbb{C}

$\mathfrak{gl}_n = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$ — standard triangular decomposition

\mathcal{O} — BGG category \mathcal{O}

S_n — the Weyl group of \mathfrak{gl}_n

Fact: S_n acts on \mathfrak{h}^* in the natural way

$M(\lambda)$ — Verma module with highest weight $\lambda - \rho$

Approach via category \mathcal{O}

\mathfrak{gl}_n — reductive Lie algebra over \mathbb{C}

$\mathfrak{gl}_n = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$ — standard triangular decomposition

\mathcal{O} — BGG category \mathcal{O}

S_n — the Weyl group of \mathfrak{gl}_n

Fact: S_n acts on \mathfrak{h}^* in the natural way

$M(\lambda)$ — Verma module with highest weight $\lambda - \rho$

Approach via category \mathcal{O}

\mathfrak{gl}_n — reductive Lie algebra over \mathbb{C}

$\mathfrak{gl}_n = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$ — standard triangular decomposition

\mathcal{O} — BGG category \mathcal{O}

S_n — the Weyl group of \mathfrak{gl}_n

Fact: S_n acts on \mathfrak{h}^* in the natural way

$M(\lambda)$ — Verma module with highest weight $\lambda - \rho$

Approach via category \mathcal{O}

\mathfrak{gl}_n — reductive Lie algebra over \mathbb{C}

$\mathfrak{gl}_n = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$ — standard triangular decomposition

\mathcal{O} — BGG category \mathcal{O}

S_n — the **Weyl group** of \mathfrak{gl}_n

Fact: S_n acts on \mathfrak{h}^* in the natural way

$M(\lambda)$ — **Verma** module with highest weight $\lambda - \rho$

Approach via category \mathcal{O}

\mathfrak{gl}_n — reductive Lie algebra over \mathbb{C}

$\mathfrak{gl}_n = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$ — standard triangular decomposition

\mathcal{O} — BGG category \mathcal{O}

S_n — the **Weyl group** of \mathfrak{gl}_n

Fact: S_n acts on \mathfrak{h}^* in the natural way

$M(\lambda)$ — **Verma** module with highest weight $\lambda - \rho$

Approach via category \mathcal{O}

\mathfrak{gl}_n — reductive Lie algebra over \mathbb{C}

$\mathfrak{gl}_n = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$ — standard triangular decomposition

\mathcal{O} — BGG category \mathcal{O}

S_n — the **Weyl group** of \mathfrak{gl}_n

Fact: S_n acts on \mathfrak{h}^* in the natural way

$M(\lambda)$ — **Verma** module with highest weight $\lambda - \rho$

Approach via category \mathcal{O}

\mathfrak{gl}_n — reductive Lie algebra over \mathbb{C}

$\mathfrak{gl}_n = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$ — standard triangular decomposition

\mathcal{O} — BGG category \mathcal{O}

S_n — the **Weyl group** of \mathfrak{gl}_n

Fact: S_n acts on \mathfrak{h}^* in the natural way

$M(\lambda)$ — **Verma** module with highest weight $\lambda - \rho$

Blocks in \mathcal{O}

\mathcal{O}_0 — the principal block of \mathcal{O}

$k \in \{0, 1, 2, \dots, n\}$

$S_k \times S_{n-k} \subset S_N$ — maximal Young subgroup

\mathfrak{p}_k — corresponding parabolic subalgebra

$\mathcal{O}_0^{(k, n-k)}$ — parabolic subcategory of locally \mathfrak{p}_k -finite modules

Definition: $\mathcal{C}_n := \bigoplus_{k=0}^n \mathcal{O}_0^{(k, n-k)}$

Fact: \mathcal{C}_n has 2^n simple objects up to isomorphism.

Definition: $\text{Gr}(\mathcal{C}_n)$ is the Grothendieck group of \mathcal{C}_n

Blocks in \mathcal{O}

\mathcal{O}_0 — the **principal block** of \mathcal{O}

$$k \in \{0, 1, 2, \dots, n\}$$

$S_k \times S_{n-k} \subset S_N$ — maximal **Young** subgroup

\mathfrak{p}_k — corresponding **parabolic** subalgebra

$\mathcal{O}_0^{(k, n-k)}$ — **parabolic** subcategory of locally \mathfrak{p}_k -finite modules

Definition: $\mathcal{C}_n := \bigoplus_{k=0}^n \mathcal{O}_0^{(k, n-k)}$

Fact: \mathcal{C}_n has 2^n simple objects up to isomorphism.

Definition: $\text{Gr}(\mathcal{C}_n)$ is the **Grothendieck group** of \mathcal{C}_n

Blocks in \mathcal{O}

\mathcal{O}_0 — the **principal block** of \mathcal{O}

$k \in \{0, 1, 2, \dots, n\}$

$S_k \times S_{n-k} \subset S_N$ — maximal **Young** subgroup

\mathfrak{p}_k — corresponding **parabolic** subalgebra

$\mathcal{O}_0^{(k, n-k)}$ — **parabolic** subcategory of locally \mathfrak{p}_k -finite modules

Definition: $\mathcal{C}_n := \bigoplus_{k=0}^n \mathcal{O}_0^{(k, n-k)}$

Fact: \mathcal{C}_n has 2^n simple objects up to isomorphism.

Definition: $\text{Gr}(\mathcal{C}_n)$ is the **Grothendieck group** of \mathcal{C}_n

Blocks in \mathcal{O}

\mathcal{O}_0 — the **principal block** of \mathcal{O}

$k \in \{0, 1, 2, \dots, n\}$

$S_k \times S_{n-k} \subset S_N$ — maximal **Young** subgroup

\mathfrak{p}_k — corresponding **parabolic** subalgebra

$\mathcal{O}_0^{(k, n-k)}$ — **parabolic** subcategory of locally \mathfrak{p}_k -finite modules

Definition: $\mathcal{C}_n := \bigoplus_{k=0}^n \mathcal{O}_0^{(k, n-k)}$

Fact: \mathcal{C}_n has 2^n simple objects up to isomorphism.

Definition: $\text{Gr}(\mathcal{C}_n)$ is the **Grothendieck group** of \mathcal{C}_n

Blocks in \mathcal{O}

\mathcal{O}_0 — the **principal block** of \mathcal{O}

$$k \in \{0, 1, 2, \dots, n\}$$

$S_k \times S_{n-k} \subset S_N$ — maximal **Young** subgroup

\mathfrak{p}_k — corresponding **parabolic** subalgebra

$\mathcal{O}_0^{(k, n-k)}$ — **parabolic** subcategory of locally \mathfrak{p}_k -finite modules

Definition:
$$\mathcal{C}_n := \bigoplus_{k=0}^n \mathcal{O}_0^{(k, n-k)}$$

Fact: \mathcal{C}_n has 2^n simple objects up to isomorphism.

Definition: $\text{Gr}(\mathcal{C}_n)$ is the **Grothendieck group** of \mathcal{C}_n

Blocks in \mathcal{O}

\mathcal{O}_0 — the **principal block** of \mathcal{O}

$$k \in \{0, 1, 2, \dots, n\}$$

$S_k \times S_{n-k} \subset S_N$ — maximal **Young** subgroup

\mathfrak{p}_k — corresponding **parabolic** subalgebra

$\mathcal{O}_0^{(k, n-k)}$ — **parabolic** subcategory of locally \mathfrak{p}_k -finite modules

Definition:
$$\mathcal{C}_n := \bigoplus_{k=0}^n \mathcal{O}_0^{(k, n-k)}$$

Fact: \mathcal{C}_n has 2^n simple objects up to isomorphism.

Definition: $\text{Gr}(\mathcal{C}_n)$ is the **Grothendieck group** of \mathcal{C}_n

Blocks in \mathcal{O}

\mathcal{O}_0 — the **principal block** of \mathcal{O}

$k \in \{0, 1, 2, \dots, n\}$

$S_k \times S_{n-k} \subset S_N$ — maximal **Young** subgroup

\mathfrak{p}_k — corresponding **parabolic** subalgebra

$\mathcal{O}_0^{(k, n-k)}$ — **parabolic** subcategory of locally \mathfrak{p}_k -finite modules

Definition: $\mathcal{C}_n := \bigoplus_{k=0}^n \mathcal{O}_0^{(k, n-k)}$

Fact: \mathcal{C}_n has 2^n simple objects up to isomorphism.

Definition: $\text{Gr}(\mathcal{C}_n)$ is the **Grothendieck group** of \mathcal{C}_n

Blocks in \mathcal{O}

\mathcal{O}_0 — the **principal block** of \mathcal{O}

$$k \in \{0, 1, 2, \dots, n\}$$

$S_k \times S_{n-k} \subset S_N$ — maximal **Young** subgroup

\mathfrak{p}_k — corresponding **parabolic** subalgebra

$\mathcal{O}_0^{(k, n-k)}$ — **parabolic** subcategory of locally \mathfrak{p}_k -finite modules

Definition:
$$\mathcal{C}_n := \bigoplus_{k=0}^n \mathcal{O}_0^{(k, n-k)}$$

Fact: \mathcal{C}_n has 2^n simple objects up to isomorphism.

Definition: $\text{Gr}(\mathcal{C}_n)$ is the **Grothendieck group** of \mathcal{C}_n

Blocks in \mathcal{O}

\mathcal{O}_0 — the **principal block** of \mathcal{O}

$k \in \{0, 1, 2, \dots, n\}$

$S_k \times S_{n-k} \subset S_N$ — maximal **Young** subgroup

\mathfrak{p}_k — corresponding **parabolic** subalgebra

$\mathcal{O}_0^{(k, n-k)}$ — **parabolic** subcategory of locally \mathfrak{p}_k -finite modules

Definition: $\mathcal{C}_n := \bigoplus_{k=0}^n \mathcal{O}_0^{(k, n-k)}$

Fact: \mathcal{C}_n has 2^n simple objects up to isomorphism.

Definition: $\text{Gr}(\mathcal{C}_n)$ is the **Grothendieck group** of \mathcal{C}_n

Blocks in \mathcal{O}

\mathcal{O}_0 — the **principal block** of \mathcal{O}

$k \in \{0, 1, 2, \dots, n\}$

$S_k \times S_{n-k} \subset S_N$ — maximal **Young** subgroup

\mathfrak{p}_k — corresponding **parabolic** subalgebra

$\mathcal{O}_0^{(k, n-k)}$ — **parabolic** subcategory of locally \mathfrak{p}_k -finite modules

Definition: $\mathcal{C}_n := \bigoplus_{k=0}^n \mathcal{O}_0^{(k, n-k)}$

Fact: \mathcal{C}_n has 2^n simple objects up to isomorphism.

Definition: $\text{Gr}(\mathcal{C}_n)$ is the **Grothendieck group** of \mathcal{C}_n

Categorification of $V^{\otimes n}$ for $v = 1$

Observation: $\dim V^{\otimes n} = \text{rank}(\text{Gr}(\mathcal{C}_n))$

$\mathfrak{p} \subset \mathfrak{q}$ — parabolic subalgebras

$I_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{q}} \subset \mathcal{O}^{\mathfrak{p}}$ — natural inclusion

$Z_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{p}} \subset \mathcal{O}^{\mathfrak{q}}$ — adjoint Zuckerman functors

Note: $Z_{(\mathfrak{p}, \mathfrak{q})}$ is only right exact

Action: $E : \mathcal{D}^b(\mathcal{O}^{(k, n-k)}) \xrightarrow{I} \mathcal{D}^b(\mathcal{O}^{(k, 1, n-k-1)}) \xrightarrow{\mathcal{L}Z} \mathcal{D}^b(\mathcal{O}^{(k+1, n-k-1)})$

Action: F — adjoint to E

Theorem. [Bernstein-Frenkel-Khovanov] This categorifies $V^{\otimes n}$ for $v = 1$.

Meaning: Taking the Grothendieck group results in $V^{\otimes n}$

Categorification of $V^{\otimes n}$ for $v = 1$

Observation: $\dim V^{\otimes n} = \text{rank}(\text{Gr}(\mathcal{C}_n))$

$\mathfrak{p} \subset \mathfrak{q}$ — parabolic subalgebras

$I_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{q}} \subset \mathcal{O}^{\mathfrak{p}}$ — natural inclusion

$Z_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{p}} \subset \mathcal{O}^{\mathfrak{q}}$ — adjoint Zuckerman functors

Note: $Z_{(\mathfrak{p}, \mathfrak{q})}$ is only right exact

Action: $E : \mathcal{D}^b(\mathcal{O}^{(k, n-k)}) \xrightarrow{I} \mathcal{D}^b(\mathcal{O}^{(k, 1, n-k-1)}) \xrightarrow{\mathcal{L}Z} \mathcal{D}^b(\mathcal{O}^{(k+1, n-k-1)})$

Action: F — adjoint to E

Theorem. [Bernstein-Frenkel-Khovanov] This categorifies $V^{\otimes n}$ for $v = 1$.

Meaning: Taking the Grothendieck group results in $V^{\otimes n}$

Categorification of $V^{\otimes n}$ for $v = 1$

Observation: $\dim V^{\otimes n} = \text{rank}(\text{Gr}(\mathcal{C}_n))$

$\mathfrak{p} \subset \mathfrak{q}$ — parabolic subalgebras

$I_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{q}} \subset \mathcal{O}^{\mathfrak{p}}$ — natural inclusion

$Z_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{p}} \subset \mathcal{O}^{\mathfrak{q}}$ — adjoint Zuckerman functors

Note: $Z_{(\mathfrak{p}, \mathfrak{q})}$ is only right exact

Action: $E : \mathcal{D}^b(\mathcal{O}^{(k, n-k)}) \xrightarrow{I} \mathcal{D}^b(\mathcal{O}^{(k, 1, n-k-1)}) \xrightarrow{\mathcal{L}Z} \mathcal{D}^b(\mathcal{O}^{(k+1, n-k-1)})$

Action: F — adjoint to E

Theorem. [Bernstein-Frenkel-Khovanov] This categorifies $V^{\otimes n}$ for $v = 1$.

Meaning: Taking the Grothendieck group results in $V^{\otimes n}$

Categorification of $V^{\otimes n}$ for $v = 1$

Observation: $\dim V^{\otimes n} = \text{rank}(\text{Gr}(\mathcal{C}_n))$

$\mathfrak{p} \subset \mathfrak{q}$ — parabolic subalgebras

$I_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{q}} \subset \mathcal{O}^{\mathfrak{p}}$ — natural **inclusion**

$Z_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{p}} \subset \mathcal{O}^{\mathfrak{q}}$ — adjoint **Zuckerman** functors

Note: $Z_{(\mathfrak{p}, \mathfrak{q})}$ is only **right exact**

Action: $E : \mathcal{D}^b(\mathcal{O}^{(k, n-k)}) \xrightarrow{I} \mathcal{D}^b(\mathcal{O}^{(k, 1, n-k-1)}) \xrightarrow{\mathcal{L}Z} \mathcal{D}^b(\mathcal{O}^{(k+1, n-k-1)})$

Action: F — adjoint to E

Theorem. [Bernstein-Frenkel-Khovanov] This **categorifies** $V^{\otimes n}$ for $v = 1$.

Meaning: Taking the Grothendieck group results in $V^{\otimes n}$

Categorification of $V^{\otimes n}$ for $v = 1$

Observation: $\dim V^{\otimes n} = \text{rank}(\text{Gr}(\mathcal{C}_n))$

$\mathfrak{p} \subset \mathfrak{q}$ — parabolic subalgebras

$I_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{q}} \subset \mathcal{O}^{\mathfrak{p}}$ — natural **inclusion**

$Z_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{p}} \subset \mathcal{O}^{\mathfrak{q}}$ — adjoint **Zuckerman** functors

Note: $Z_{(\mathfrak{p}, \mathfrak{q})}$ is only **right exact**

Action: $E : \mathcal{D}^b(\mathcal{O}^{(k, n-k)}) \xrightarrow{I} \mathcal{D}^b(\mathcal{O}^{(k, 1, n-k-1)}) \xrightarrow{\mathcal{L}Z} \mathcal{D}^b(\mathcal{O}^{(k+1, n-k-1)})$

Action: F — adjoint to E

Theorem. [Bernstein-Frenkel-Khovanov] This **categorifies** $V^{\otimes n}$ for $v = 1$.

Meaning: Taking the Grothendieck group results in $V^{\otimes n}$

Categorification of $V^{\otimes n}$ for $v = 1$

Observation: $\dim V^{\otimes n} = \text{rank}(\text{Gr}(\mathcal{C}_n))$

$\mathfrak{p} \subset \mathfrak{q}$ — parabolic subalgebras

$I_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{q}} \subset \mathcal{O}^{\mathfrak{p}}$ — natural **inclusion**

$Z_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{p}} \subset \mathcal{O}^{\mathfrak{q}}$ — adjoint **Zuckerman** functors

Note: $Z_{(\mathfrak{p}, \mathfrak{q})}$ is only **right** exact

Action: $E : \mathcal{D}^b(\mathcal{O}^{(k, n-k)}) \xrightarrow{I} \mathcal{D}^b(\mathcal{O}^{(k, 1, n-k-1)}) \xrightarrow{\mathcal{L}Z} \mathcal{D}^b(\mathcal{O}^{(k+1, n-k-1)})$

Action: F — adjoint to E

Theorem. [Bernstein-Frenkel-Khovanov] This categorifies $V^{\otimes n}$ for $v = 1$.

Meaning: Taking the Grothendieck group results in $V^{\otimes n}$

Categorification of $V^{\otimes n}$ for $v = 1$

Observation: $\dim V^{\otimes n} = \text{rank}(\text{Gr}(\mathcal{C}_n))$

$\mathfrak{p} \subset \mathfrak{q}$ — parabolic subalgebras

$I_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{q}} \subset \mathcal{O}^{\mathfrak{p}}$ — natural **inclusion**

$Z_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{p}} \subset \mathcal{O}^{\mathfrak{q}}$ — adjoint **Zuckerman** functors

Note: $Z_{(\mathfrak{p}, \mathfrak{q})}$ is only **right** exact

Action: $E : \mathcal{D}^b(\mathcal{O}^{(k, n-k)}) \xrightarrow{I} \mathcal{D}^b(\mathcal{O}^{(k, 1, n-k-1)}) \xrightarrow{\mathcal{LZ}} \mathcal{D}^b(\mathcal{O}^{(k+1, n-k-1)})$

Action: F — adjoint to E

Theorem. [Bernstein-Frenkel-Khovanov] This **categorifies** $V^{\otimes n}$ for $v = 1$.

Meaning: Taking the Grothendieck group results in $V^{\otimes n}$

Categorification of $V^{\otimes n}$ for $v = 1$

Observation: $\dim V^{\otimes n} = \text{rank}(\text{Gr}(\mathcal{C}_n))$

$\mathfrak{p} \subset \mathfrak{q}$ — parabolic subalgebras

$I_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{q}} \subset \mathcal{O}^{\mathfrak{p}}$ — natural **inclusion**

$Z_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{p}} \subset \mathcal{O}^{\mathfrak{q}}$ — adjoint **Zuckerman** functors

Note: $Z_{(\mathfrak{p}, \mathfrak{q})}$ is only **right** exact

Action: $E : \mathcal{D}^b(\mathcal{O}^{(k, n-k)}) \xrightarrow{I} \mathcal{D}^b(\mathcal{O}^{(k, 1, n-k-1)}) \xrightarrow{\mathcal{L}Z} \mathcal{D}^b(\mathcal{O}^{(k+1, n-k-1)})$

Action: F — adjoint to E

Theorem. [Bernstein-Frenkel-Khovanov] This **categorifies** $V^{\otimes n}$ for $v = 1$.

Meaning: Taking the Grothendieck group results in $V^{\otimes n}$

Categorification of $V^{\otimes n}$ for $v = 1$

Observation: $\dim V^{\otimes n} = \text{rank}(\text{Gr}(\mathcal{C}_n))$

$\mathfrak{p} \subset \mathfrak{q}$ — parabolic subalgebras

$I_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{q}} \subset \mathcal{O}^{\mathfrak{p}}$ — natural **inclusion**

$Z_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{p}} \subset \mathcal{O}^{\mathfrak{q}}$ — adjoint **Zuckerman** functors

Note: $Z_{(\mathfrak{p}, \mathfrak{q})}$ is only **right** exact

Action: $E : \mathcal{D}^b(\mathcal{O}^{(k, n-k)}) \xrightarrow{I} \mathcal{D}^b(\mathcal{O}^{(k, 1, n-k-1)}) \xrightarrow{\mathcal{LZ}} \mathcal{D}^b(\mathcal{O}^{(k+1, n-k-1)})$

Action: F — adjoint to E

Theorem. [Bernstein-Frenkel-Khovanov] This **categorifies** $V^{\otimes n}$ for $v = 1$.

Meaning: Taking the Grothendieck group results in $V^{\otimes n}$

Categorification of $V^{\otimes n}$ for $v = 1$

Observation: $\dim V^{\otimes n} = \text{rank}(\text{Gr}(\mathcal{C}_n))$

$\mathfrak{p} \subset \mathfrak{q}$ — parabolic subalgebras

$I_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{q}} \subset \mathcal{O}^{\mathfrak{p}}$ — natural **inclusion**

$Z_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{p}} \subset \mathcal{O}^{\mathfrak{q}}$ — adjoint **Zuckerman** functors

Note: $Z_{(\mathfrak{p}, \mathfrak{q})}$ is only **right** exact

Action: $E : \mathcal{D}^b(\mathcal{O}^{(k, n-k)}) \xrightarrow{I} \mathcal{D}^b(\mathcal{O}^{(k, 1, n-k-1)}) \xrightarrow{\mathcal{L}Z} \mathcal{D}^b(\mathcal{O}^{(k+1, n-k-1)})$

Action: F — adjoint to E

Theorem. [Bernstein-Frenkel-Khovanov] This **categorifies** $V^{\otimes n}$ for $v = 1$.

Meaning: Taking the Grothendieck group results in $V^{\otimes n}$

Categorification of $V^{\otimes n}$ for $v = 1$

Observation: $\dim V^{\otimes n} = \text{rank}(\text{Gr}(\mathcal{C}_n))$

$\mathfrak{p} \subset \mathfrak{q}$ — parabolic subalgebras

$I_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{q}} \subset \mathcal{O}^{\mathfrak{p}}$ — natural **inclusion**

$Z_{(\mathfrak{p}, \mathfrak{q})} : \mathcal{O}^{\mathfrak{p}} \subset \mathcal{O}^{\mathfrak{q}}$ — adjoint **Zuckerman** functors

Note: $Z_{(\mathfrak{p}, \mathfrak{q})}$ is only **right** exact

Action: $E : \mathcal{D}^b(\mathcal{O}^{(k, n-k)}) \xrightarrow{I} \mathcal{D}^b(\mathcal{O}^{(k, 1, n-k-1)}) \xrightarrow{\mathcal{L}Z} \mathcal{D}^b(\mathcal{O}^{(k+1, n-k-1)})$

Action: F — adjoint to E

Theorem. [Bernstein-Frenkel-Khovanov] This **categorifies** $V^{\otimes n}$ for $v = 1$.

Meaning: Taking the Grothendieck group results in $V^{\otimes n}$

Categorification of v

Question: Where can we find v ?

Answer: Introduce **grading**.

Theorem. [Soergel] Each block of (parabolic) \mathcal{O} is equivalent to the category of (ungraded) modules over a finite dimensional positively graded and even Koszul algebra.

$\tilde{\mathcal{C}}_n$ — **graded version** of \mathcal{C}_n

Theorem. [Stroppel] The action of graded Zuckerman functors on $\mathcal{D}^b(\tilde{\mathcal{C}}_n)$ **categorifies** $V^{\otimes n}$

Here: v corresponds to shift of grading.

Categorification of v

Question: Where can we find v ?

Answer: Introduce **grading**.

Theorem. [Soergel] Each block of (parabolic) \mathcal{O} is equivalent to the category of (ungraded) modules over a finite dimensional **positively graded** and even Koszul algebra.

$\tilde{\mathcal{C}}_n$ — **graded version** of \mathcal{C}_n

Theorem. [Stroppel] The action of graded Zuckerman functors on $\mathcal{D}^b(\tilde{\mathcal{C}}_n)$ **categorifies** $V^{\otimes n}$

Here: v corresponds to shift of grading.

Categorification of v

Question: Where can we find v ?

Answer: Introduce **grading**.

Theorem. [Soergel] Each block of (parabolic) \mathcal{O} is equivalent to the category of (ungraded) modules over a finite dimensional positively graded and even Koszul algebra.

$\tilde{\mathcal{C}}_n$ — **graded version** of \mathcal{C}_n

Theorem. [Stroppel] The action of graded Zuckerman functors on $\mathcal{D}^b(\tilde{\mathcal{C}}_n)$ **categorifies** $V^{\otimes n}$

Here: v corresponds to shift of grading.

Categorification of v

Question: Where can we find v ?

Answer: Introduce **grading**.

Theorem. [Soergel] Each block of (parabolic) \mathcal{O} is equivalent to the category of (ungraded) modules over a finite dimensional **positively graded** and even **Koszul** algebra.

$\tilde{\mathcal{C}}_n$ — **graded version** of \mathcal{C}_n

Theorem. [Stroppel] The action of graded Zuckerman functors on $\mathcal{D}^b(\tilde{\mathcal{C}}_n)$ **categorifies** $V^{\otimes n}$

Here: v corresponds to shift of grading.

Categorification of v

Question: Where can we find v ?

Answer: Introduce **grading**.

Theorem. [Soergel] Each block of (parabolic) \mathcal{O} is equivalent to the category of (ungraded) modules over a finite dimensional **positively graded** and even **Koszul** algebra.

$\tilde{\mathcal{C}}_n$ — **graded version** of \mathcal{C}_n

Theorem. [Stroppel] The action of graded Zuckerman functors on $\mathcal{D}^b(\tilde{\mathcal{C}}_n)$ **categorifies** $V^{\otimes n}$

Here: v corresponds to shift of grading.

Categorification of v

Question: Where can we find v ?

Answer: Introduce **grading**.

Theorem. [Soergel] Each block of (parabolic) \mathcal{O} is equivalent to the category of (ungraded) modules over a finite dimensional **positively graded** and even **Koszul** algebra.

$\tilde{\mathcal{C}}_n$ — **graded version** of \mathcal{C}_n

Theorem. [Stroppel] The action of graded Zuckerman functors on $\mathcal{D}^b(\tilde{\mathcal{C}}_n)$ **categorifies** $V^{\otimes n}$

Here: v corresponds to shift of grading.

Categorification of v

Question: Where can we find v ?

Answer: Introduce **grading**.

Theorem. [Soergel] Each block of (parabolic) \mathcal{O} is equivalent to the category of (ungraded) modules over a finite dimensional **positively graded** and even **Koszul** algebra.

$\tilde{\mathcal{C}}_n$ — **graded version** of \mathcal{C}_n

Theorem. [Stroppel] The action of graded Zuckerman functors on $\mathcal{D}^b(\tilde{\mathcal{C}}_n)$ **categorifies** $V^{\otimes n}$

Here: v corresponds to **shift of grading**.

Categorification of v

Question: Where can we find v ?

Answer: Introduce **grading**.

Theorem. [Soergel] Each block of (parabolic) \mathcal{O} is equivalent to the category of (ungraded) modules over a finite dimensional **positively graded** and even **Koszul** algebra.

$\tilde{\mathcal{C}}_n$ — **graded version** of \mathcal{C}_n

Theorem. [Stroppel] The action of graded Zuckerman functors on $\mathcal{D}^b(\tilde{\mathcal{C}}_n)$ **categorifies** $V^{\otimes n}$

Here: v corresponds to **shift of grading**.

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — setup

Definition. [Bernstein-S. Gelfand] A projective functor on \mathcal{O} is a functor isomorphic to a direct summand of tensoring with a finite dimensional module.

Fact. Projective functors commute with Zuckerman functors.

Need: Categorification of $V^{\otimes m}$ for $m < n$

Use: Singular and singular-parabolic blocks of \mathcal{O}

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — setup

Definition. [Bernstein-S. Gelfand] A **projective functor** on \mathcal{O} is a functor isomorphic to a direct summand of tensoring with a finite dimensional module.

Fact. Projective functors commute with Zuckerman functors.

Need: Categorification of $V^{\otimes m}$ for $m < n$

Use: Singular and singular-parabolic blocks of \mathcal{O}

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — setup

Definition. [Bernstein-S. Gelfand] A **projective functor** on \mathcal{O} is a functor isomorphic to a direct summand of tensoring with a finite dimensional module.

Fact. Projective functors commute with Zuckerman functors.

Need: Categorification of $V^{\otimes m}$ for $m < n$

Use: Singular and singular-parabolic blocks of \mathcal{O}

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — setup

Definition. [Bernstein-S. Gelfand] A **projective functor** on \mathcal{O} is a functor isomorphic to a direct summand of tensoring with a finite dimensional module.

Fact. Projective functors commute with Zuckerman functors.

Need: Categorification of $V^{\otimes m}$ for $m < n$

Use: Singular and singular-parabolic blocks of \mathcal{O}

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — setup

Definition. [Bernstein-S. Gelfand] A **projective functor** on \mathcal{O} is a functor isomorphic to a direct summand of tensoring with a finite dimensional module.

Fact. Projective functors commute with Zuckerman functors.

Need: Categorification of $V^{\otimes m}$ for $m < n$

Use: Singular and singular-parabolic blocks of \mathcal{O}

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — setup

Definition. [Bernstein-S. Gelfand] A **projective functor** on \mathcal{O} is a functor isomorphic to a direct summand of tensoring with a finite dimensional module.

Fact. Projective functors commute with Zuckerman functors.

Need: Categorification of $V^{\otimes m}$ for $m < n$

Use: Singular and singular-parabolic blocks of \mathcal{O}

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — shuffling functors

$s \in S_n$ — simple reflection

θ_s — wall-crossing functor

Fact There are adjunctions $\theta_s \rightarrow \text{Id}$ and $\text{Id} \rightarrow \theta_s$

Definition.[Carlin] Shuffling functor $C_s := \text{Coker}(\text{Id} \rightarrow \theta_s)$ (adjoint: coshuffling)

Fact. Shuffling is right exact.

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — shuffling functors

$s \in S_n$ — simple reflection

θ_s — wall-crossing functor

Fact There are adjunctions $\theta_s \rightarrow \text{Id}$ and $\text{Id} \rightarrow \theta_s$

Definition. [Carlin] Shuffling functor $C_s := \text{Coker}(\text{Id} \rightarrow \theta_s)$ (adjoint: coshuffling)

Fact. Shuffling is right exact.

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — shuffling functors

$s \in S_n$ — simple reflection

θ_s — wall-crossing functor

Fact There are adjunctions $\theta_s \rightarrow \text{Id}$ and $\text{Id} \rightarrow \theta_s$

Definition.[Carlin] Shuffling functor $C_s := \text{Coker}(\text{Id} \rightarrow \theta_s)$ (adjoint: coshuffling)

Fact. Shuffling is right exact.

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — shuffling functors

$s \in S_n$ — simple reflection

θ_s — wall-crossing functor

Fact There are adjunctions $\theta_s \rightarrow \text{Id}$ and $\text{Id} \rightarrow \theta_s$

Definition. [Carlin] Shuffling functor $C_s := \text{Coker}(\text{Id} \rightarrow \theta_s)$ (adjoint: coshuffling)

Fact. Shuffling is right exact.

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — shuffling functors

$s \in S_n$ — simple reflection

θ_s — wall-crossing functor

Fact There are adjunctions $\theta_s \rightarrow \text{Id}$ and $\text{Id} \rightarrow \theta_s$

Definition. [Carlin] Shuffling functor $C_s := \text{Coker}(\text{Id} \rightarrow \theta_s)$ (adjoint: coshuffling)

Fact. Shuffling is right exact.

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — shuffling functors

$s \in S_n$ — simple reflection

θ_s — wall-crossing functor

Fact There are adjunctions $\theta_s \rightarrow \text{Id}$ and $\text{Id} \rightarrow \theta_s$

Definition. [Carlin] Shuffling functor $C_s := \text{Coker}(\text{Id} \rightarrow \theta_s)$ (adjoint: coshuffling)

Fact. Shuffling is right exact.

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — shuffling functors

$s \in S_n$ — simple reflection

θ_s — wall-crossing functor

Fact There are adjunctions $\theta_s \rightarrow \text{Id}$ and $\text{Id} \rightarrow \theta_s$

Definition. [Carlin] Shuffling functor $C_s := \text{Coker}(\text{Id} \rightarrow \theta_s)$ (adjoint: coshuffling)

Fact. Shuffling is right exact.

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — assignments

Assign:

Cap diagram: Translation onto a wall.

Cup diagram: Translation out of a wall.

Right crossing: Derived shuffling functor.

Right crossing: Derived coshuffling functor.

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — assignments

Assign:

Cap diagram: Translation onto a wall.

Cup diagram: Translation out of a wall.

Right crossing: Derived shuffling functor.

Right crossing: Derived coshuffling functor.

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — assignments

Assign:

Cap diagram: Translation onto a wall.

Cup diagram: Translation out of a wall.

Right crossing: Derived shuffling functor.

Right crossing: Derived coshuffling functor.

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — assignments

Assign:

Cap diagram: Translation onto a wall.

Cup diagram: Translation out of a wall.

Right crossing: Derived shuffling functor.

Right crossing: Derived coshuffling functor.

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — assignments

Assign:

Cap diagram: Translation onto a wall.

Cup diagram: Translation out of a wall.

Right crossing: Derived shuffling functor.

Right crossing: Derived coshuffling functor.

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — assignments

Assign:

Cap diagram: Translation onto a wall.

Cup diagram: Translation out of a wall.

Right crossing: Derived shuffling functor.

Right crossing: Derived coshuffling functor.

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants — assignments

Assign:

Cap diagram: Translation onto a wall.

Cup diagram: Translation out of a wall.

Right crossing: Derived shuffling functor.

Right crossing: Derived coshuffling functor.

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants

Theorem.[Stroppel] This gives rise to a functor $\mathcal{F} : \mathbf{Tang} \rightarrow \mathbf{Cat}$ (up to position and degree shifts).

Theorem.[Stroppel] For L oriented link, the functor $\mathcal{F}(T_L)[n_-]\langle n_+ - 2n_- \rangle$ is an invariant of L .

Theorem.[Brundan-Stroppel] This is equivalent to Khovanov's categorification of Jones polynomial.

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants

Theorem. [Stroppel] This gives rise to a functor $\mathcal{F} : \mathbf{Tang} \rightarrow \mathbf{Cat}$ (up to position and degree shifts).

Theorem. [Stroppel] For L oriented link, the functor $\mathcal{F}(T_L)[n_-]\langle n_+ - 2n_- \rangle$ is an invariant of L .

Theorem. [Brundan-Stroppel] This is equivalent to Khovanov's categorification of Jones polynomial.

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants

Theorem. [Stroppel] This gives rise to a functor $\mathcal{F} : \mathbf{Tang} \rightarrow \mathbf{Cat}$ (up to position and degree shifts).

Theorem. [Stroppel] For L oriented link, the functor $\mathcal{F}(T_L)[n_-]\langle n_+ - 2n_- \rangle$ is an invariant of L .

Theorem. [Brundan-Stroppel] This is equivalent to Khovanov's categorification of Jones polynomial.

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants

Theorem. [Stroppel] This gives rise to a functor $\mathcal{F} : \mathbf{Tang} \rightarrow \mathbf{Cat}$ (up to position and degree shifts).

Theorem. [Stroppel] For L oriented link, the functor $\mathcal{F}(T_L)[n_-]\langle n_+ - 2n_- \rangle$ is an invariant of L .

Theorem. [Brundan-Stroppel] This is equivalent to Khovanov's categorification of Jones polynomial.

Categorification of quantum $U_v(\mathfrak{sl}_2)$ -invariants

Theorem. [Stroppel] This gives rise to a functor $\mathcal{F} : \mathbf{Tang} \rightarrow \mathbf{Cat}$ (up to position and degree shifts).

Theorem. [Stroppel] For L oriented link, the functor $\mathcal{F}(T_L)[n_-]\langle n_+ - 2n_- \rangle$ is an invariant of L .

Theorem. [Brundan-Stroppel] This is equivalent to Khovanov's categorification of Jones polynomial.

THANK YOU!!!