

# Algebraic categorification and its applications, III

Volodymyr Mazorchuk  
(Uppsala University)

Winter School “Geometry and physics”  
January 17-24, 2015, Srní, Czech Republic

# 2-categories

**Definition.** A **2-category** is a category enriched over the monoidal category **Cat** of small categories (in the latter the monoidal structure is induced by the cartesian product).

This means that a 2-category  $\mathcal{C}$  is given by the following data:

- ▶ objects of  $\mathcal{C}$ ;
- ▶ small categories  $\mathcal{C}(i, j)$  of morphisms;
- ▶ bifunctorial composition  $\mathcal{C}(j, k) \times \mathcal{C}(i, j) \rightarrow \mathcal{C}(i, k)$ ;
- ▶ identity objects  $\mathbb{1}_j$ ;

which are subject to the obvious set of (strict) axioms.

# 2-categories

**Definition.** A **2-category** is a category enriched over the monoidal category **Cat** of **small** categories (in the latter the monoidal structure is induced by the cartesian product).

This means that a 2-category  $\mathcal{C}$  is given by the following data:

- ▶ objects of  $\mathcal{C}$ ;
- ▶ small categories  $\mathcal{C}(i, j)$  of morphisms;
- ▶ bifunctorial composition  $\mathcal{C}(j, k) \times \mathcal{C}(i, j) \rightarrow \mathcal{C}(i, k)$ ;
- ▶ identity objects  $1_j$ ;

which are subject to the obvious set of (strict) axioms.

# 2-categories

**Definition.** A **2-category** is a category enriched over the monoidal category **Cat** of **small** categories (in the latter the monoidal structure is induced by the cartesian product).

This means that a **2-category**  $\mathcal{C}$  is given by the following data:

- ▶ objects of  $\mathcal{C}$ ;
- ▶ small categories  $\mathcal{C}(i, j)$  of morphisms;
- ▶ bifunctorial composition  $\mathcal{C}(j, k) \times \mathcal{C}(i, j) \rightarrow \mathcal{C}(i, k)$ ;
- ▶ identity objects  $1_j$ ;

which are subject to the obvious set of (strict) axioms.

# 2-categories

**Definition.** A **2-category** is a category enriched over the monoidal category **Cat** of **small** categories (in the latter the monoidal structure is induced by the cartesian product).

This means that a **2-category**  $\mathcal{C}$  is given by the following data:

- ▶ **objects** of  $\mathcal{C}$ ;
- ▶ **small categories**  $\mathcal{C}(i, j)$  of morphisms;
- ▶ **bifunctorial composition**  $\mathcal{C}(j, k) \times \mathcal{C}(i, j) \rightarrow \mathcal{C}(i, k)$ ;
- ▶ **identity objects**  $1_j$ ;

which are subject to the obvious set of (strict) axioms.

# 2-categories

**Definition.** A **2-category** is a category enriched over the monoidal category **Cat** of **small** categories (in the latter the monoidal structure is induced by the cartesian product).

This means that a **2-category**  $\mathcal{C}$  is given by the following data:

- ▶ **objects** of  $\mathcal{C}$ ;
- ▶ **small categories**  $\mathcal{C}(i, j)$  of morphisms;
- ▶ **bifunctorial composition**  $\mathcal{C}(j, k) \times \mathcal{C}(i, j) \rightarrow \mathcal{C}(i, k)$ ;
- ▶ **identity objects**  $\mathbb{1}_j$ ;

which are subject to the obvious set of (strict) axioms.

# 2-categories

**Definition.** A **2-category** is a category enriched over the monoidal category **Cat** of **small** categories (in the latter the monoidal structure is induced by the cartesian product).

This means that a **2-category**  $\mathcal{C}$  is given by the following data:

- ▶ **objects** of  $\mathcal{C}$ ;
- ▶ small **categories**  $\mathcal{C}(i, j)$  of morphisms;
- ▶ **bifunctorial** composition  $\mathcal{C}(j, k) \times \mathcal{C}(i, j) \rightarrow \mathcal{C}(i, k)$ ;
- ▶ **identity objects**  $\mathbb{1}_j$ ;

which are subject to the obvious set of (strict) axioms.

# 2-categories

**Definition.** A **2-category** is a category enriched over the monoidal category **Cat** of **small** categories (in the latter the monoidal structure is induced by the cartesian product).

This means that a **2-category**  $\mathcal{C}$  is given by the following data:

- ▶ **objects** of  $\mathcal{C}$ ;
- ▶ small **categories**  $\mathcal{C}(i, j)$  of morphisms;
- ▶ **bifunctorial** composition  $\mathcal{C}(j, k) \times \mathcal{C}(i, j) \rightarrow \mathcal{C}(i, k)$ ;
- ▶ **identity objects**  $\mathbb{1}_j$ ;

which are subject to the obvious set of (strict) axioms.

# 2-categories

**Definition.** A **2-category** is a category enriched over the monoidal category **Cat** of **small** categories (in the latter the monoidal structure is induced by the cartesian product).

This means that a **2-category**  $\mathcal{C}$  is given by the following data:

- ▶ **objects** of  $\mathcal{C}$ ;
- ▶ small **categories**  $\mathcal{C}(i, j)$  of morphisms;
- ▶ **bifunctorial** composition  $\mathcal{C}(j, k) \times \mathcal{C}(i, j) \rightarrow \mathcal{C}(i, k)$ ;
- ▶ **identity objects**  $\mathbb{1}_j$ ;

which are subject to the obvious set of (**strict**) axioms.

# 2-categories

**Definition.** A **2-category** is a category enriched over the monoidal category **Cat** of **small** categories (in the latter the monoidal structure is induced by the cartesian product).

This means that a **2-category**  $\mathcal{C}$  is given by the following data:

- ▶ **objects** of  $\mathcal{C}$ ;
- ▶ small **categories**  $\mathcal{C}(i, j)$  of morphisms;
- ▶ **bifunctorial** composition  $\mathcal{C}(j, k) \times \mathcal{C}(i, j) \rightarrow \mathcal{C}(i, k)$ ;
- ▶ **identity objects**  $\mathbb{1}_j$ ;

which are subject to the obvious set of (**strict**) axioms.

# Terminology and the principal example

## Terminology.

- ▶ An object in  $\mathcal{C}(i, j)$  is called a **1-morphism** of  $\mathcal{C}$ .
- ▶ A morphism in  $\mathcal{C}(i, j)$  is called a **2-morphism** of  $\mathcal{C}$ .
- ▶ Composition in  $\mathcal{C}(i, j)$  is called **vertical** and denoted  $\circ_1$ .
- ▶ Composition in  $\mathcal{C}$  is called **horizontal** and denoted  $\circ_0$ .

## Principal example.

The category **Cat** is a 2-category.

- ▶ Objects of **Cat** are small categories.
- ▶ 1-morphisms in **Cat** are functors.
- ▶ 2-morphisms in **Cat** are natural transformations.
- ▶ Composition is the usual composition.
- ▶ Identity 1-morphisms are the identity functors.

# Terminology and the principal example

## Terminology.

- ▶ An object in  $\mathcal{C}(i, j)$  is called a **1-morphism** of  $\mathcal{C}$ .
- ▶ A morphism in  $\mathcal{C}(i, j)$  is called a **2-morphism** of  $\mathcal{C}$ .
- ▶ Composition in  $\mathcal{C}(i, j)$  is called **vertical** and denoted  $\circ_1$ .
- ▶ Composition in  $\mathcal{C}$  is called **horizontal** and denoted  $\circ_0$ .

**Principal example.** The category **Cat** is a 2-category.

- ▶ Objects of **Cat** are small categories.
- ▶ 1-morphisms in **Cat** are functors.
- ▶ 2-morphisms in **Cat** are natural transformations.
- ▶ Composition is the usual composition.
- ▶ Identity 1-morphisms are the identity functors.

# Terminology and the principal example

## Terminology.

- ▶ An object in  $\mathcal{C}(i, j)$  is called a **1-morphism** of  $\mathcal{C}$ .
- ▶ A morphism in  $\mathcal{C}(i, j)$  is called a **2-morphism** of  $\mathcal{C}$ .
- ▶ Composition in  $\mathcal{C}(i, j)$  is called **vertical** and denoted  $\circ_1$ .
- ▶ Composition in  $\mathcal{C}$  is called **horizontal** and denoted  $\circ_0$ .

**Principal example.** The category  $\mathbf{Cat}$  is a 2-category.

- ▶ Objects of  $\mathbf{Cat}$  are small categories.
- ▶ 1-morphisms in  $\mathbf{Cat}$  are functors.
- ▶ 2-morphisms in  $\mathbf{Cat}$  are natural transformations.
- ▶ Composition is the usual composition.
- ▶ Identity 1-morphisms are the identity functors.

# Terminology and the principal example

## Terminology.

- ▶ An object in  $\mathcal{C}(i, j)$  is called a **1-morphism** of  $\mathcal{C}$ .
- ▶ A morphism in  $\mathcal{C}(i, j)$  is called a **2-morphism** of  $\mathcal{C}$ .
- ▶ Composition in  $\mathcal{C}(i, j)$  is called **vertical** and denoted  $\circ_1$ .
- ▶ Composition in  $\mathcal{C}$  is called **horizontal** and denoted  $\circ_0$ .

**Principal example.** The category  $\mathbf{Cat}$  is a 2-category.

- ▶ Objects of  $\mathbf{Cat}$  are small categories.
- ▶ 1-morphisms in  $\mathbf{Cat}$  are functors.
- ▶ 2-morphisms in  $\mathbf{Cat}$  are natural transformations.
- ▶ Composition is the usual composition.
- ▶ Identity 1-morphisms are the identity functors.

# Terminology and the principal example

## Terminology.

- ▶ An object in  $\mathcal{C}(i, j)$  is called a **1-morphism** of  $\mathcal{C}$ .
- ▶ A morphism in  $\mathcal{C}(i, j)$  is called a **2-morphism** of  $\mathcal{C}$ .
- ▶ Composition in  $\mathcal{C}(i, j)$  is called **vertical** and denoted  $\circ_1$ .
- ▶ Composition in  $\mathcal{C}$  is called **horizontal** and denoted  $\circ_0$ .

**Principal example.** The category **Cat** is a 2-category.

- ▶ Objects of **Cat** are small categories.
- ▶ 1-morphisms in **Cat** are functors.
- ▶ 2-morphisms in **Cat** are natural transformations.
- ▶ Composition is the usual composition.
- ▶ Identity 1-morphisms are the identity functors.

# Terminology and the principal example

## Terminology.

- ▶ An object in  $\mathcal{C}(i, j)$  is called a **1-morphism** of  $\mathcal{C}$ .
- ▶ A morphism in  $\mathcal{C}(i, j)$  is called a **2-morphism** of  $\mathcal{C}$ .
- ▶ Composition in  $\mathcal{C}(i, j)$  is called **vertical** and denoted  $\circ_1$ .
- ▶ Composition in  $\mathcal{C}$  is called **horizontal** and denoted  $\circ_0$ .

**Principal example.** The category **Cat** is a 2-category.

- ▶ Objects of **Cat** are **small categories**.
- ▶ 1-morphisms in **Cat** are **functors**.
- ▶ 2-morphisms in **Cat** are **natural transformations**.
- ▶ Composition is the usual composition.
- ▶ Identity 1-morphisms are the **identity functors**.

# Terminology and the principal example

## Terminology.

- ▶ An object in  $\mathcal{C}(i, j)$  is called a **1-morphism** of  $\mathcal{C}$ .
- ▶ A morphism in  $\mathcal{C}(i, j)$  is called a **2-morphism** of  $\mathcal{C}$ .
- ▶ Composition in  $\mathcal{C}(i, j)$  is called **vertical** and denoted  $\circ_1$ .
- ▶ Composition in  $\mathcal{C}$  is called **horizontal** and denoted  $\circ_0$ .

**Principal example.** The category **Cat** is a 2-category.

- ▶ Objects of **Cat** are *small categories*.
- ▶ 1-morphisms in **Cat** are *functors*.
- ▶ 2-morphisms in **Cat** are *natural transformations*.
- ▶ Composition is the *usual composition*.
- ▶ Identity 1-morphisms are the *identity functors*.

# Terminology and the principal example

## Terminology.

- ▶ An object in  $\mathcal{C}(i, j)$  is called a **1-morphism** of  $\mathcal{C}$ .
- ▶ A morphism in  $\mathcal{C}(i, j)$  is called a **2-morphism** of  $\mathcal{C}$ .
- ▶ Composition in  $\mathcal{C}(i, j)$  is called **vertical** and denoted  $\circ_1$ .
- ▶ Composition in  $\mathcal{C}$  is called **horizontal** and denoted  $\circ_0$ .

**Principal example.** The category **Cat** is a 2-category.

- ▶ Objects of **Cat** are **small categories**.
- ▶ 1-morphisms in **Cat** are **functors**.
- ▶ 2-morphisms in **Cat** are **natural transformations**.
- ▶ Composition is the usual composition.
- ▶ Identity 1-morphisms are the identity functors.

# Terminology and the principal example

## Terminology.

- ▶ An object in  $\mathcal{C}(i, j)$  is called a **1-morphism** of  $\mathcal{C}$ .
- ▶ A morphism in  $\mathcal{C}(i, j)$  is called a **2-morphism** of  $\mathcal{C}$ .
- ▶ Composition in  $\mathcal{C}(i, j)$  is called **vertical** and denoted  $\circ_1$ .
- ▶ Composition in  $\mathcal{C}$  is called **horizontal** and denoted  $\circ_0$ .

**Principal example.** The category **Cat** is a 2-category.

- ▶ Objects of **Cat** are **small categories**.
- ▶ 1-morphisms in **Cat** are **functors**.
- ▶ 2-morphisms in **Cat** are **natural transformations**.
- ▶ Composition is the usual **composition**.
- ▶ Identity 1-morphisms are the **identity functors**.

# Terminology and the principal example

## Terminology.

- ▶ An object in  $\mathcal{C}(i, j)$  is called a **1-morphism** of  $\mathcal{C}$ .
- ▶ A morphism in  $\mathcal{C}(i, j)$  is called a **2-morphism** of  $\mathcal{C}$ .
- ▶ Composition in  $\mathcal{C}(i, j)$  is called **vertical** and denoted  $\circ_1$ .
- ▶ Composition in  $\mathcal{C}$  is called **horizontal** and denoted  $\circ_0$ .

**Principal example.** The category **Cat** is a 2-category.

- ▶ Objects of **Cat** are **small categories**.
- ▶ 1-morphisms in **Cat** are **functors**.
- ▶ 2-morphisms in **Cat** are **natural transformations**.
- ▶ Composition is the usual composition.
- ▶ Identity 1-morphisms are the identity functors.

# Terminology and the principal example

## Terminology.

- ▶ An object in  $\mathcal{C}(i, j)$  is called a **1-morphism** of  $\mathcal{C}$ .
- ▶ A morphism in  $\mathcal{C}(i, j)$  is called a **2-morphism** of  $\mathcal{C}$ .
- ▶ Composition in  $\mathcal{C}(i, j)$  is called **vertical** and denoted  $\circ_1$ .
- ▶ Composition in  $\mathcal{C}$  is called **horizontal** and denoted  $\circ_0$ .

**Principal example.** The category **Cat** is a 2-category.

- ▶ Objects of **Cat** are **small categories**.
- ▶ 1-morphisms in **Cat** are **functors**.
- ▶ 2-morphisms in **Cat** are **natural transformations**.
- ▶ Composition is the usual **composition**.
- ▶ Identity 1-morphisms are the **identity functors**.

# Terminology and the principal example

## Terminology.

- ▶ An object in  $\mathcal{C}(i, j)$  is called a **1-morphism** of  $\mathcal{C}$ .
- ▶ A morphism in  $\mathcal{C}(i, j)$  is called a **2-morphism** of  $\mathcal{C}$ .
- ▶ Composition in  $\mathcal{C}(i, j)$  is called **vertical** and denoted  $\circ_1$ .
- ▶ Composition in  $\mathcal{C}$  is called **horizontal** and denoted  $\circ_0$ .

**Principal example.** The category **Cat** is a 2-category.

- ▶ Objects of **Cat** are **small categories**.
- ▶ 1-morphisms in **Cat** are **functors**.
- ▶ 2-morphisms in **Cat** are **natural transformations**.
- ▶ Composition is the usual **composition**.
- ▶ Identity 1-morphisms are the **identity functors**.

# Terminology and the principal example

## Terminology.

- ▶ An object in  $\mathcal{C}(i, j)$  is called a **1-morphism** of  $\mathcal{C}$ .
- ▶ A morphism in  $\mathcal{C}(i, j)$  is called a **2-morphism** of  $\mathcal{C}$ .
- ▶ Composition in  $\mathcal{C}(i, j)$  is called **vertical** and denoted  $\circ_1$ .
- ▶ Composition in  $\mathcal{C}$  is called **horizontal** and denoted  $\circ_0$ .

**Principal example.** The category **Cat** is a 2-category.

- ▶ Objects of **Cat** are **small categories**.
- ▶ 1-morphisms in **Cat** are **functors**.
- ▶ 2-morphisms in **Cat** are **natural transformations**.
- ▶ Composition is the usual **composition**.
- ▶ Identity 1-morphisms are the **identity functors**.

# Decategorification of additive 2-categories

**Definition.** A 2-category  $\mathcal{C}$  is **additive** if:

- ▶ Each  $\mathcal{C}(i, j)$  is additive and idempotent split.
- ▶ Horizontal composition is biadditive.

**Definition.** The **split Gorthendieck group**  $[\mathcal{A}]_{\oplus}$  of an additive category  $\mathcal{A}$  is the quotient of the free abelian group generated by  $[X]$ , where  $X$  is an object of  $\mathcal{A}$ , modulo relations  $[X] = [Y] + [Z]$  whenever  $X \cong Y \oplus Z$ .

**Definition.** The **decategorification**  $[\mathcal{C}]$  of an additive 2-category  $\mathcal{C}$  is the (usual 1-) category defined as follows:

- ▶  $[\mathcal{C}]$  has the same objects as  $\mathcal{C}$ ;
- ▶  $[\mathcal{C}](i, j) := [\mathcal{C}(i, j)]_{\oplus}$ ;
- ▶ composition in  $[\mathcal{C}]$  is induced from that in  $\mathcal{C}$ .

# Decategorification of additive 2-categories

**Definition.** A 2-category  $\mathcal{C}$  is **additive** if:

- ▶ Each  $\mathcal{C}(i, j)$  is additive and idempotent split.
- ▶ Horizontal composition is biadditive.

**Definition.** The **split Grothendieck group**  $[\mathcal{A}]_{\oplus}$  of an additive category  $\mathcal{A}$  is the quotient of the free abelian group generated by  $[X]$ , where  $X$  is an object of  $\mathcal{A}$ , modulo relations  $[X] = [Y] + [Z]$  whenever  $X \cong Y \oplus Z$ .

**Definition.** The **decategorification**  $[\mathcal{C}]$  of an additive 2-category  $\mathcal{C}$  is the (usual 1-) category defined as follows:

- ▶  $[\mathcal{C}]$  has the same objects as  $\mathcal{C}$ ;
- ▶  $[\mathcal{C}](i, j) := [\mathcal{C}(i, j)]_{\oplus}$ ;
- ▶ composition in  $[\mathcal{C}]$  is induced from that in  $\mathcal{C}$ .

# Decategorification of additive 2-categories

**Definition.** A 2-category  $\mathcal{C}$  is **additive** if:

- ▶ Each  $\mathcal{C}(i, j)$  is **additive** and **idempotent split**.
- ▶ Horizontal composition is biadditive.

**Definition.** The **split Gorthendieck group**  $[\mathcal{A}]_{\oplus}$  of an additive category  $\mathcal{A}$  is the quotient of the free abelian group generated by  $[X]$ , where  $X$  is an object of  $\mathcal{A}$ , modulo relations  $[X] = [Y] + [Z]$  whenever  $X \cong Y \oplus Z$ .

**Definition.** The **decategorification**  $[\mathcal{C}]$  of an additive 2-category  $\mathcal{C}$  is the (usual 1-) category defined as follows:

- ▶  $[\mathcal{C}]$  has the same objects as  $\mathcal{C}$ ;
- ▶  $[\mathcal{C}](i, j) := [\mathcal{C}(i, j)]_{\oplus}$ ;
- ▶ composition in  $[\mathcal{C}]$  is induced from that in  $\mathcal{C}$ .

# Decategorification of additive 2-categories

**Definition.** A 2-category  $\mathcal{C}$  is **additive** if:

- ▶ Each  $\mathcal{C}(i, j)$  is **additive** and **idempotent split**.
- ▶ Horizontal composition is **biadditive**.

**Definition.** The **split Gorthendieck group**  $[\mathcal{A}]_{\oplus}$  of an additive category  $\mathcal{A}$  is the quotient of the free abelian group generated by  $[X]$ , where  $X$  is an object of  $\mathcal{A}$ , modulo relations  $[X] = [Y] + [Z]$  whenever  $X \cong Y \oplus Z$ .

**Definition.** The **decategorification**  $[\mathcal{C}]$  of an additive 2-category  $\mathcal{C}$  is the (usual 1-) category defined as follows:

- ▶  $[\mathcal{C}]$  has the same objects as  $\mathcal{C}$ ;
- ▶  $[\mathcal{C}](i, j) := [\mathcal{C}(i, j)]_{\oplus}$ ;
- ▶ composition in  $[\mathcal{C}]$  is induced from that in  $\mathcal{C}$ .

# Decategorification of additive 2-categories

**Definition.** A 2-category  $\mathcal{C}$  is **additive** if:

- ▶ Each  $\mathcal{C}(i, j)$  is **additive** and **idempotent split**.
- ▶ Horizontal composition is **biadditive**.

**Definition.** The **split Gorthendieck group**  $[\mathcal{A}]_{\oplus}$  of an **additive** category  $\mathcal{A}$  is the quotient of the free abelian group generated by  $[X]$ , where  $X$  is an object of  $\mathcal{A}$ , modulo relations  $[X] = [Y] + [Z]$  whenever  $X \cong Y \oplus Z$ .

**Definition.** The **decategorification**  $[\mathcal{C}]$  of an additive 2-category  $\mathcal{C}$  is the (usual 1-) category defined as follows:

- ▶  $[\mathcal{C}]$  has the same objects as  $\mathcal{C}$ ;
- ▶  $[\mathcal{C}](i, j) := [\mathcal{C}(i, j)]_{\oplus}$ ;
- ▶ composition in  $[\mathcal{C}]$  is induced from that in  $\mathcal{C}$ .

# Decategorification of additive 2-categories

**Definition.** A 2-category  $\mathcal{C}$  is **additive** if:

- ▶ Each  $\mathcal{C}(i, j)$  is **additive** and **idempotent split**.
- ▶ Horizontal composition is **biadditive**.

**Definition.** The **split Gorthendieck group**  $[\mathcal{A}]_{\oplus}$  of an **additive** category  $\mathcal{A}$  is the quotient of the free abelian group generated by  $[X]$ , where  $X$  is an object of  $\mathcal{A}$ , modulo relations  $[X] = [Y] + [Z]$  whenever  $X \cong Y \oplus Z$ .

**Definition.** The **decategorification**  $[\mathcal{C}]$  of an **additive 2-category**  $\mathcal{C}$  is the (**usual 1-**) category defined as follows:

- ▶  $[\mathcal{C}]$  has the same objects as  $\mathcal{C}$ ;
- ▶  $[\mathcal{C}](i, j) := [\mathcal{C}(i, j)]_{\oplus}$ ;
- ▶ composition in  $[\mathcal{C}]$  is induced from that in  $\mathcal{C}$ .

# Decategorification of additive 2-categories

**Definition.** A 2-category  $\mathcal{C}$  is **additive** if:

- ▶ Each  $\mathcal{C}(i, j)$  is **additive** and **idempotent split**.
- ▶ Horizontal composition is **biadditive**.

**Definition.** The **split Gorthendieck group**  $[\mathcal{A}]_{\oplus}$  of an **additive** category  $\mathcal{A}$  is the quotient of the free abelian group generated by  $[X]$ , where  $X$  is an object of  $\mathcal{A}$ , modulo relations  $[X] = [Y] + [Z]$  whenever  $X \cong Y \oplus Z$ .

**Definition.** The **decategorification**  $[\mathcal{C}]$  of an **additive 2-category**  $\mathcal{C}$  is the (**usual 1-**) category defined as follows:

- ▶  $[\mathcal{C}]$  has **the same objects** as  $\mathcal{C}$ ;
- ▶  $[\mathcal{C}](i, j) := [\mathcal{C}(i, j)]_{\oplus}$ ;
- ▶ composition in  $[\mathcal{C}]$  is induced from that in  $\mathcal{C}$ .

# Decategorification of additive 2-categories

**Definition.** A 2-category  $\mathcal{C}$  is **additive** if:

- ▶ Each  $\mathcal{C}(i, j)$  is **additive** and **idempotent split**.
- ▶ Horizontal composition is **biadditive**.

**Definition.** The **split Gorthendieck group**  $[\mathcal{A}]_{\oplus}$  of an **additive** category  $\mathcal{A}$  is the quotient of the free abelian group generated by  $[X]$ , where  $X$  is an object of  $\mathcal{A}$ , modulo relations  $[X] = [Y] + [Z]$  whenever  $X \cong Y \oplus Z$ .

**Definition.** The **decategorification**  $[\mathcal{C}]$  of an **additive 2-category**  $\mathcal{C}$  is the (**usual 1-**) category defined as follows:

- ▶  $[\mathcal{C}]$  has **the same objects** as  $\mathcal{C}$ ;
- ▶  $[\mathcal{C}](i, j) := [\mathcal{C}(i, j)]_{\oplus}$ ;
- ▶ composition in  $[\mathcal{C}]$  is induced from that in  $\mathcal{C}$ .

# Decategorification of additive 2-categories

**Definition.** A 2-category  $\mathcal{C}$  is **additive** if:

- ▶ Each  $\mathcal{C}(i, j)$  is **additive** and **idempotent split**.
- ▶ Horizontal composition is **biadditive**.

**Definition.** The **split Gorthendieck group**  $[\mathcal{A}]_{\oplus}$  of an **additive** category  $\mathcal{A}$  is the quotient of the free abelian group generated by  $[X]$ , where  $X$  is an object of  $\mathcal{A}$ , modulo relations  $[X] = [Y] + [Z]$  whenever  $X \cong Y \oplus Z$ .

**Definition.** The **decategorification**  $[\mathcal{C}]$  of an **additive 2-category**  $\mathcal{C}$  is the (**usual 1-**) category defined as follows:

- ▶  $[\mathcal{C}]$  has **the same objects** as  $\mathcal{C}$ ;
- ▶  $[\mathcal{C}](i, j) := [\mathcal{C}(i, j)]_{\oplus}$ ;
- ▶ composition in  $[\mathcal{C}]$  is **induced** from that in  $\mathcal{C}$ .

# Decategorification of additive 2-categories

**Definition.** A 2-category  $\mathcal{C}$  is **additive** if:

- ▶ Each  $\mathcal{C}(i, j)$  is **additive** and **idempotent split**.
- ▶ Horizontal composition is **biadditive**.

**Definition.** The **split Gorthendieck group**  $[\mathcal{A}]_{\oplus}$  of an **additive** category  $\mathcal{A}$  is the quotient of the free abelian group generated by  $[X]$ , where  $X$  is an object of  $\mathcal{A}$ , modulo relations  $[X] = [Y] + [Z]$  whenever  $X \cong Y \oplus Z$ .

**Definition.** The **decategorification**  $[\mathcal{C}]$  of an **additive 2-category**  $\mathcal{C}$  is the (**usual 1-**) category defined as follows:

- ▶  $[\mathcal{C}]$  has **the same objects** as  $\mathcal{C}$ ;
- ▶  $[\mathcal{C}](i, j) := [\mathcal{C}(i, j)]_{\oplus}$ ;
- ▶ composition in  $[\mathcal{C}]$  is **induced** from that in  $\mathcal{C}$ .

# Categorification

$\mathcal{C}$  — additive 2-category

$[\mathcal{C}]$  — decategorification of  $\mathcal{C}$

**Definition.**  $\mathcal{C}$  is called a **categorification** of  $[\mathcal{C}]$ .

**Put differently:** **Categorification** is just the formal “inverse” of decategorification.

**Warning:** Categorification is “multi-valued” in general.

# Categorification

$\mathcal{C}$  — additive 2-category

$[\mathcal{C}]$  — decategorification of  $\mathcal{C}$

**Definition.**  $\mathcal{C}$  is called a categorification of  $[\mathcal{C}]$ .

**Put differently:** Categorification is just the formal “inverse” of decategorification.

**Warning:** Categorification is “multi-valued” in general.

# Categorification

$\mathcal{C}$  — additive 2-category

$[\mathcal{C}]$  — decategorification of  $\mathcal{C}$

**Definition.**  $\mathcal{C}$  is called a categorification of  $[\mathcal{C}]$ .

**Put differently:** Categorification is just the formal “inverse” of decategorification.

**Warning:** Categorification is “multi-valued” in general.

# Categorification

$\mathcal{C}$  — additive 2-category

$[\mathcal{C}]$  — decategorification of  $\mathcal{C}$

**Definition.**  $\mathcal{C}$  is called a **categorification** of  $[\mathcal{C}]$ .

**Put differently:** Categorification is just the formal “inverse” of decategorification.

**Warning:** Categorification is “multi-valued” in general.

# Categorification

$\mathcal{C}$  — additive 2-category

$[\mathcal{C}]$  — decategorification of  $\mathcal{C}$

**Definition.**  $\mathcal{C}$  is called a **categorification** of  $[\mathcal{C}]$ .

**Put differently:** **Categorification** is just the formal “inverse” of decategorification.

**Warning:** Categorification is “multi-valued” in general.

# Categorification

$\mathcal{C}$  — additive 2-category

$[\mathcal{C}]$  — decategorification of  $\mathcal{C}$

**Definition.**  $\mathcal{C}$  is called a categorification of  $[\mathcal{C}]$ .

**Put differently:** Categorification is just the formal “inverse” of decategorification.

**Warning:** Categorification is “multi-valued” in general.

# Categorification

$\mathcal{C}$  — additive 2-category

$[\mathcal{C}]$  — decategorification of  $\mathcal{C}$

**Definition.**  $\mathcal{C}$  is called a categorification of  $[\mathcal{C}]$ .

**Put differently:** Categorification is just the formal “inverse” of decategorification.

**Warning:** Categorification is “multi-valued” in general.

# Example: projective functors on $\mathcal{O}_0$

$$\mathfrak{g} = \mathfrak{sl}_n$$

$\mathcal{O}_0$  — principal block of category  $\mathcal{O}$  for  $\mathfrak{g}$

$\mathcal{S}$  — the 2-category of projective functors on  $\mathcal{O}_0$ , that is:

- ▶  $\mathcal{S}$  has one object  $\clubsuit$  (identified with some small category  $\mathcal{C} \cong \mathcal{O}_0$ );
- ▶ 1-morphisms in  $\mathcal{S}$  are functors isomorphic to projective functors;
- ▶ 2-morphisms in  $\mathcal{S}$  are natural transformations of functors;
- ▶ horizontal composition in  $\mathcal{S}$  is composition of functors.

**Fact.**  $\mathcal{S}$  is an additive 2-category.

**Theorem.**  $[\mathcal{S}](\clubsuit, \clubsuit) \cong \mathbb{Z}[S_n]$

**Consequence:** In this sense  $\mathcal{S}$  is a categorification of  $\mathbb{Z}[S_n]$ .

# Example: projective functors on $\mathcal{O}_0$

$$\mathfrak{g} = \mathfrak{sl}_n$$

$\mathcal{O}_0$  — principal block of category  $\mathcal{O}$  for  $\mathfrak{g}$

$\mathcal{S}$  — the 2-category of projective functors on  $\mathcal{O}_0$ , that is:

- ▶  $\mathcal{S}$  has one object  $\clubsuit$  (identified with some small category  $\mathcal{C} \cong \mathcal{O}_0$ );
- ▶ 1-morphisms in  $\mathcal{S}$  are functors isomorphic to projective functors;
- ▶ 2-morphisms in  $\mathcal{S}$  are natural transformations of functors;
- ▶ horizontal composition in  $\mathcal{S}$  is composition of functors.

**Fact.**  $\mathcal{S}$  is an additive 2-category.

**Theorem.**  $[\mathcal{S}](\clubsuit, \clubsuit) \cong \mathbb{Z}[S_n]$

**Consequence:** In this sense  $\mathcal{S}$  is a categorification of  $\mathbb{Z}[S_n]$ .

# Example: projective functors on $\mathcal{O}_0$

$$\mathfrak{g} = \mathfrak{sl}_n$$

$\mathcal{O}_0$  — principal block of category  $\mathcal{O}$  for  $\mathfrak{g}$

$\mathcal{S}$  — the 2-category of projective functors on  $\mathcal{O}_0$ , that is:

- ▶  $\mathcal{S}$  has one object  $\clubsuit$  (identified with some small category  $\mathcal{C} \cong \mathcal{O}_0$ );
- ▶ 1-morphisms in  $\mathcal{S}$  are functors isomorphic to projective functors;
- ▶ 2-morphisms in  $\mathcal{S}$  are natural transformations of functors;
- ▶ horizontal composition in  $\mathcal{S}$  is composition of functors.

**Fact.**  $\mathcal{S}$  is an additive 2-category.

**Theorem.**  $[\mathcal{S}](\clubsuit, \clubsuit) \cong \mathbb{Z}[S_n]$

**Consequence:** In this sense  $\mathcal{S}$  is a categorification of  $\mathbb{Z}[S_n]$ .

# Example: projective functors on $\mathcal{O}_0$

$$\mathfrak{g} = \mathfrak{sl}_n$$

$\mathcal{O}_0$  — **principal block** of category  $\mathcal{O}$  for  $\mathfrak{g}$

$\mathcal{S}$  — the 2-category of **projective functors** on  $\mathcal{O}_0$ , that is:

- ▶  $\mathcal{S}$  has one object  $\clubsuit$  (identified with some **small category**  $\mathcal{C} \cong \mathcal{O}_0$ );
- ▶ 1-morphisms in  $\mathcal{S}$  are functors isomorphic to **projective functors**;
- ▶ 2-morphisms in  $\mathcal{S}$  are **natural transformations** of functors;
- ▶ horizontal composition in  $\mathcal{S}$  is **composition of functors**.

**Fact.**  $\mathcal{S}$  is an additive 2-category.

**Theorem.**  $[\mathcal{S}](\clubsuit, \clubsuit) \cong \mathbb{Z}[S_n]$

**Consequence:** In this sense  $\mathcal{S}$  is a **categorification** of  $\mathbb{Z}[S_n]$ .

# Example: projective functors on $\mathcal{O}_0$

$$\mathfrak{g} = \mathfrak{sl}_n$$

$\mathcal{O}_0$  — principal block of category  $\mathcal{O}$  for  $\mathfrak{g}$

$\mathcal{S}$  — the 2-category of projective functors on  $\mathcal{O}_0$ , that is:

- ▶  $\mathcal{S}$  has one object  $\clubsuit$  (identified with some small category  $\mathcal{C} \cong \mathcal{O}_0$ );
- ▶ 1-morphisms in  $\mathcal{S}$  are functors isomorphic to projective functors;
- ▶ 2-morphisms in  $\mathcal{S}$  are natural transformations of functors;
- ▶ horizontal composition in  $\mathcal{S}$  is composition of functors.

**Fact.**  $\mathcal{S}$  is an additive 2-category.

**Theorem.**  $[\mathcal{S}](\clubsuit, \clubsuit) \cong \mathbb{Z}[S_n]$

**Consequence:** In this sense  $\mathcal{S}$  is a categorification of  $\mathbb{Z}[S_n]$ .

# Example: projective functors on $\mathcal{O}_0$

$$\mathfrak{g} = \mathfrak{sl}_n$$

$\mathcal{O}_0$  — principal block of category  $\mathcal{O}$  for  $\mathfrak{g}$

$\mathcal{S}$  — the 2-category of projective functors on  $\mathcal{O}_0$ , that is:

- ▶  $\mathcal{S}$  has one object  $\clubsuit$  (identified with some small category  $\mathcal{C} \cong \mathcal{O}_0$ );
- ▶ 1-morphisms in  $\mathcal{S}$  are functors isomorphic to projective functors;
- ▶ 2-morphisms in  $\mathcal{S}$  are natural transformations of functors;
- ▶ horizontal composition in  $\mathcal{S}$  is composition of functors.

Fact.  $\mathcal{S}$  is an additive 2-category.

Theorem.  $[\mathcal{S}](\clubsuit, \clubsuit) \cong \mathbb{Z}[S_n]$

Consequence: In this sense  $\mathcal{S}$  is a categorification of  $\mathbb{Z}[S_n]$ .

# Example: projective functors on $\mathcal{O}_0$

$$\mathfrak{g} = \mathfrak{sl}_n$$

$\mathcal{O}_0$  — principal block of category  $\mathcal{O}$  for  $\mathfrak{g}$

$\mathcal{S}$  — the 2-category of projective functors on  $\mathcal{O}_0$ , that is:

- ▶  $\mathcal{S}$  has one object  $\clubsuit$  (identified with some small category  $\mathcal{C} \cong \mathcal{O}_0$ );
- ▶ 1-morphisms in  $\mathcal{S}$  are functors isomorphic to projective functors;
- ▶ 2-morphisms in  $\mathcal{S}$  are natural transformations of functors;
- ▶ horizontal composition in  $\mathcal{S}$  is composition of functors.

**Fact.**  $\mathcal{S}$  is an additive 2-category.

**Theorem.**  $[\mathcal{S}](\clubsuit, \clubsuit) \cong \mathbb{Z}[S_n]$

**Consequence:** In this sense  $\mathcal{S}$  is a categorification of  $\mathbb{Z}[S_n]$ .

# Example: projective functors on $\mathcal{O}_0$

$$\mathfrak{g} = \mathfrak{sl}_n$$

$\mathcal{O}_0$  — principal block of category  $\mathcal{O}$  for  $\mathfrak{g}$

$\mathcal{S}$  — the 2-category of projective functors on  $\mathcal{O}_0$ , that is:

- ▶  $\mathcal{S}$  has one object  $\clubsuit$  (identified with some small category  $\mathcal{C} \cong \mathcal{O}_0$ );
- ▶ 1-morphisms in  $\mathcal{S}$  are functors isomorphic to projective functors;
- ▶ 2-morphisms in  $\mathcal{S}$  are natural transformations of functors;
- ▶ horizontal composition in  $\mathcal{S}$  is composition of functors.

**Fact.**  $\mathcal{S}$  is an additive 2-category.

**Theorem.**  $[\mathcal{S}](\clubsuit, \clubsuit) \cong \mathbb{Z}[S_n]$

**Consequence:** In this sense  $\mathcal{S}$  is a categorification of  $\mathbb{Z}[S_n]$ .

# Example: projective functors on $\mathcal{O}_0$

$$\mathfrak{g} = \mathfrak{sl}_n$$

$\mathcal{O}_0$  — principal block of category  $\mathcal{O}$  for  $\mathfrak{g}$

$\mathcal{S}$  — the 2-category of projective functors on  $\mathcal{O}_0$ , that is:

- ▶  $\mathcal{S}$  has one object  $\clubsuit$  (identified with some small category  $\mathcal{C} \cong \mathcal{O}_0$ );
- ▶ 1-morphisms in  $\mathcal{S}$  are functors isomorphic to projective functors;
- ▶ 2-morphisms in  $\mathcal{S}$  are natural transformations of functors;
- ▶ horizontal composition in  $\mathcal{S}$  is composition of functors.

**Fact.**  $\mathcal{S}$  is an additive 2-category.

**Theorem.**  $[\mathcal{S}](\clubsuit, \clubsuit) \cong \mathbb{Z}[S_n]$

**Consequence:** In this sense  $\mathcal{S}$  is a categorification of  $\mathbb{Z}[S_n]$ .

# Example: projective functors on $\mathcal{O}_0$

$$\mathfrak{g} = \mathfrak{sl}_n$$

$\mathcal{O}_0$  — **principal block** of category  $\mathcal{O}$  for  $\mathfrak{g}$

$\mathcal{S}$  — the 2-category of **projective functors** on  $\mathcal{O}_0$ , that is:

- ▶  $\mathcal{S}$  has one object  $\clubsuit$  (identified with some **small** category  $\mathcal{C} \cong \mathcal{O}_0$ );
- ▶ 1-morphisms in  $\mathcal{S}$  are functors isomorphic to **projective functors**;
- ▶ 2-morphisms in  $\mathcal{S}$  are **natural transformations** of functors;
- ▶ horizontal composition in  $\mathcal{S}$  is **composition** of functors.

**Fact.**  $\mathcal{S}$  is an **additive** 2-category.

**Theorem.**  $[\mathcal{S}](\clubsuit, \clubsuit) \cong \mathbb{Z}[S_n]$

**Consequence:** In this sense  $\mathcal{S}$  is a **categorification** of  $\mathbb{Z}[S_n]$ .

# Example: projective functors on $\mathcal{O}_0$

$$\mathfrak{g} = \mathfrak{sl}_n$$

$\mathcal{O}_0$  — **principal block** of category  $\mathcal{O}$  for  $\mathfrak{g}$

$\mathcal{S}$  — the 2-category of **projective functors** on  $\mathcal{O}_0$ , that is:

- ▶  $\mathcal{S}$  has one object  $\clubsuit$  (identified with some **small** category  $\mathcal{C} \cong \mathcal{O}_0$ );
- ▶ 1-morphisms in  $\mathcal{S}$  are functors isomorphic to **projective functors**;
- ▶ 2-morphisms in  $\mathcal{S}$  are **natural transformations** of functors;
- ▶ horizontal composition in  $\mathcal{S}$  is **composition** of functors.

**Fact.**  $\mathcal{S}$  is an **additive** 2-category.

**Theorem.**  $[\mathcal{S}](\clubsuit, \clubsuit) \cong \mathbb{Z}[S_n]$

**Consequence:** In this sense  $\mathcal{S}$  is a **categorification** of  $\mathbb{Z}[S_n]$ .

# Example: projective functors on $\mathcal{O}_0$

$$\mathfrak{g} = \mathfrak{sl}_n$$

$\mathcal{O}_0$  — **principal block** of category  $\mathcal{O}$  for  $\mathfrak{g}$

$\mathcal{S}$  — the 2-category of **projective functors** on  $\mathcal{O}_0$ , that is:

- ▶  $\mathcal{S}$  has one object  $\clubsuit$  (identified with some **small** category  $\mathcal{C} \cong \mathcal{O}_0$ );
- ▶ 1-morphisms in  $\mathcal{S}$  are functors isomorphic to **projective functors**;
- ▶ 2-morphisms in  $\mathcal{S}$  are **natural transformations** of functors;
- ▶ horizontal composition in  $\mathcal{S}$  is **composition** of functors.

**Fact.**  $\mathcal{S}$  is an **additive** 2-category.

**Theorem.**  $[\mathcal{S}](\clubsuit, \clubsuit) \cong \mathbb{Z}[S_n]$

**Consequence:** In this sense  $\mathcal{S}$  is a **categorification** of  $\mathbb{Z}[S_n]$ .

# Finitary $\mathbb{k}$ -linear categories.

$\mathbb{k}$  — field

**Definition:** An additive  $\mathbb{k}$ -linear category  $\mathcal{A}$  is **finitary** if

- ▶  $\mathcal{A}$  is **idempotent split**;
- ▶  $\mathcal{A}$  has **finitely many** indecomposables;
- ▶ all morphism spaces in  $\mathcal{A}$  are **finite dimensional** (over  $\mathbb{k}$ ).

**Example:**  $A\text{-proj}$  for a finite dimensional  $\mathbb{k}$ -algebra  $A$ .

**Fact:** If  $\mathcal{A}$  is finitary, then  $\mathcal{A} \cong A\text{-proj}$  for some  $A$ .

# Finitary $\mathbb{k}$ -linear categories.

$\mathbb{k}$  — field

**Definition:** An additive  $\mathbb{k}$ -linear category  $\mathcal{A}$  is **finitary** if

- ▶  $\mathcal{A}$  is **idempotent split**;
- ▶  $\mathcal{A}$  has **finitely many** indecomposables;
- ▶ all morphism spaces in  $\mathcal{A}$  are **finite dimensional** (over  $\mathbb{k}$ ).

**Example:**  $A\text{-proj}$  for a finite dimensional  $\mathbb{k}$ -algebra  $A$ .

**Fact:** If  $\mathcal{A}$  is finitary, then  $\mathcal{A} \cong A\text{-proj}$  for some  $A$ .

# Finitary $\mathbb{k}$ -linear categories.

$\mathbb{k}$  — field

**Definition:** An additive  $\mathbb{k}$ -linear category  $\mathcal{A}$  is **finitary** if

- ▶  $\mathcal{A}$  is **idempotent split**;
- ▶  $\mathcal{A}$  has **finitely many** indecomposables;
- ▶ all morphism spaces in  $\mathcal{A}$  are **finite dimensional** (over  $\mathbb{k}$ ).

**Example:**  $A\text{-proj}$  for a finite dimensional  $\mathbb{k}$ -algebra  $A$ .

**Fact:** If  $\mathcal{A}$  is finitary, then  $\mathcal{A} \cong A\text{-proj}$  for some  $A$ .

# Finitary $\mathbb{k}$ -linear categories.

$\mathbb{k}$  — field

**Definition:** An additive  $\mathbb{k}$ -linear category  $\mathcal{A}$  is **finitary** if

- ▶  $\mathcal{A}$  is **idempotent split**;
- ▶  $\mathcal{A}$  has **finitely many** indecomposables;
- ▶ all morphism spaces in  $\mathcal{A}$  are **finite dimensional** (over  $\mathbb{k}$ ).

**Example:**  $A\text{-proj}$  for a finite dimensional  $\mathbb{k}$ -algebra  $A$ .

**Fact:** If  $\mathcal{A}$  is finitary, then  $\mathcal{A} \cong A\text{-proj}$  for some  $A$ .

# Finitary $\mathbb{k}$ -linear categories.

$\mathbb{k}$  — field

**Definition:** An additive  $\mathbb{k}$ -linear category  $\mathcal{A}$  is **finitary** if

- ▶  $\mathcal{A}$  is **idempotent split**;
- ▶  $\mathcal{A}$  has **finitely many** indecomposables;
- ▶ all morphism spaces in  $\mathcal{A}$  are **finite dimensional** (over  $\mathbb{k}$ ).

**Example:**  $A\text{-proj}$  for a finite dimensional  $\mathbb{k}$ -algebra  $A$ .

**Fact:** If  $\mathcal{A}$  is finitary, then  $\mathcal{A} \cong A\text{-proj}$  for some  $A$ .

# Finitary $\mathbb{k}$ -linear categories.

$\mathbb{k}$  — field

**Definition:** An additive  $\mathbb{k}$ -linear category  $\mathcal{A}$  is **finitary** if

- ▶  $\mathcal{A}$  is **idempotent split**;
- ▶  $\mathcal{A}$  has **finitely many** indecomposables;
- ▶ all morphism spaces in  $\mathcal{A}$  are **finite dimensional** (over  $\mathbb{k}$ ).

**Example:**  $A\text{-proj}$  for a finite dimensional  $\mathbb{k}$ -algebra  $A$ .

**Fact:** If  $\mathcal{A}$  is finitary, then  $\mathcal{A} \cong A\text{-proj}$  for some  $A$ .

# Finitary $\mathbb{k}$ -linear categories.

$\mathbb{k}$  — field

**Definition:** An additive  $\mathbb{k}$ -linear category  $\mathcal{A}$  is **finitary** if

- ▶  $\mathcal{A}$  is **idempotent split**;
- ▶  $\mathcal{A}$  has **finitely many** indecomposables;
- ▶ all morphism spaces in  $\mathcal{A}$  are **finite dimensional** (over  $\mathbb{k}$ ).

**Example:**  $A\text{-proj}$  for a finite dimensional  $\mathbb{k}$ -algebra  $A$ .

**Fact:** If  $\mathcal{A}$  is finitary, then  $\mathcal{A} \cong A\text{-proj}$  for some  $A$ .

# Finitary $\mathbb{k}$ -linear categories.

$\mathbb{k}$  — field

**Definition:** An additive  $\mathbb{k}$ -linear category  $\mathcal{A}$  is **finitary** if

- ▶  $\mathcal{A}$  is **idempotent split**;
- ▶  $\mathcal{A}$  has **finitely many** indecomposables;
- ▶ all morphism spaces in  $\mathcal{A}$  are **finite dimensional** (over  $\mathbb{k}$ ).

**Example:**  $A\text{-proj}$  for a finite dimensional  $\mathbb{k}$ -algebra  $A$ .

**Fact:** If  $\mathcal{A}$  is finitary, then  $\mathcal{A} \cong A\text{-proj}$  for **some**  $A$ .

# Finitary $\mathbb{k}$ -linear categories.

$\mathbb{k}$  — field

**Definition:** An additive  $\mathbb{k}$ -linear category  $\mathcal{A}$  is **finitary** if

- ▶  $\mathcal{A}$  is **idempotent split**;
- ▶  $\mathcal{A}$  has **finitely many** indecomposables;
- ▶ all morphism spaces in  $\mathcal{A}$  are **finite dimensional** (over  $\mathbb{k}$ ).

**Example:**  $A\text{-proj}$  for a finite dimensional  $\mathbb{k}$ -algebra  $A$ .

**Fact:** If  $\mathcal{A}$  is finitary, then  $\mathcal{A} \cong A\text{-proj}$  for **some**  $A$ .

# Finitary 2-categories.

**Definition:** A 2-category  $\mathcal{C}$  is **finitary** over  $\mathbb{k}$  if

- ▶  $\mathcal{C}$  has **finitely many** objects;
- ▶ each  $\mathcal{C}(i, j)$  is **finitary**  $\mathbb{k}$ -linear;
- ▶ composition is **biadditive** and  **$\mathbb{k}$ -bilinear**;
- ▶ identity 1-morphisms are **indecomposable**.

**Examples:**

- ▶ Projective functors on  $\mathcal{O}_0$ ;
- ▶ Soergel bimodules over the coinvariant algebra.

Finitary 2-categories are 2-analogues of finite dimensional algebras

# Finitary 2-categories.

**Definition:** A 2-category  $\mathcal{C}$  is **finitary** over  $\mathbb{k}$  if

- ▶  $\mathcal{C}$  has **finitely many** objects;
- ▶ each  $\mathcal{C}(i, j)$  is **finitary**  $\mathbb{k}$ -linear;
- ▶ composition is **biadditive** and  $\mathbb{k}$ -bilinear;
- ▶ identity 1-morphisms are **indecomposable**.

**Examples:**

- ▶ Projective functors on  $\mathcal{O}_0$ ;
- ▶ Soergel bimodules over the coinvariant algebra.

Finitary 2-categories are 2-analogues of finite dimensional algebras

# Finitary 2-categories.

**Definition:** A 2-category  $\mathcal{C}$  is **finitary** over  $\mathbb{k}$  if

- ▶  $\mathcal{C}$  has **finitely many** objects;
- ▶ each  $\mathcal{C}(i, j)$  is **finitary**  $\mathbb{k}$ -linear;
- ▶ composition is **biadditive** and  **$\mathbb{k}$ -bilinear**;
- ▶ identity 1-morphisms are **indecomposable**.

**Examples:**

- ▶ Projective functors on  $\mathcal{O}_0$ ;
- ▶ Soergel bimodules over the coinvariant algebra.

Finitary 2-categories are 2-analogues of finite dimensional algebras

# Finitary 2-categories.

**Definition:** A 2-category  $\mathcal{C}$  is **finitary** over  $\mathbb{k}$  if

- ▶  $\mathcal{C}$  has **finitely many** objects;
- ▶ each  $\mathcal{C}(i, j)$  is **finitary**  $\mathbb{k}$ -linear;
- ▶ composition is **biadditive** and  **$\mathbb{k}$ -bilinear**;
- ▶ identity 1-morphisms are **indecomposable**.

**Examples:**

- ▶ Projective functors on  $\mathcal{O}_0$ ;
- ▶ Soergel bimodules over the coinvariant algebra.

Finitary 2-categories are 2-analogues of finite dimensional algebras

# Finitary 2-categories.

**Definition:** A 2-category  $\mathcal{C}$  is **finitary** over  $\mathbb{k}$  if

- ▶  $\mathcal{C}$  has **finitely many** objects;
- ▶ each  $\mathcal{C}(i, j)$  is **finitary**  $\mathbb{k}$ -linear;
- ▶ composition is **biadditive** and  **$\mathbb{k}$ -bilinear**;
- ▶ identity 1-morphisms are **indecomposable**.

**Examples:**

- ▶ Projective functors on  $\mathcal{O}_0$ ;
- ▶ Soergel bimodules over the coinvariant algebra.

Finitary 2-categories are 2-analogues of finite dimensional algebras

# Finitary 2-categories.

**Definition:** A 2-category  $\mathcal{C}$  is **finitary** over  $\mathbb{k}$  if

- ▶  $\mathcal{C}$  has **finitely many** objects;
- ▶ each  $\mathcal{C}(i, j)$  is **finitary**  $\mathbb{k}$ -linear;
- ▶ composition is **biadditive** and  **$\mathbb{k}$ -bilinear**;
- ▶ identity 1-morphisms are **indecomposable**.

**Examples:**

- ▶ Projective functors on  $\mathcal{O}_0$ ;
- ▶ Soergel bimodules over the coinvariant algebra.

Finitary 2-categories are 2-analogues of finite dimensional algebras

# Finitary 2-categories.

**Definition:** A 2-category  $\mathcal{C}$  is **finitary** over  $\mathbb{k}$  if

- ▶  $\mathcal{C}$  has **finitely many** objects;
- ▶ each  $\mathcal{C}(i, j)$  is **finitary**  $\mathbb{k}$ -linear;
- ▶ composition is **biadditive** and  **$\mathbb{k}$ -bilinear**;
- ▶ identity 1-morphisms are **indecomposable**.

**Examples:**

- ▶ Projective functors on  $\mathcal{O}_0$ ;
- ▶ Soergel bimodules over the coinvariant algebra.

Finitary 2-categories are 2-analogues of finite dimensional algebras

# Finitary 2-categories.

**Definition:** A 2-category  $\mathcal{C}$  is **finitary** over  $\mathbb{k}$  if

- ▶  $\mathcal{C}$  has **finitely many** objects;
- ▶ each  $\mathcal{C}(i, j)$  is **finitary**  $\mathbb{k}$ -linear;
- ▶ composition is **biadditive** and  **$\mathbb{k}$ -bilinear**;
- ▶ identity 1-morphisms are **indecomposable**.

**Examples:**

- ▶ Projective functors on  $\mathcal{O}_0$ ;
- ▶ Soergel bimodules over the coinvariant algebra.

Finitary 2-categories are 2-analogues of finite dimensional algebras

# Finitary 2-categories.

**Definition:** A 2-category  $\mathcal{C}$  is **finitary** over  $\mathbb{k}$  if

- ▶  $\mathcal{C}$  has **finitely many** objects;
- ▶ each  $\mathcal{C}(i, j)$  is **finitary**  $\mathbb{k}$ -linear;
- ▶ composition is **biadditive** and  **$\mathbb{k}$ -bilinear**;
- ▶ identity 1-morphisms are **indecomposable**.

**Examples:**

- ▶ Projective functors on  $\mathcal{O}_0$ ;
- ▶ Soergel bimodules over the coinvariant algebra.

Finitary 2-categories are 2-analogues of finite dimensional algebras

# Finitary 2-categories.

**Definition:** A 2-category  $\mathcal{C}$  is **finitary** over  $\mathbb{k}$  if

- ▶  $\mathcal{C}$  has **finitely many** objects;
- ▶ each  $\mathcal{C}(i, j)$  is **finitary**  $\mathbb{k}$ -linear;
- ▶ composition is **biadditive** and  **$\mathbb{k}$ -bilinear**;
- ▶ identity 1-morphisms are **indecomposable**.

**Examples:**

- ▶ Projective functors on  $\mathcal{O}_0$ ;
- ▶ Soergel bimodules over the coinvariant algebra.

Finitary 2-categories are 2-analogues of finite dimensional algebras

# Finitary 2-categories.

**Definition:** A 2-category  $\mathcal{C}$  is **finitary** over  $\mathbb{k}$  if

- ▶  $\mathcal{C}$  has **finitely many** objects;
- ▶ each  $\mathcal{C}(i, j)$  is **finitary**  $\mathbb{k}$ -linear;
- ▶ composition is **biadditive** and  **$\mathbb{k}$ -bilinear**;
- ▶ identity 1-morphisms are **indecomposable**.

**Examples:**

- ▶ Projective functors on  $\mathcal{O}_0$ ;
- ▶ Soergel bimodules over the coinvariant algebra.

Finitary 2-categories are 2-analogues of finite dimensional algebras

# Example: projective functors for finite dimensional algebras.

$\mathbb{k}$  — algebraically closed field

$A$  — finite dimensional  $\mathbb{k}$ -algebra

**Definition.** A **projective** endofunctor of  $A\text{-mod}$  is tensoring with a projective  $A$ - $A$ -bimodule, up to isomorphism

$\mathcal{C}$  — a small category **equivalent** to  $A\text{-mod}$

**Definition.** The 2-category  $\mathcal{C}_A$  has:

- ▶ one object  $\clubsuit$  (identified with  $\mathcal{C}$ );
- ▶ as 1-morphisms — functors in the additive closure of the identity and projective functors;
- ▶ as 2-morphisms — natural transformations of functors;
- ▶ as horizontal composition — composition of functors.

**Fact.** The 2-category  $\mathcal{C}_A$  is finitary.

# Example: projective functors for finite dimensional algebras.

$\mathbb{k}$  — algebraically closed field

$A$  — finite dimensional  $\mathbb{k}$ -algebra

**Definition.** A projective endofunctor of  $A\text{-mod}$  is tensoring with a projective  $A$ - $A$ -bimodule, up to isomorphism

$\mathcal{C}$  — a small category equivalent to  $A\text{-mod}$

**Definition.** The 2-category  $\mathcal{C}_A$  has:

- ▶ one object  $\clubsuit$  (identified with  $\mathcal{C}$ );
- ▶ as 1-morphisms — functors in the additive closure of the identity and projective functors;
- ▶ as 2-morphisms — natural transformations of functors;
- ▶ as horizontal composition — composition of functors.

**Fact.** The 2-category  $\mathcal{C}_A$  is finitary.

# Example: projective functors for finite dimensional algebras.

$\mathbb{k}$  — algebraically closed field

$A$  — finite dimensional  $\mathbb{k}$ -algebra

**Definition.** A projective endofunctor of  $A\text{-mod}$  is tensoring with a projective  $A$ - $A$ -bimodule, up to isomorphism

$\mathcal{C}$  — a small category equivalent to  $A\text{-mod}$

**Definition.** The 2-category  $\mathcal{C}_A$  has:

- ▶ one object  $\clubsuit$  (identified with  $\mathcal{C}$ );
- ▶ as 1-morphisms — functors in the additive closure of the identity and projective functors;
- ▶ as 2-morphisms — natural transformations of functors;
- ▶ as horizontal composition — composition of functors.

**Fact.** The 2-category  $\mathcal{C}_A$  is finitary.

# Example: projective functors for finite dimensional algebras.

$\mathbb{k}$  — algebraically closed field

$A$  — finite dimensional  $\mathbb{k}$ -algebra

**Definition.** A **projective** endofunctor of  $A\text{-mod}$  is tensoring with a projective  $A$ - $A$ -bimodule, up to isomorphism

$\mathcal{C}$  — a small category equivalent to  $A\text{-mod}$

**Definition.** The 2-category  $\mathcal{C}_A$  has:

- ▶ one object  $\clubsuit$  (identified with  $\mathcal{C}$ );
- ▶ as 1-morphisms — functors in the additive closure of the identity and projective functors;
- ▶ as 2-morphisms — natural transformations of functors;
- ▶ as horizontal composition — composition of functors.

**Fact.** The 2-category  $\mathcal{C}_A$  is finitary.

# Example: projective functors for finite dimensional algebras.

$\mathbb{k}$  — algebraically closed field

$A$  — finite dimensional  $\mathbb{k}$ -algebra

**Definition.** A **projective** endofunctor of  $A\text{-mod}$  is tensoring with a projective  $A$ - $A$ -bimodule, up to isomorphism

$\mathcal{C}$  — a small category **equivalent** to  $A\text{-mod}$

**Definition.** The 2-category  $\mathcal{C}_A$  has:

- ▶ one object  $\clubsuit$  (identified with  $\mathcal{C}$ );
- ▶ as 1-morphisms — functors in the additive closure of the identity and projective functors;
- ▶ as 2-morphisms — natural transformations of functors;
- ▶ as horizontal composition — composition of functors.

**Fact.** The 2-category  $\mathcal{C}_A$  is finitary.

# Example: projective functors for finite dimensional algebras.

$\mathbb{k}$  — algebraically closed field

$A$  — finite dimensional  $\mathbb{k}$ -algebra

**Definition.** A **projective** endofunctor of  $A\text{-mod}$  is tensoring with a projective  $A$ - $A$ -bimodule, up to isomorphism

$\mathcal{C}$  — a small category **equivalent** to  $A\text{-mod}$

**Definition.** The 2-category  $\mathcal{C}_A$  has:

- ▶ one object  $\clubsuit$  (identified with  $\mathcal{C}$ );
- ▶ as 1-morphisms — functors in the additive closure of the identity and projective functors;
- ▶ as 2-morphisms — natural transformations of functors;
- ▶ as horizontal composition — composition of functors.

**Fact.** The 2-category  $\mathcal{C}_A$  is finitary.

# Example: projective functors for finite dimensional algebras.

$\mathbb{k}$  — algebraically closed field

$A$  — finite dimensional  $\mathbb{k}$ -algebra

**Definition.** A **projective** endofunctor of  $A\text{-mod}$  is tensoring with a projective  $A$ - $A$ -bimodule, up to isomorphism

$\mathcal{C}$  — a small category **equivalent** to  $A\text{-mod}$

**Definition.** The 2-category  $\mathcal{C}_A$  has:

- ▶ one object  $\clubsuit$  (identified with  $\mathcal{C}$ );
- ▶ as 1-morphisms — functors in the additive closure of the identity and projective functors;
- ▶ as 2-morphisms — natural transformations of functors;
- ▶ as horizontal composition — composition of functors.

**Fact.** The 2-category  $\mathcal{C}_A$  is finitary.

# Example: projective functors for finite dimensional algebras.

$\mathbb{k}$  — algebraically closed field

$A$  — finite dimensional  $\mathbb{k}$ -algebra

**Definition.** A **projective** endofunctor of  $A\text{-mod}$  is tensoring with a projective  $A$ - $A$ -bimodule, up to isomorphism

$\mathcal{C}$  — a small category **equivalent** to  $A\text{-mod}$

**Definition.** The 2-category  $\mathcal{C}_A$  has:

- ▶ one object  $\clubsuit$  (identified with  $\mathcal{C}$ );
- ▶ as **1-morphisms** — functors in the additive closure of the **identity** and **projective** functors;
- ▶ as 2-morphisms — natural transformations of functors;
- ▶ as horizontal composition — composition of functors.

**Fact.** The 2-category  $\mathcal{C}_A$  is finitary.

# Example: projective functors for finite dimensional algebras.

$\mathbb{k}$  — algebraically closed field

$A$  — finite dimensional  $\mathbb{k}$ -algebra

**Definition.** A **projective** endofunctor of  $A\text{-mod}$  is tensoring with a projective  $A$ - $A$ -bimodule, up to isomorphism

$\mathcal{C}$  — a small category **equivalent** to  $A\text{-mod}$

**Definition.** The 2-category  $\mathcal{C}_A$  has:

- ▶ one **object**  $\clubsuit$  (identified with  $\mathcal{C}$ );
- ▶ as **1-morphisms** — functors in the additive closure of the **identity** and **projective** functors;
- ▶ as **2-morphisms** — **natural transformations** of functors;
- ▶ as **horizontal composition** — composition of functors.

**Fact.** The 2-category  $\mathcal{C}_A$  is finitary.

# Example: projective functors for finite dimensional algebras.

$\mathbb{k}$  — algebraically closed field

$A$  — finite dimensional  $\mathbb{k}$ -algebra

**Definition.** A **projective** endofunctor of  $A\text{-mod}$  is tensoring with a projective  $A$ - $A$ -bimodule, up to isomorphism

$\mathcal{C}$  — a small category **equivalent** to  $A\text{-mod}$

**Definition.** The 2-category  $\mathcal{C}_A$  has:

- ▶ one **object**  $\clubsuit$  (identified with  $\mathcal{C}$ );
- ▶ as **1-morphisms** — functors in the additive closure of the **identity** and **projective** functors;
- ▶ as **2-morphisms** — **natural transformations** of functors;
- ▶ as **horizontal composition** — **composition** of functors.

**Fact.** The 2-category  $\mathcal{C}_A$  is finitary.

# Example: projective functors for finite dimensional algebras.

$\mathbb{k}$  — algebraically closed field

$A$  — finite dimensional  $\mathbb{k}$ -algebra

**Definition.** A **projective** endofunctor of  $A\text{-mod}$  is tensoring with a projective  $A$ - $A$ -bimodule, up to isomorphism

$\mathcal{C}$  — a small category **equivalent** to  $A\text{-mod}$

**Definition.** The 2-category  $\mathcal{C}_A$  has:

- ▶ one **object**  $\clubsuit$  (identified with  $\mathcal{C}$ );
- ▶ as **1-morphisms** — functors in the additive closure of the **identity** and **projective** functors;
- ▶ as **2-morphisms** — **natural transformations** of functors;
- ▶ as **horizontal composition** — **composition** of functors.

**Fact.** The 2-category  $\mathcal{C}_A$  is **finitary**.

# Example: projective functors for finite dimensional algebras.

$\mathbb{k}$  — algebraically closed field

$A$  — finite dimensional  $\mathbb{k}$ -algebra

**Definition.** A **projective** endofunctor of  $A\text{-mod}$  is tensoring with a projective  $A$ - $A$ -bimodule, up to isomorphism

$\mathcal{C}$  — a small category **equivalent** to  $A\text{-mod}$

**Definition.** The 2-category  $\mathcal{C}_A$  has:

- ▶ one **object**  $\clubsuit$  (identified with  $\mathcal{C}$ );
- ▶ as **1-morphisms** — functors in the additive closure of the **identity** and **projective** functors;
- ▶ as **2-morphisms** — **natural transformations** of functors;
- ▶ as **horizontal composition** — **composition** of functors.

**Fact.** The 2-category  $\mathcal{C}_A$  is **finitary**.

# Combinatorics of finitary 2-categories.

$\mathcal{C}$  — finitary 2-category

$\Sigma(\mathcal{C})$  — isoclasses of indecomposable 1-morphisms in  $\mathcal{C}$

**Fact:**  $\Sigma(\mathcal{C})$  is a **multisemigroup** under

$$F \star G = \{H : H \text{ is isomorphic to a direct summand of } FG\}$$

**Left preorder:**  $F \geq_L G$  if  $F \in \Sigma(\mathcal{C}) \star G$

**Left cells:** equivalence classes w.r.t.  $\geq_L$  (a.k.a. Green's  $\mathcal{L}$ -classes)

**Similarly:** **right** and **two-sided** preorders  $\geq_R$  and  $\geq_J$  and **right** and **two-sided** cells

**Example:** For Soergel bimodules (projective functors on  $\mathcal{O}_0$ ) these are Kazhdan-Lusztig orders and cells

# Combinatorics of finitary 2-categories.

$\mathcal{C}$  — finitary 2-category

$\Sigma(\mathcal{C})$  — isoclasses of indecomposable 1-morphisms in  $\mathcal{C}$

**Fact:**  $\Sigma(\mathcal{C})$  is a multisemigroup under

$$F \star G = \{H : H \text{ is isomorphic to a direct summand of } FG\}$$

**Left preorder:**  $F \geq_L G$  if  $F \in \Sigma(\mathcal{C}) \star G$

**Left cells:** equivalence classes w.r.t.  $\geq_L$  (a.k.a. Green's  $\mathcal{L}$ -classes)

**Similarly:** right and two-sided preorders  $\geq_R$  and  $\geq_J$  and right and two-sided cells

**Example:** For Soergel bimodules (projective functors on  $\mathcal{O}_0$ ) these are Kazhdan-Lusztig orders and cells

# Combinatorics of finitary 2-categories.

$\mathcal{C}$  — finitary 2-category

$\Sigma(\mathcal{C})$  — isoclasses of indecomposable 1-morphisms in  $\mathcal{C}$

**Fact:**  $\Sigma(\mathcal{C})$  is a multisemigroup under

$$F \star G = \{H : H \text{ is isomorphic to a direct summand of } FG\}$$

**Left preorder:**  $F \geq_L G$  if  $F \in \Sigma(\mathcal{C}) \star G$

**Left cells:** equivalence classes w.r.t.  $\geq_L$  (a.k.a. Green's  $\mathcal{L}$ -classes)

**Similarly:** right and two-sided preorders  $\geq_R$  and  $\geq_J$  and right and two-sided cells

**Example:** For Soergel bimodules (projective functors on  $\mathcal{O}_0$ ) these are Kazhdan-Lusztig orders and cells

# Combinatorics of finitary 2-categories.

$\mathcal{C}$  — finitary 2-category

$\Sigma(\mathcal{C})$  — isoclasses of indecomposable 1-morphisms in  $\mathcal{C}$

**Fact:**  $\Sigma(\mathcal{C})$  is a multisemigroup under

$$F \star G = \{H : H \text{ is isomorphic to a direct summand of } FG\}$$

Left preorder:  $F \geq_L G$  if  $F \in \Sigma(\mathcal{C}) \star G$

Left cells: equivalence classes w.r.t.  $\geq_L$  (a.k.a. Green's  $\mathcal{L}$ -classes)

Similarly: right and two-sided preorders  $\geq_R$  and  $\geq_J$  and right and two-sided cells

**Example:** For Soergel bimodules (projective functors on  $\mathcal{O}_0$ ) these are Kazhdan-Lusztig orders and cells

# Combinatorics of finitary 2-categories.

$\mathcal{C}$  — finitary 2-category

$\Sigma(\mathcal{C})$  — isoclasses of indecomposable 1-morphisms in  $\mathcal{C}$

**Fact:**  $\Sigma(\mathcal{C})$  is a multisemigroup under

$$F \star G = \{H : H \text{ is isomorphic to a direct summand of } FG\}$$

Left preorder:  $F \geq_L G$  if  $F \in \Sigma(\mathcal{C}) \star G$

Left cells: equivalence classes w.r.t.  $\geq_L$  (a.k.a. Green's  $\mathcal{L}$ -classes)

Similarly: right and two-sided preorders  $\geq_R$  and  $\geq_J$  and right and two-sided cells

**Example:** For Soergel bimodules (projective functors on  $\mathcal{O}_0$ ) these are Kazhdan-Lusztig orders and cells

# Combinatorics of finitary 2-categories.

$\mathcal{C}$  — finitary 2-category

$\Sigma(\mathcal{C})$  — isoclasses of indecomposable 1-morphisms in  $\mathcal{C}$

**Fact:**  $\Sigma(\mathcal{C})$  is a multisemigroup under

$$F \star G = \{H : H \text{ is isomorphic to a direct summand of } FG\}$$

**Left preorder:**  $F \geq_L G$  if  $F \in \Sigma(\mathcal{C}) \star G$

**Left cells:** equivalence classes w.r.t.  $\geq_L$  (a.k.a. Green's  $\mathcal{L}$ -classes)

**Similarly:** right and two-sided preorders  $\geq_R$  and  $\geq_J$  and right and two-sided cells

**Example:** For Soergel bimodules (projective functors on  $\mathcal{O}_0$ ) these are Kazhdan-Lusztig orders and cells

# Combinatorics of finitary 2-categories.

$\mathcal{C}$  — finitary 2-category

$\Sigma(\mathcal{C})$  — isoclasses of indecomposable 1-morphisms in  $\mathcal{C}$

**Fact:**  $\Sigma(\mathcal{C})$  is a multisemigroup under

$$F \star G = \{H : H \text{ is isomorphic to a direct summand of } FG\}$$

**Left preorder:**  $F \geq_L G$  if  $F \in \Sigma(\mathcal{C}) \star G$

**Left cells:** equivalence classes w.r.t.  $\geq_L$  (a.k.a. Green's  $\mathcal{L}$ -classes)

**Similarly:** right and two-sided preorders  $\geq_R$  and  $\geq_J$  and right and two-sided cells

**Example:** For Soergel bimodules (projective functors on  $\mathcal{O}_0$ ) these are Kazhdan-Lusztig orders and cells

# Combinatorics of finitary 2-categories.

$\mathcal{C}$  — finitary 2-category

$\Sigma(\mathcal{C})$  — isoclasses of indecomposable 1-morphisms in  $\mathcal{C}$

**Fact:**  $\Sigma(\mathcal{C})$  is a multisemigroup under

$$F \star G = \{H : H \text{ is isomorphic to a direct summand of } FG\}$$

**Left preorder:**  $F \geq_L G$  if  $F \in \Sigma(\mathcal{C}) \star G$

**Left cells:** equivalence classes w.r.t.  $\geq_L$  (a.k.a. Green's  $\mathcal{L}$ -classes)

**Similarly:** right and two-sided preorders  $\geq_R$  and  $\geq_J$  and right and two-sided cells

**Example:** For Soergel bimodules (projective functors on  $\mathcal{O}_0$ ) these are Kazhdan-Lusztig orders and cells

# Combinatorics of finitary 2-categories.

$\mathcal{C}$  — finitary 2-category

$\Sigma(\mathcal{C})$  — isoclasses of indecomposable 1-morphisms in  $\mathcal{C}$

**Fact:**  $\Sigma(\mathcal{C})$  is a multisemigroup under

$$F \star G = \{H : H \text{ is isomorphic to a direct summand of } FG\}$$

**Left preorder:**  $F \geq_L G$  if  $F \in \Sigma(\mathcal{C}) \star G$

**Left cells:** equivalence classes w.r.t.  $\geq_L$  (a.k.a. Green's  $\mathcal{L}$ -classes)

**Similarly:** right and two-sided preorders  $\geq_R$  and  $\geq_J$  and right and two-sided cells

**Example:** For Soergel bimodules (projective functors on  $\mathcal{O}_0$ ) these are Kazhdan-Lusztig orders and cells

# Combinatorics of finitary 2-categories.

$\mathcal{C}$  — finitary 2-category

$\Sigma(\mathcal{C})$  — isoclasses of indecomposable 1-morphisms in  $\mathcal{C}$

**Fact:**  $\Sigma(\mathcal{C})$  is a multisemigroup under

$$F \star G = \{H : H \text{ is isomorphic to a direct summand of } FG\}$$

**Left preorder:**  $F \geq_L G$  if  $F \in \Sigma(\mathcal{C}) \star G$

**Left cells:** equivalence classes w.r.t.  $\geq_L$  (a.k.a. Green's  $\mathcal{L}$ -classes)

**Similarly:** right and two-sided preorders  $\geq_R$  and  $\geq_J$  and right and two-sided cells

**Example:** For Soergel bimodules (projective functors on  $\mathcal{O}_0$ ) these are Kazhdan-Lusztig orders and cells

## More detailed example: $\mathcal{C}_A$

$A$  — basic, connected finite dimensional  $\mathbb{k}$ -algebra

$1 = e_1 + e_2 + \cdots + e_n$  — primitive decomposition of  $1 \in A$

$B_{ij} := Ae_i \otimes_{\mathbb{k}} e_j A$  for  $i, j = 1, 2, \dots, n$

**Fact:**  $\Sigma(\mathcal{C}_A) = \{A, B_{ij} : i, j = 1, 2, \dots, n\}$

For  $\mathcal{I}_1 = \{A\}$  and  $\mathcal{I}_2 = \{B_{ij}\}$  we have  $\mathcal{I}_2 \geq_J \mathcal{I}_1$

$\mathcal{L}_j := \{B_{ij} : i = 1, 2, \dots, n\}$  and  $\mathcal{R}_i := \{B_{ij} : j = 1, 2, \dots, n\}$

$\mathcal{I}_2 = \mathcal{L}_1 \cup \cdots \cup \mathcal{L}_n = \mathcal{R}_1 \cup \cdots \cup \mathcal{R}_n$

**Note:**  $\mathcal{L}_j$  and  $\mathcal{L}_{j'}$  are not  $\geq_L$ -comparable if  $j \neq j'$ , similarly for the  $\mathcal{R}_i$ 's

# More detailed example: $\mathcal{C}_A$

$A$  — basic, connected **finite dimensional**  $\mathbb{k}$ -algebra

$1 = e_1 + e_2 + \dots + e_n$  — **primitive decomposition** of  $1 \in A$

$B_{ij} := Ae_i \otimes_{\mathbb{k}} e_j A$  for  $i, j = 1, 2, \dots, n$

**Fact:**  $\Sigma(\mathcal{C}_A) = \{A, B_{ij} : i, j = 1, 2, \dots, n\}$

For  $\mathcal{I}_1 = \{A\}$  and  $\mathcal{I}_2 = \{B_{ij}\}$  we have  $\mathcal{I}_2 \geq_J \mathcal{I}_1$

$\mathcal{L}_j := \{B_{ij} : i = 1, 2, \dots, n\}$  and  $\mathcal{R}_i := \{B_{ij} : j = 1, 2, \dots, n\}$

$\mathcal{I}_2 = \mathcal{L}_1 \cup \dots \cup \mathcal{L}_n = \mathcal{R}_1 \cup \dots \cup \mathcal{R}_n$

**Note:**  $\mathcal{L}_j$  and  $\mathcal{L}_{j'}$  are not  $\geq_L$ -comparable if  $j \neq j'$ , similarly for the  $\mathcal{R}_i$ 's

# More detailed example: $\mathcal{C}_A$

$A$  — basic, connected **finite dimensional**  $\mathbb{k}$ -algebra

$1 = e_1 + e_2 + \cdots + e_n$  — **primitive** decomposition of  $1 \in A$

$B_{ij} := Ae_i \otimes_{\mathbb{k}} e_j A$  for  $i, j = 1, 2, \dots, n$

**Fact:**  $\Sigma(\mathcal{C}_A) = \{A, B_{ij} : i, j = 1, 2, \dots, n\}$

For  $\mathcal{I}_1 = \{A\}$  and  $\mathcal{I}_2 = \{B_{ij}\}$  we have  $\mathcal{I}_2 \geq_J \mathcal{I}_1$

$\mathcal{L}_j := \{B_{ij} : i = 1, 2, \dots, n\}$  and  $\mathcal{R}_i := \{B_{ij} : j = 1, 2, \dots, n\}$

$\mathcal{I}_2 = \mathcal{L}_1 \cup \cdots \cup \mathcal{L}_n = \mathcal{R}_1 \cup \cdots \cup \mathcal{R}_n$

**Note:**  $\mathcal{L}_j$  and  $\mathcal{L}_{j'}$  are not  $\geq_L$ -comparable if  $j \neq j'$ , similarly for the  $\mathcal{R}_i$ 's

## More detailed example: $\mathcal{C}_A$

$A$  — basic, connected **finite dimensional**  $\mathbb{k}$ -algebra

$1 = e_1 + e_2 + \cdots + e_n$  — **primitive** decomposition of  $1 \in A$

$B_{ij} := Ae_i \otimes_{\mathbb{k}} e_j A$  for  $i, j = 1, 2, \dots, n$

**Fact:**  $\Sigma(\mathcal{C}_A) = \{A, B_{ij} : i, j = 1, 2, \dots, n\}$

For  $\mathcal{I}_1 = \{A\}$  and  $\mathcal{I}_2 = \{B_{ij}\}$  we have  $\mathcal{I}_2 \geq_J \mathcal{I}_1$

$\mathcal{L}_j := \{B_{ij} : i = 1, 2, \dots, n\}$  and  $\mathcal{R}_i := \{B_{ij} : j = 1, 2, \dots, n\}$

$\mathcal{I}_2 = \mathcal{L}_1 \cup \cdots \cup \mathcal{L}_n = \mathcal{R}_1 \cup \cdots \cup \mathcal{R}_n$

**Note:**  $\mathcal{L}_j$  and  $\mathcal{L}_{j'}$  are not  $\geq_L$ -comparable if  $j \neq j'$ , similarly for the  $\mathcal{R}_i$ 's

## More detailed example: $\mathcal{C}_A$

$A$  — basic, connected **finite dimensional**  $\mathbb{k}$ -algebra

$1 = e_1 + e_2 + \cdots + e_n$  — **primitive** decomposition of  $1 \in A$

$B_{ij} := Ae_i \otimes_{\mathbb{k}} e_j A$  for  $i, j = 1, 2, \dots, n$

**Fact:**  $\Sigma(\mathcal{C}_A) = \{A, B_{ij} : i, j = 1, 2, \dots, n\}$

For  $\mathcal{I}_1 = \{A\}$  and  $\mathcal{I}_2 = \{B_{ij}\}$  we have  $\mathcal{I}_2 \geq_J \mathcal{I}_1$

$\mathcal{L}_j := \{B_{ij} : i = 1, 2, \dots, n\}$  and  $\mathcal{R}_i := \{B_{ij} : j = 1, 2, \dots, n\}$

$\mathcal{I}_2 = \mathcal{L}_1 \cup \cdots \cup \mathcal{L}_n = \mathcal{R}_1 \cup \cdots \cup \mathcal{R}_n$

**Note:**  $\mathcal{L}_j$  and  $\mathcal{L}_{j'}$  are not  $\geq_L$ -comparable if  $j \neq j'$ , similarly for the  $\mathcal{R}_i$ 's

## More detailed example: $\mathcal{C}_A$

$A$  — basic, connected **finite dimensional**  $\mathbb{k}$ -algebra

$1 = e_1 + e_2 + \cdots + e_n$  — **primitive** decomposition of  $1 \in A$

$B_{ij} := Ae_i \otimes_{\mathbb{k}} e_j A$  for  $i, j = 1, 2, \dots, n$

**Fact:**  $\Sigma(\mathcal{C}_A) = \{A, B_{ij} : i, j = 1, 2, \dots, n\}$

For  $\mathcal{J}_1 = \{A\}$  and  $\mathcal{J}_2 = \{B_{ij}\}$  we have  $\mathcal{J}_2 \geq_J \mathcal{J}_1$

$\mathcal{L}_j := \{B_{ij} : i = 1, 2, \dots, n\}$  and  $\mathcal{R}_i := \{B_{ij} : j = 1, 2, \dots, n\}$

$\mathcal{J}_2 = \mathcal{L}_1 \cup \cdots \cup \mathcal{L}_n = \mathcal{R}_1 \cup \cdots \cup \mathcal{R}_n$

**Note:**  $\mathcal{L}_j$  and  $\mathcal{L}_{j'}$  are not  $\geq_L$ -comparable if  $j \neq j'$ , similarly for the  $\mathcal{R}_i$ 's

## More detailed example: $\mathcal{C}_A$

$A$  — basic, connected **finite dimensional**  $\mathbb{k}$ -algebra

$1 = e_1 + e_2 + \cdots + e_n$  — **primitive** decomposition of  $1 \in A$

$B_{ij} := Ae_i \otimes_{\mathbb{k}} e_j A$  for  $i, j = 1, 2, \dots, n$

**Fact:**  $\Sigma(\mathcal{C}_A) = \{A, B_{ij} : i, j = 1, 2, \dots, n\}$

For  $\mathcal{J}_1 = \{A\}$  and  $\mathcal{J}_2 = \{B_{ij}\}$  we have  $\mathcal{J}_2 \geq_J \mathcal{J}_1$

$\mathcal{L}_j := \{B_{ij} : i = 1, 2, \dots, n\}$  and  $\mathcal{R}_i := \{B_{ij} : j = 1, 2, \dots, n\}$

$\mathcal{J}_2 = \mathcal{L}_1 \cup \cdots \cup \mathcal{L}_n = \mathcal{R}_1 \cup \cdots \cup \mathcal{R}_n$

**Note:**  $\mathcal{L}_j$  and  $\mathcal{L}_{j'}$  are not  $\geq_L$ -comparable if  $j \neq j'$ , similarly for the  $\mathcal{R}_i$ 's

## More detailed example: $\mathcal{C}_A$

$A$  — basic, connected **finite dimensional**  $\mathbb{k}$ -algebra

$1 = e_1 + e_2 + \cdots + e_n$  — **primitive** decomposition of  $1 \in A$

$B_{ij} := Ae_i \otimes_{\mathbb{k}} e_j A$  for  $i, j = 1, 2, \dots, n$

**Fact:**  $\Sigma(\mathcal{C}_A) = \{A, B_{ij} : i, j = 1, 2, \dots, n\}$

For  $\mathcal{J}_1 = \{A\}$  and  $\mathcal{J}_2 = \{B_{ij}\}$  we have  $\mathcal{J}_2 \geq_J \mathcal{J}_1$

$\mathcal{L}_j := \{B_{ij} : i = 1, 2, \dots, n\}$  and  $\mathcal{R}_i := \{B_{ij} : j = 1, 2, \dots, n\}$

$\mathcal{J}_2 = \mathcal{L}_1 \cup \cdots \cup \mathcal{L}_n = \mathcal{R}_1 \cup \cdots \cup \mathcal{R}_n$

**Note:**  $\mathcal{L}_j$  and  $\mathcal{L}_{j'}$  are not  $\geq_L$ -comparable if  $j \neq j'$ , similarly for the  $\mathcal{R}_i$ 's

## More detailed example: $\mathcal{C}_A$

$A$  — basic, connected **finite dimensional**  $\mathbb{k}$ -algebra

$1 = e_1 + e_2 + \cdots + e_n$  — **primitive** decomposition of  $1 \in A$

$B_{ij} := Ae_i \otimes_{\mathbb{k}} e_j A$  for  $i, j = 1, 2, \dots, n$

**Fact:**  $\Sigma(\mathcal{C}_A) = \{A, B_{ij} : i, j = 1, 2, \dots, n\}$

For  $\mathcal{J}_1 = \{A\}$  and  $\mathcal{J}_2 = \{B_{ij}\}$  we have  $\mathcal{J}_2 \geq_J \mathcal{J}_1$

$\mathcal{L}_j := \{B_{ij} : i = 1, 2, \dots, n\}$  and  $\mathcal{R}_i := \{B_{ij} : j = 1, 2, \dots, n\}$

$\mathcal{J}_2 = \mathcal{L}_1 \cup \cdots \cup \mathcal{L}_n = \mathcal{R}_1 \cup \cdots \cup \mathcal{R}_n$

**Note:**  $\mathcal{L}_j$  and  $\mathcal{L}_{j'}$  are not  $\geq_L$ -comparable if  $j \neq j'$ , similarly for the  $\mathcal{R}_i$ 's

## More detailed example: $\mathcal{C}_A$

$A$  — basic, connected **finite dimensional**  $\mathbb{k}$ -algebra

$1 = e_1 + e_2 + \cdots + e_n$  — **primitive** decomposition of  $1 \in A$

$B_{ij} := Ae_i \otimes_{\mathbb{k}} e_j A$  for  $i, j = 1, 2, \dots, n$

**Fact:**  $\Sigma(\mathcal{C}_A) = \{A, B_{ij} : i, j = 1, 2, \dots, n\}$


For  $\mathcal{J}_1 = \{A\}$  and  $\mathcal{J}_2 = \{B_{ij}\}$  we have  $\mathcal{J}_2 \geq_J \mathcal{J}_1$

$\mathcal{L}_j := \{B_{ij} : i = 1, 2, \dots, n\}$  and  $\mathcal{R}_i := \{B_{ij} : j = 1, 2, \dots, n\}$


$\mathcal{J}_2 = \mathcal{L}_1 \cup \cdots \cup \mathcal{L}_n = \mathcal{R}_1 \cup \cdots \cup \mathcal{R}_n$

**Note:**  $\mathcal{L}_j$  and  $\mathcal{L}_{j'}$  are not  $\geq_L$ -comparable if  $j \neq j'$ , similarly for the  $\mathcal{R}_i$ 's


# The egg-box diagram



# The egg-box diagram



# The egg-box diagram



# Fiat 2-categories

$\mathcal{C}$  — finitary 2-category

**Definition:**  $\mathcal{C}$  is fiat if

- ▶ there is a weak involution  $*$  :  $\mathcal{C} \rightarrow \mathcal{C}$ ;
- ▶ there are adjunction 2-morphisms  $\alpha : \mathbb{1}_i \rightarrow FF^*$  and  $\beta : F^*F \rightarrow \mathbb{1}_j$  such that

$$F(\beta) \circ_1 \alpha_F = \text{id}_F \quad \text{and} \quad \beta_{F^*} \circ_1 F^*(\alpha) = \text{id}_{F^*}$$

**Note:** This makes  $F$  and  $F^*$  always biadjoint

**Examples:**

- ▶ Soergel bimodules (projective functors on  $\mathcal{O}_0$ )
- ▶  $\mathcal{C}_A$  for  $A$  self-injective and weakly symmetric

# Fiat 2-categories

$\mathcal{C}$  — finitary 2-category

**Definition:**  $\mathcal{C}$  is fiat if

- ▶ there is a weak involution  $*$  :  $\mathcal{C} \rightarrow \mathcal{C}$ ;
- ▶ there are adjunction 2-morphisms  $\alpha : \mathbb{1}_i \rightarrow FF^*$  and  $\beta : F^*F \rightarrow \mathbb{1}_j$  such that

$$F(\beta) \circ_1 \alpha_F = \text{id}_F \quad \text{and} \quad \beta_{F^*} \circ_1 F^*(\alpha) = \text{id}_{F^*}$$

**Note:** This makes  $F$  and  $F^*$  always biadjoint

**Examples:**

- ▶ Soergel bimodules (projective functors on  $\mathcal{O}_0$ )
- ▶  $\mathcal{C}_A$  for  $A$  self-injective and weakly symmetric

# Fiat 2-categories

$\mathcal{C}$  — finitary 2-category

**Definition:**  $\mathcal{C}$  is fiat if

- ▶ there is a weak involution  $*$  :  $\mathcal{C} \rightarrow \mathcal{C}$ ;
- ▶ there are adjunction 2-morphisms  $\alpha : \mathbb{1}_i \rightarrow FF^*$  and  $\beta : F^*F \rightarrow \mathbb{1}_j$  such that

$$F(\beta) \circ_1 \alpha_F = \text{id}_F \quad \text{and} \quad \beta_{F^*} \circ_1 F^*(\alpha) = \text{id}_{F^*}$$

**Note:** This makes  $F$  and  $F^*$  always biadjoint

**Examples:**

- ▶ Soergel bimodules (projective functors on  $\mathcal{O}_0$ )
- ▶  $\mathcal{C}_A$  for  $A$  self-injective and weakly symmetric

# Fiat 2-categories

$\mathcal{C}$  — finitary 2-category

**Definition:**  $\mathcal{C}$  is fiat if

- ▶ there is a weak involution  $*$  :  $\mathcal{C} \rightarrow \mathcal{C}$ ;
- ▶ there are adjunction 2-morphisms  $\alpha : \mathbb{1}_i \rightarrow FF^*$  and  $\beta : F^*F \rightarrow \mathbb{1}_j$  such that

$$F(\beta) \circ_1 \alpha_F = \text{id}_F \quad \text{and} \quad \beta_{F^*} \circ_1 F^*(\alpha) = \text{id}_{F^*}$$

**Note:** This makes  $F$  and  $F^*$  always biadjoint

**Examples:**

- ▶ Soergel bimodules (projective functors on  $\mathcal{O}_0$ )
- ▶  $\mathcal{C}_A$  for  $A$  self-injective and weakly symmetric

# Fiat 2-categories

$\mathcal{C}$  — finitary 2-category

**Definition:**  $\mathcal{C}$  is fiat if

- ▶ there is a weak involution  $*$  :  $\mathcal{C} \rightarrow \mathcal{C}$ ;
- ▶ there are adjunction 2-morphisms  $\alpha : \mathbb{1}_i \rightarrow FF^*$  and  $\beta : F^*F \rightarrow \mathbb{1}_j$  such that

$$F(\beta) \circ_1 \alpha_F = \text{id}_F \quad \text{and} \quad \beta_{F^*} \circ_1 F^*(\alpha) = \text{id}_{F^*}$$

**Note:** This makes  $F$  and  $F^*$  always biadjoint

**Examples:**

- ▶ Soergel bimodules (projective functors on  $\mathcal{O}_0$ )
- ▶  $\mathcal{C}_A$  for  $A$  self-injective and weakly symmetric

# Fiat 2-categories

$\mathcal{C}$  — finitary 2-category

**Definition:**  $\mathcal{C}$  is fiat if

- ▶ there is a weak involution  $*$  :  $\mathcal{C} \rightarrow \mathcal{C}$ ;
- ▶ there are adjunction 2-morphisms  $\alpha : \mathbb{1}_i \rightarrow FF^*$  and  $\beta : F^*F \rightarrow \mathbb{1}_j$  such that

$$F(\beta) \circ_1 \alpha_F = \text{id}_F \quad \text{and} \quad \beta_{F^*} \circ_1 F^*(\alpha) = \text{id}_{F^*}$$

**Note:** This makes  $F$  and  $F^*$  always biadjoint

Examples:

- ▶ Soergel bimodules (projective functors on  $\mathcal{O}_0$ )
- ▶  $\mathcal{C}_A$  for  $A$  self-injective and weakly symmetric

# Fiat 2-categories

$\mathcal{C}$  — finitary 2-category

**Definition:**  $\mathcal{C}$  is fiat if

- ▶ there is a weak involution  $*$  :  $\mathcal{C} \rightarrow \mathcal{C}$ ;
- ▶ there are adjunction 2-morphisms  $\alpha : \mathbb{1}_i \rightarrow FF^*$  and  $\beta : F^*F \rightarrow \mathbb{1}_j$  such that

$$F(\beta) \circ_1 \alpha_F = \text{id}_F \quad \text{and} \quad \beta_{F^*} \circ_1 F^*(\alpha) = \text{id}_{F^*}$$

**Note:** This makes  $F$  and  $F^*$  always biadjoint

**Examples:**

- ▶ Soergel bimodules (projective functors on  $\mathcal{O}_0$ )
- ▶  $\mathcal{C}_A$  for  $A$  self-injective and weakly symmetric

# Fiat 2-categories

$\mathcal{C}$  — finitary 2-category

**Definition:**  $\mathcal{C}$  is fiat if

- ▶ there is a weak involution  $*$  :  $\mathcal{C} \rightarrow \mathcal{C}$ ;
- ▶ there are adjunction 2-morphisms  $\alpha : \mathbb{1}_i \rightarrow FF^*$  and  $\beta : F^*F \rightarrow \mathbb{1}_j$  such that

$$F(\beta) \circ_1 \alpha_F = \text{id}_F \quad \text{and} \quad \beta_{F^*} \circ_1 F^*(\alpha) = \text{id}_{F^*}$$

**Note:** This makes  $F$  and  $F^*$  always biadjoint

**Examples:**

- ▶ Soergel bimodules (projective functors on  $\mathcal{O}_0$ )
- ▶  $\mathcal{C}_A$  for  $A$  self-injective and weakly symmetric

# Fiat 2-categories

$\mathcal{C}$  — finitary 2-category

**Definition:**  $\mathcal{C}$  is fiat if

- ▶ there is a weak involution  $*$  :  $\mathcal{C} \rightarrow \mathcal{C}$ ;
- ▶ there are adjunction 2-morphisms  $\alpha : \mathbb{1}_i \rightarrow FF^*$  and  $\beta : F^*F \rightarrow \mathbb{1}_j$  such that

$$F(\beta) \circ_1 \alpha_F = \text{id}_F \quad \text{and} \quad \beta_{F^*} \circ_1 F^*(\alpha) = \text{id}_{F^*}$$

**Note:** This makes  $F$  and  $F^*$  always biadjoint

**Examples:**

- ▶ Soergel bimodules (projective functors on  $\mathcal{O}_0$ )
- ▶  $\mathcal{C}_A$  for  $A$  self-injective and weakly symmetric

# Fiat 2-categories

$\mathcal{C}$  — finitary 2-category

**Definition:**  $\mathcal{C}$  is fiat if

- ▶ there is a weak involution  $*$  :  $\mathcal{C} \rightarrow \mathcal{C}$ ;
- ▶ there are adjunction 2-morphisms  $\alpha : \mathbb{1}_i \rightarrow FF^*$  and  $\beta : F^*F \rightarrow \mathbb{1}_j$  such that

$$F(\beta) \circ_1 \alpha_F = \text{id}_F \quad \text{and} \quad \beta_{F^*} \circ_1 F^*(\alpha) = \text{id}_{F^*}$$

**Note:** This makes  $F$  and  $F^*$  always biadjoint

**Examples:**

- ▶ Soergel bimodules (projective functors on  $\mathcal{O}_0$ )
- ▶  $\mathcal{C}_A$  for  $A$  self-injective and weakly symmetric

# Example: 2-Kac-Moody algebras

$\mathfrak{g}$  — Kac-Moody algebra

$U_q(\mathfrak{g})$  — the corresponding quantum group

$\dot{U}$  — the idempotent completion of  $U_q(\mathfrak{g})$

$\dot{U}_{\mathbb{Z}}$  — the integral form for  $\dot{U}$

There is a number of 2-categories associated to  $\mathfrak{g}$ .

**Due to:** Khovanov-Lauda, Rouquier, Webster, Cautis-Lauda

**Some of these** categorify  $\dot{U}_{\mathbb{Z}}$ .

**Remark.** They have involution and adjunctions but are not finitary.

**Remark.** Some of them have finitary quotients.

# Example: 2-Kac-Moody algebras

$\mathfrak{g}$  — Kac-Moody algebra

$U_q(\mathfrak{g})$  — the corresponding quantum group

$\dot{U}$  — the idempotent completion of  $U_q(\mathfrak{g})$

$\dot{U}_{\mathbb{Z}}$  — the integral form for  $\dot{U}$

There is a number of 2-categories associated to  $\mathfrak{g}$ .

**Due to:** Khovanov-Lauda, Rouquier, Webster, Cautis-Lauda

**Some of these categorify**  $\dot{U}_{\mathbb{Z}}$ .

**Remark.** They have involution and adjunctions but are not finitary.

**Remark.** Some of them have finitary quotients.

# Example: 2-Kac-Moody algebras

$\mathfrak{g}$  — Kac-Moody algebra

$U_q(\mathfrak{g})$  — the corresponding quantum group

$\dot{U}$  — the idempotent completion of  $U_q(\mathfrak{g})$

$\dot{U}_{\mathbb{Z}}$  — the integral form for  $\dot{U}$

There is a number of 2-categories associated to  $\mathfrak{g}$ .

**Due to:** Khovanov-Lauda, Rouquier, Webster, Cautis-Lauda

**Some of these categorify**  $\dot{U}_{\mathbb{Z}}$ .

**Remark.** They have involution and adjunctions but are not finitary.

**Remark.** Some of them have finitary quotients.

# Example: 2-Kac-Moody algebras

$\mathfrak{g}$  — Kac-Moody algebra

$U_q(\mathfrak{g})$  — the corresponding quantum group

$\dot{U}$  — the idempotent completion of  $U_q(\mathfrak{g})$

$\dot{U}_{\mathbb{Z}}$  — the integral form for  $\dot{U}$

There is a number of 2-categories associated to  $\mathfrak{g}$ .

Due to: Khovanov-Lauda, Rouquier, Webster, Cautis-Lauda

Some of these categorify  $\dot{U}_{\mathbb{Z}}$ .

Remark. They have involution and adjunctions but are not finitary.

Remark. Some of them have finitary quotients.

# Example: 2-Kac-Moody algebras

$\mathfrak{g}$  — Kac-Moody algebra

$U_q(\mathfrak{g})$  — the corresponding quantum group

$\dot{U}$  — the idempotent completion of  $U_q(\mathfrak{g})$

$\dot{U}_{\mathbb{Z}}$  — the integral form for  $\dot{U}$

There is a number of 2-categories associated to  $\mathfrak{g}$ .

**Due to:** Khovanov-Lauda, Rouquier, Webster, Cautis-Lauda

Some of these categorify  $\dot{U}_{\mathbb{Z}}$ .

**Remark.** They have involution and adjunctions but are not finitary.

**Remark.** Some of them have finitary quotients.

# Example: 2-Kac-Moody algebras

$\mathfrak{g}$  — Kac-Moody algebra

$U_q(\mathfrak{g})$  — the corresponding quantum group

$\dot{U}$  — the idempotent completion of  $U_q(\mathfrak{g})$

$\dot{U}_{\mathbb{Z}}$  — the integral form for  $\dot{U}$

There is a number of 2-categories associated to  $\mathfrak{g}$ .

Due to: Khovanov-Lauda, Rouquier, Webster, Cautis-Lauda

Some of these categorify  $\dot{U}_{\mathbb{Z}}$ .

Remark. They have involution and adjunctions but are not finitary.

Remark. Some of them have finitary quotients.

# Example: 2-Kac-Moody algebras

$\mathfrak{g}$  — Kac-Moody algebra

$U_q(\mathfrak{g})$  — the corresponding quantum group

$\dot{U}$  — the idempotent completion of  $U_q(\mathfrak{g})$

$\dot{U}_{\mathbb{Z}}$  — the integral form for  $\dot{U}$

There is a number of 2-categories associated to  $\mathfrak{g}$ .

**Due to:** Khovanov-Lauda, Rouquier, Webster, Cautis-Lauda

Some of these categorify  $\dot{U}_{\mathbb{Z}}$ .

**Remark.** They have involution and adjunctions but are not finitary.

**Remark.** Some of them have finitary quotients.

# Example: 2-Kac-Moody algebras

$\mathfrak{g}$  — Kac-Moody algebra

$U_q(\mathfrak{g})$  — the corresponding quantum group

$\dot{U}$  — the idempotent completion of  $U_q(\mathfrak{g})$

$\dot{U}_{\mathbb{Z}}$  — the integral form for  $\dot{U}$

There is a number of 2-categories associated to  $\mathfrak{g}$ .

**Due to:** Khovanov-Lauda, Rouquier, Webster, Cautis-Lauda

**Some of these** categorify  $\dot{U}_{\mathbb{Z}}$ .

**Remark.** They have involution and adjunctions but are not finitary.

**Remark.** Some of them have finitary quotients.

# Example: 2-Kac-Moody algebras

$\mathfrak{g}$  — Kac-Moody algebra

$U_q(\mathfrak{g})$  — the corresponding quantum group

$\dot{U}$  — the idempotent completion of  $U_q(\mathfrak{g})$

$\dot{U}_{\mathbb{Z}}$  — the integral form for  $\dot{U}$

There is a number of 2-categories associated to  $\mathfrak{g}$ .

**Due to:** Khovanov-Lauda, Rouquier, Webster, Cautis-Lauda

**Some of these** categorify  $\dot{U}_{\mathbb{Z}}$ .

**Remark.** They have involutions and adjunctions but are not finitary.

**Remark.** Some of them have finitary quotients.

# Example: 2-Kac-Moody algebras

$\mathfrak{g}$  — Kac-Moody algebra

$U_q(\mathfrak{g})$  — the corresponding quantum group

$\dot{U}$  — the idempotent completion of  $U_q(\mathfrak{g})$

$\dot{U}_{\mathbb{Z}}$  — the integral form for  $\dot{U}$

There is a number of 2-categories associated to  $\mathfrak{g}$ .

**Due to:** Khovanov-Lauda, Rouquier, Webster, Cautis-Lauda

**Some of these** categorify  $\dot{U}_{\mathbb{Z}}$ .

**Remark.** They have involutions and adjunctions but are not finitary.

**Remark.** Some of them have finitary quotients.

# Example: 2-Kac-Moody algebras

$\mathfrak{g}$  — Kac-Moody algebra

$U_q(\mathfrak{g})$  — the corresponding quantum group

$\dot{U}$  — the idempotent completion of  $U_q(\mathfrak{g})$

$\dot{U}_{\mathbb{Z}}$  — the integral form for  $\dot{U}$

There is a number of 2-categories associated to  $\mathfrak{g}$ .

**Due to:** Khovanov-Lauda, Rouquier, Webster, Cautis-Lauda

**Some of these** categorify  $\dot{U}_{\mathbb{Z}}$ .

**Remark.** They have involutions and adjunctions but are not finitary.

**Remark.** Some of them have finitary quotients.

# Artin-Wedderburn Theorem

## Artin-Wedderburn Theorem.

Every simple complex finite-dimensional algebra is isomorphic to  $\text{Mat}_{n \times n}(\mathbb{C})$  for some  $n$ .

## Theorem. [M.-Miemietz]

Every "simple" fiat 2-category with a strongly regular maximal two-sided cell is "essentially"  $\mathcal{C}_A$  for  $A$  self-injective and weakly symmetric.

# Artin-Wedderburn Theorem

## Artin-Wedderburn Theorem.

Every simple complex finite-dimensional algebra is isomorphic to  $\text{Mat}_{n \times n}(\mathbb{C})$  for some  $n$ .

## Theorem. [M.-Miemietz]

Every "simple" fiat 2-category with a strongly regular maximal two-sided cell is "essentially"  $\mathcal{C}_A$  for  $A$  self-injective and weakly symmetric.

# Artin-Wedderburn Theorem

## Artin-Wedderburn Theorem.

Every **simple** complex finite-dimensional algebra is isomorphic to  $\text{Mat}_{n \times n}(\mathbb{C})$  for some  $n$ .

**Theorem.** [M.-Miemietz]

Every “simple” fiat 2-category with a strongly regular maximal two-sided cell is “essentially”  $\mathcal{C}_A$  for  $A$  self-injective and weakly symmetric.

# Artin-Wedderburn Theorem

## Artin-Wedderburn Theorem.

Every **simple** complex finite-dimensional algebra is isomorphic to  $\text{Mat}_{n \times n}(\mathbb{C})$  for some  $n$ .

## Theorem. [M.-Miemietz]

Every “simple” fiat 2-category with a strongly regular maximal two-sided cell is “essentially”  $\mathcal{C}_A$  for  $A$  self-injective and weakly symmetric.

# Artin-Wedderburn Theorem

## Artin-Wedderburn Theorem.

Every **simple** complex finite-dimensional algebra is isomorphic to  $\text{Mat}_{n \times n}(\mathbb{C})$  for some  $n$ .

## Theorem. [M.-Miemietz]

Every “**simple**” fiat 2-category with a strongly regular maximal two-sided cell is “**essentially**”  $\mathcal{C}_A$  for  $A$  self-injective and weakly symmetric.

# Artin-Wedderburn Theorem

## Artin-Wedderburn Theorem.

Every **simple** complex finite-dimensional algebra is isomorphic to  $\text{Mat}_{n \times n}(\mathbb{C})$  for some  $n$ .

## Theorem. [M.-Miemietz]

Every “**simple**” fiat 2-category with a strongly regular maximal two-sided cell is “**essentially**”  $\mathcal{C}_A$  for  $A$  self-injective and weakly symmetric.

# 2-representations

$\mathcal{C}$  — finitary 2-category

**“Definition”:** A **2-representation** of  $\mathcal{C}$  is a functorial action of  $\mathcal{C}$  on a suitable category(ies).

**Example:** **Principal 2-representation**  $\mathbf{P}_i := \mathcal{C}(i, \_)$  for  $i \in \mathcal{C}$

**Note:** 2-representations of  $\mathcal{C}$  form a 2-category where

- ▶ 1-morphisms are 2-natural transformations
- ▶ 2-morphisms are modifications

**Note:** There is a natural notion of equivalence for 2-representations

# 2-representations

$\mathcal{C}$  — finitary 2-category

**“Definition”:** A 2-representation of  $\mathcal{C}$  is a functorial action of  $\mathcal{C}$  on a suitable category(ies).

**Example:** Principal 2-representation  $\mathbf{P}_i := \mathcal{C}(i, -)$  for  $i \in \mathcal{C}$

**Note:** 2-representations of  $\mathcal{C}$  form a 2-category where

- ▶ 1-morphisms are 2-natural transformations
- ▶ 2-morphisms are modifications

**Note:** There is a natural notion of equivalence for 2-representations

# 2-representations

$\mathcal{C}$  — finitary 2-category

**“Definition”:** A **2-representation** of  $\mathcal{C}$  is a functorial action of  $\mathcal{C}$  on a suitable category(ies).

**Example:** **Principal 2-representation**  $\mathbf{P}_i := \mathcal{C}(i, -)$  for  $i \in \mathcal{C}$

**Note:** 2-representations of  $\mathcal{C}$  form a 2-category where

- ▶ 1-morphisms are 2-natural transformations
- ▶ 2-morphisms are modifications

**Note:** There is a natural notion of equivalence for 2-representations

# 2-representations

$\mathcal{C}$  — finitary 2-category

**“Definition”:** A **2-representation** of  $\mathcal{C}$  is a functorial action of  $\mathcal{C}$  on a suitable category(ies).

**Example:** **Principal 2-representation**  $\mathbf{P}_i := \mathcal{C}(i, -)$  for  $i \in \mathcal{C}$

**Note:** 2-representations of  $\mathcal{C}$  form a 2-category where

- ▶ 1-morphisms are 2-natural transformations
- ▶ 2-morphisms are modifications

**Note:** There is a natural notion of equivalence for 2-representations

# 2-representations

$\mathcal{C}$  — finitary 2-category

**“Definition”:** A **2-representation** of  $\mathcal{C}$  is a functorial action of  $\mathcal{C}$  on a suitable category(ies).

**Example:** **Principal 2-representation**  $\mathbf{P}_i := \mathcal{C}(i, -)$  for  $i \in \mathcal{C}$

**Note:** 2-representations of  $\mathcal{C}$  form a **2-category** where

- ▶ 1-morphisms are 2-natural transformations
- ▶ 2-morphisms are modifications

**Note:** There is a natural notion of equivalence for 2-representations

# 2-representations

$\mathcal{C}$  — finitary 2-category

**“Definition”:** A **2-representation** of  $\mathcal{C}$  is a functorial action of  $\mathcal{C}$  on a suitable category(ies).

**Example:** **Principal 2-representation**  $\mathbf{P}_i := \mathcal{C}(i, -)$  for  $i \in \mathcal{C}$

**Note:** 2-representations of  $\mathcal{C}$  form a **2-category** where

- ▶ 1-morphisms are **2-natural transformations**
- ▶ 2-morphisms are **modifications**

**Note:** There is a natural notion of equivalence for 2-representations

# 2-representations

$\mathcal{C}$  — finitary 2-category

**“Definition”:** A **2-representation** of  $\mathcal{C}$  is a functorial action of  $\mathcal{C}$  on a suitable category(ies).

**Example:** **Principal 2-representation**  $\mathbf{P}_i := \mathcal{C}(i, -)$  for  $i \in \mathcal{C}$

**Note:** 2-representations of  $\mathcal{C}$  form a **2-category** where

- ▶ 1-morphisms are **2-natural transformations**
- ▶ 2-morphisms are **modifications**

**Note:** There is a natural notion of equivalence for 2-representations

# 2-representations

$\mathcal{C}$  — finitary 2-category

**“Definition”:** A **2-representation** of  $\mathcal{C}$  is a functorial action of  $\mathcal{C}$  on a suitable category(ies).

**Example:** **Principal 2-representation**  $\mathbf{P}_i := \mathcal{C}(i, -)$  for  $i \in \mathcal{C}$

**Note:** 2-representations of  $\mathcal{C}$  form a **2-category** where

- ▶ 1-morphisms are **2-natural transformations**
- ▶ 2-morphisms are **modifications**

**Note:** There is a natural notion of **equivalence** for 2-representations

# 2-representations

$\mathcal{C}$  — finitary 2-category

**“Definition”:** A **2-representation** of  $\mathcal{C}$  is a functorial action of  $\mathcal{C}$  on a suitable category(ies).

**Example:** **Principal 2-representation**  $\mathbf{P}_i := \mathcal{C}(i, -)$  for  $i \in \mathcal{C}$

**Note:** 2-representations of  $\mathcal{C}$  form a **2-category** where

- ▶ 1-morphisms are **2-natural transformations**
- ▶ 2-morphisms are **modifications**

**Note:** There is a natural notion of **equivalence** for 2-representations

# Cell 2-representations

$\mathcal{C}$  — finitary 2-category

$\mathcal{L}$  — left cell in  $\mathcal{C}$

$i$  — the source for 1-morphisms in  $\mathcal{C}$

$\mathbf{P}_i$  — the  $i$ -th principal 2-representation

$\mathbf{Q}_{\mathcal{L}}$  — 2-subrepresentation of  $\mathbf{P}_i$  generated by  $F \geq_{\mathcal{L}} \mathcal{L}$

$\mathbf{I}$  — the unique maximal  $\mathcal{C}$ -invariant ideal in  $\mathbf{Q}_{\mathcal{L}}$

**Definition:**  $\mathbf{C}_{\mathcal{L}} := \mathbf{Q}_{\mathcal{L}}/\mathbf{I}$  — the cell 2-representation of  $\mathcal{C}$  for  $\mathcal{L}$

**Example:** The defining (tautological) 2-representation of  $\mathcal{C}_A$  is equivalent to  $\mathbf{C}_{\mathcal{L}_j}$  for any  $j = 1, 2, \dots, n$ .

# Cell 2-representations

$\mathcal{C}$  — finitary 2-category

$\mathcal{L}$  — left cell in  $\mathcal{C}$

$i$  — the source for 1-morphisms in  $\mathcal{C}$

$\mathbf{P}_i$  — the  $i$ -th principal 2-representation

$\mathbf{Q}_{\mathcal{L}}$  — 2-subrepresentation of  $\mathbf{P}_i$  generated by  $F \geq_{\mathcal{L}} \mathcal{L}$

$\mathbf{I}$  — the unique maximal  $\mathcal{C}$ -invariant ideal in  $\mathbf{Q}_{\mathcal{L}}$

**Definition:**  $\mathbf{C}_{\mathcal{L}} := \mathbf{Q}_{\mathcal{L}}/\mathbf{I}$  — the cell 2-representation of  $\mathcal{C}$  for  $\mathcal{L}$

**Example:** The defining (tautological) 2-representation of  $\mathcal{C}_A$  is equivalent to  $\mathbf{C}_{\mathcal{L}_j}$  for any  $j = 1, 2, \dots, n$ .

# Cell 2-representations

$\mathcal{C}$  — finitary 2-category

$\mathcal{L}$  — left cell in  $\mathcal{C}$

$i$  — the source for 1-morphisms in  $\mathcal{C}$

$P_i$  — the  $i$ -th principal 2-representation

$Q_{\mathcal{L}}$  — 2-subrepresentation of  $P_i$  generated by  $F \geq_{\mathcal{L}} \mathcal{L}$

$I$  — the unique maximal  $\mathcal{C}$ -invariant ideal in  $Q_{\mathcal{L}}$

**Definition:**  $C_{\mathcal{L}} := Q_{\mathcal{L}}/I$  — the cell 2-representation of  $\mathcal{C}$  for  $\mathcal{L}$

**Example:** The defining (tautological) 2-representation of  $\mathcal{C}_A$  is equivalent to  $C_{\mathcal{L}_j}$  for any  $j = 1, 2, \dots, n$ .

# Cell 2-representations

$\mathcal{C}$  — finitary 2-category

$\mathcal{L}$  — left cell in  $\mathcal{C}$

$i$  — the source for 1-morphisms in  $\mathcal{C}$

$\mathbf{P}_i$  — the  $i$ -th principal 2-representation

$\mathbf{Q}_{\mathcal{L}}$  — 2-subrepresentation of  $\mathbf{P}_i$  generated by  $F \geq_{\mathcal{L}} \mathcal{L}$

$\mathbf{I}$  — the unique maximal  $\mathcal{C}$ -invariant ideal in  $\mathbf{Q}_{\mathcal{L}}$

**Definition:**  $\mathbf{C}_{\mathcal{L}} := \mathbf{Q}_{\mathcal{L}}/\mathbf{I}$  — the cell 2-representation of  $\mathcal{C}$  for  $\mathcal{L}$

**Example:** The defining (tautological) 2-representation of  $\mathcal{C}_A$  is equivalent to  $\mathbf{C}_{\mathcal{L}_j}$  for any  $j = 1, 2, \dots, n$ .

# Cell 2-representations

$\mathcal{C}$  — finitary 2-category

$\mathcal{L}$  — left cell in  $\mathcal{C}$

$i$  — the source for 1-morphisms in  $\mathcal{C}$

$\mathbf{P}_i$  — the  $i$ -th principal 2-representation

$\mathbf{Q}_{\mathcal{L}}$  — 2-subrepresentation of  $\mathbf{P}_i$  generated by  $F \geq_{\mathcal{L}} \mathcal{L}$

$\mathbf{I}$  — the unique maximal  $\mathcal{C}$ -invariant ideal in  $\mathbf{Q}_{\mathcal{L}}$

**Definition:**  $\mathbf{C}_{\mathcal{L}} := \mathbf{Q}_{\mathcal{L}}/\mathbf{I}$  — the cell 2-representation of  $\mathcal{C}$  for  $\mathcal{L}$

**Example:** The defining (tautological) 2-representation of  $\mathcal{C}_A$  is equivalent to  $\mathbf{C}_{\mathcal{L}_j}$  for any  $j = 1, 2, \dots, n$ .

# Cell 2-representations

$\mathcal{C}$  — finitary 2-category

$\mathcal{L}$  — left cell in  $\mathcal{C}$

$i$  — the source for 1-morphisms in  $\mathcal{C}$

$\mathbf{P}_i$  — the  $i$ -th principal 2-representation

$\mathbf{Q}_{\mathcal{L}}$  — 2-subrepresentation of  $\mathbf{P}_i$  generated by  $F \geq_L \mathcal{L}$

$I$  — the unique maximal  $\mathcal{C}$ -invariant ideal in  $\mathbf{Q}_{\mathcal{L}}$

**Definition:**  $\mathbf{C}_{\mathcal{L}} := \mathbf{Q}_{\mathcal{L}}/I$  — the cell 2-representation of  $\mathcal{C}$  for  $\mathcal{L}$

**Example:** The defining (tautological) 2-representation of  $\mathcal{C}_A$  is equivalent to  $\mathbf{C}_{\mathcal{L}_j}$  for any  $j = 1, 2, \dots, n$ .

# Cell 2-representations

$\mathcal{C}$  — finitary 2-category

$\mathcal{L}$  — left cell in  $\mathcal{C}$

$i$  — the source for 1-morphisms in  $\mathcal{C}$

$\mathbf{P}_i$  — the  $i$ -th principal 2-representation

$\mathbf{Q}_{\mathcal{L}}$  — 2-subrepresentation of  $\mathbf{P}_i$  generated by  $F \geq_L \mathcal{L}$

$\mathbf{I}$  — the unique maximal  $\mathcal{C}$ -invariant ideal in  $\mathbf{Q}_{\mathcal{L}}$

**Definition:**  $\mathbf{C}_{\mathcal{L}} := \mathbf{Q}_{\mathcal{L}}/\mathbf{I}$  — the cell 2-representation of  $\mathcal{C}$  for  $\mathcal{L}$

**Example:** The defining (tautological) 2-representation of  $\mathcal{C}_A$  is equivalent to  $\mathbf{C}_{\mathcal{L}_j}$  for any  $j = 1, 2, \dots, n$ .

# Cell 2-representations

$\mathcal{C}$  — finitary 2-category

$\mathcal{L}$  — left cell in  $\mathcal{C}$

$i$  — the source for 1-morphisms in  $\mathcal{C}$

$\mathbf{P}_i$  — the  $i$ -th principal 2-representation

$\mathbf{Q}_{\mathcal{L}}$  — 2-subrepresentation of  $\mathbf{P}_i$  generated by  $F \geq_L \mathcal{L}$

$\mathbf{I}$  — the unique maximal  $\mathcal{C}$ -invariant ideal in  $\mathbf{Q}_{\mathcal{L}}$

**Definition:**  $\mathbf{C}_{\mathcal{L}} := \mathbf{Q}_{\mathcal{L}}/\mathbf{I}$  — the cell 2-representation of  $\mathcal{C}$  for  $\mathcal{L}$

**Example:** The defining (tautological) 2-representation of  $\mathcal{C}_A$  is equivalent to  $\mathbf{C}_{\mathcal{L}_j}$  for any  $j = 1, 2, \dots, n$ .

# Cell 2-representations

$\mathcal{C}$  — finitary 2-category

$\mathcal{L}$  — left cell in  $\mathcal{C}$

$i$  — the source for 1-morphisms in  $\mathcal{C}$

$\mathbf{P}_i$  — the  $i$ -th principal 2-representation

$\mathbf{Q}_{\mathcal{L}}$  — 2-subrepresentation of  $\mathbf{P}_i$  generated by  $F \geq_L \mathcal{L}$

$\mathbf{I}$  — the unique maximal  $\mathcal{C}$ -invariant ideal in  $\mathbf{Q}_{\mathcal{L}}$

**Definition:**  $\mathbf{C}_{\mathcal{L}} := \mathbf{Q}_{\mathcal{L}}/\mathbf{I}$  — the cell 2-representation of  $\mathcal{C}$  for  $\mathcal{L}$

**Example:** The defining (tautological) 2-representation of  $\mathcal{C}_A$  is equivalent to  $\mathbf{C}_{\mathcal{L}_j}$  for any  $j = 1, 2, \dots, n$ .

# Cell 2-representations

$\mathcal{C}$  — finitary 2-category

$\mathcal{L}$  — left cell in  $\mathcal{C}$

$i$  — the source for 1-morphisms in  $\mathcal{C}$

$\mathbf{P}_i$  — the  $i$ -th principal 2-representation

$\mathbf{Q}_{\mathcal{L}}$  — 2-subrepresentation of  $\mathbf{P}_i$  generated by  $F \geq_L \mathcal{L}$

$\mathbf{I}$  — the unique maximal  $\mathcal{C}$ -invariant ideal in  $\mathbf{Q}_{\mathcal{L}}$

**Definition:**  $\mathbf{C}_{\mathcal{L}} := \mathbf{Q}_{\mathcal{L}}/\mathbf{I}$  — the cell 2-representation of  $\mathcal{C}$  for  $\mathcal{L}$

**Example:** The defining (tautological) 2-representation of  $\mathcal{C}_A$  is equivalent to  $\mathbf{C}_{\mathcal{L}_j}$  for any  $j = 1, 2, \dots, n$ .

# Transitive 2-representations

$\mathcal{C}$  — finitary 2-category

$M$  — 2-representation of  $\mathcal{C}$

**Definition:**  $M$  is **finitary** if  $M(i)$  is finitary  $\mathbb{k}$ -linear for all  $i$

**Definition:**  $M$  is **transitive** if  $M$  is finitary and for any indecomposable  $X, Y$  in  $M$  there is a 1-morphism  $F$  such that  $X$  is isomorphic to a direct summand of  $F Y$

**Intuition:** Transitive action of a group (for us: a multisemigroup)

**Definition:**  $M$  is **simple transitive** if  $M$  is transitive and has no non-trivial  $\mathcal{C}$ -invariant ideals.

**Example:** Cell 2-representations are simple transitive.

# Transitive 2-representations

$\mathcal{C}$  — finitary 2-category

$M$  — 2-representation of  $\mathcal{C}$

**Definition:**  $M$  is finitary if  $M(i)$  is finitary  $\mathbb{k}$ -linear for all  $i$

**Definition:**  $M$  is transitive if  $M$  is finitary and for any indecomposable  $X, Y$  in  $M$  there is a 1-morphism  $F$  such that  $X$  is isomorphic to a direct summand of  $F Y$

**Intuition:** Transitive action of a group (for us: a multisemigroup)

**Definition:**  $M$  is simple transitive if  $M$  is transitive and has no non-trivial  $\mathcal{C}$ -invariant ideals.

**Example:** Cell 2-representations are simple transitive.

# Transitive 2-representations

$\mathcal{C}$  — finitary 2-category

$\mathbf{M}$  — 2-representation of  $\mathcal{C}$

**Definition:**  $\mathbf{M}$  is finitary if  $\mathbf{M}(i)$  is finitary  $\mathbb{k}$ -linear for all  $i$

**Definition:**  $\mathbf{M}$  is transitive if  $\mathbf{M}$  is finitary and for any indecomposable  $X, Y$  in  $\mathbf{M}$  there is a 1-morphism  $F$  such that  $X$  is isomorphic to a direct summand of  $F Y$

**Intuition:** Transitive action of a group (for us: a multisemigroup)

**Definition:**  $\mathbf{M}$  is simple transitive if  $\mathbf{M}$  is transitive and has no non-trivial  $\mathcal{C}$ -invariant ideals.

**Example:** Cell 2-representations are simple transitive.

# Transitive 2-representations

$\mathcal{C}$  — finitary 2-category

$\mathbf{M}$  — 2-representation of  $\mathcal{C}$

**Definition:**  $\mathbf{M}$  is finitary if  $\mathbf{M}(i)$  is finitary  $\mathbb{k}$ -linear for all  $i$

**Definition:**  $\mathbf{M}$  is transitive if  $\mathbf{M}$  is finitary and for any indecomposable  $X, Y$  in  $\mathbf{M}$  there is a 1-morphism  $F$  such that  $X$  is isomorphic to a direct summand of  $F Y$

**Intuition:** Transitive action of a group (for us: a multisemigroup)

**Definition:**  $\mathbf{M}$  is simple transitive if  $\mathbf{M}$  is transitive and has no non-trivial  $\mathcal{C}$ -invariant ideals.

**Example:** Cell 2-representations are simple transitive.

# Transitive 2-representations

$\mathcal{C}$  — finitary 2-category

$\mathbf{M}$  — 2-representation of  $\mathcal{C}$

**Definition:**  $\mathbf{M}$  is finitary if  $\mathbf{M}(i)$  is finitary  $\mathbb{k}$ -linear for all  $i$

**Definition:**  $\mathbf{M}$  is transitive if  $\mathbf{M}$  is finitary and for any indecomposable  $X, Y$  in  $\mathbf{M}$  there is a 1-morphism  $F$  such that  $X$  is isomorphic to a direct summand of  $F Y$

**Intuition:** Transitive action of a group (for us: a multisemigroup)

**Definition:**  $\mathbf{M}$  is simple transitive if  $\mathbf{M}$  is transitive and has no non-trivial  $\mathcal{C}$ -invariant ideals.

**Example:** Cell 2-representations are simple transitive.

# Transitive 2-representations

$\mathcal{C}$  — finitary 2-category

$\mathbf{M}$  — 2-representation of  $\mathcal{C}$

**Definition:**  $\mathbf{M}$  is finitary if  $\mathbf{M}(i)$  is finitary  $\mathbb{k}$ -linear for all  $i$

**Definition:**  $\mathbf{M}$  is transitive if  $\mathbf{M}$  is finitary and for any indecomposable  $X, Y$  in  $\mathbf{M}$  there is a 1-morphism  $F$  such that  $X$  is isomorphic to a direct summand of  $F Y$

**Intuition:** Transitive action of a group (for us: a multisemigroup)

**Definition:**  $\mathbf{M}$  is simple transitive if  $\mathbf{M}$  is transitive and has no non-trivial  $\mathcal{C}$ -invariant ideals.

**Example:** Cell 2-representations are simple transitive.

# Transitive 2-representations

$\mathcal{C}$  — finitary 2-category

$\mathbf{M}$  — 2-representation of  $\mathcal{C}$

**Definition:**  $\mathbf{M}$  is finitary if  $\mathbf{M}(i)$  is finitary  $\mathbb{k}$ -linear for all  $i$

**Definition:**  $\mathbf{M}$  is transitive if  $\mathbf{M}$  is finitary and for any indecomposable  $X, Y$  in  $\mathbf{M}$  there is a 1-morphism  $F$  such that  $X$  is isomorphic to a direct summand of  $F Y$

**Intuition:** Transitive action of a group (for us: a multisemigroup)

**Definition:**  $\mathbf{M}$  is simple transitive if  $\mathbf{M}$  is transitive and has no non-trivial  $\mathcal{C}$ -invariant ideals.

**Example:** Cell 2-representations are simple transitive.

# Transitive 2-representations

$\mathcal{C}$  — finitary 2-category

$\mathbf{M}$  — 2-representation of  $\mathcal{C}$

**Definition:**  $\mathbf{M}$  is finitary if  $\mathbf{M}(i)$  is finitary  $\mathbb{k}$ -linear for all  $i$

**Definition:**  $\mathbf{M}$  is transitive if  $\mathbf{M}$  is finitary and for any indecomposable  $X, Y$  in  $\mathbf{M}$  there is a 1-morphism  $F$  such that  $X$  is isomorphic to a direct summand of  $F Y$

**Intuition:** Transitive action of a group (for us: a multisemigroup)

**Definition:**  $\mathbf{M}$  is simple transitive if  $\mathbf{M}$  is transitive and has no non-trivial  $\mathcal{C}$ -invariant ideals.

**Example:** Cell 2-representations are simple transitive.

# Transitive 2-representations

$\mathcal{C}$  — finitary 2-category

$\mathbf{M}$  — 2-representation of  $\mathcal{C}$

**Definition:**  $\mathbf{M}$  is finitary if  $\mathbf{M}(i)$  is finitary  $\mathbb{k}$ -linear for all  $i$

**Definition:**  $\mathbf{M}$  is transitive if  $\mathbf{M}$  is finitary and for any indecomposable  $X, Y$  in  $\mathbf{M}$  there is a 1-morphism  $F$  such that  $X$  is isomorphic to a direct summand of  $F Y$

**Intuition:** Transitive action of a group (for us: a multisemigroup)

**Definition:**  $\mathbf{M}$  is simple transitive if  $\mathbf{M}$  is transitive and has no non-trivial  $\mathcal{C}$ -invariant ideals.

**Example:** Cell 2-representations are simple transitive.

# Classification of simple transitive 2-representations

**Theorem.** [M.-Miemietz]

Under some natural assumption, cell 2-representations are the only simple transitive 2-representation.

**Applies to:**

- ▶ Soergel bimodules in type  $A$ .
- ▶  $\mathcal{C}_A$ .
- ▶ Finitary quotients of 2-Kac-Moody algebras of finite type.

# Classification of simple transitive 2-representations

**Theorem.** [M.-Miemietz]

Under some natural assumption, cell 2-representations are the only simple transitive 2-representation.

Applies to:

- ▶ Soergel bimodules in type  $A$ .
- ▶  $\mathcal{C}_A$ .
- ▶ Finitary quotients of 2-Kac-Moody algebras of finite type.

# Classification of simple transitive 2-representations

**Theorem.** [M.-Miemietz]

Under some **natural** assumption, cell 2-representations are the only simple transitive 2-representation.

Applies to:

- ▶ Soergel bimodules in type  $A$ .
- ▶  $\mathcal{C}_A$ .
- ▶ Finitary quotients of 2-Kac-Moody algebras of finite type.

# Classification of simple transitive 2-representations

**Theorem.** [M.-Miemietz]

Under some **natural** assumption, cell 2-representations are the only simple transitive 2-representation.

**Applies to:**

- ▶ Soergel bimodules in type  $A$ .
- ▶  $\mathcal{C}_A$ .
- ▶ Finitary quotients of 2-Kac-Moody algebras of finite type.

# Classification of simple transitive 2-representations

**Theorem.** [M.-Miemietz]

Under some **natural** assumption, cell 2-representations are the only simple transitive 2-representation.

**Applies to:**

- ▶ Soergel bimodules in **type A**.
- ▶  $\mathcal{C}_A$ .
- ▶ Finitary quotients of 2-Kac-Moody algebras of finite type.

# Classification of simple transitive 2-representations

**Theorem.** [M.-Miemietz]

Under some **natural** assumption, cell 2-representations are the only simple transitive 2-representation.

**Applies to:**

- ▶ Soergel bimodules in **type A**.
- ▶  $\mathcal{C}_A$ .
- ▶ Finitary quotients of 2-Kac-Moody algebras of finite type.

# Classification of simple transitive 2-representations

**Theorem.** [M.-Miemietz]

Under some **natural** assumption, cell 2-representations are the only simple transitive 2-representation.

**Applies to:**

- ▶ Soergel bimodules in **type A**.
- ▶  $\mathcal{C}_A$ .
- ▶ **Finitary quotients** of 2-Kac-Moody algebras of finite type.

# Classification of simple transitive 2-representations

**Theorem.** [M.-Miemietz]

Under some **natural** assumption, cell 2-representations are the only simple transitive 2-representation.

**Applies to:**

- ▶ Soergel bimodules in **type A**.
- ▶  $\mathcal{C}_A$ .
- ▶ **Finitary quotients** of 2-Kac-Moody algebras of finite type.

# Other results and future challenges

**Known:** Morita theory for finitary 2-categories ([M.-Miemietz]).

**Known:** Classification of isotypic 2-representations of certain fiat 2-categories ([M.-Miemietz]).

**Known:** An analogue of Schur's lemma for cell 2-representations of certain fiat 2-categories ([M.-Miemietz]).

Classification of 2-representations for finitary 2-categories?

Classification of simple transitive 2-representations for finitary 2-categories?

Is the number of simple transitive 2-representations always finite?

Homological algebra for 2-representations?

# Other results and future challenges

**Known:** Morita theory for finitary 2-categories ([M.-Miemietz]).

**Known:** Classification of isotypic 2-representations of certain fiat 2-categories ([M.-Miemietz]).

**Known:** An analogue of Schur's lemma for cell 2-representations of certain fiat 2-categories ([M.-Miemietz]).

Classification of 2-representations for finitary 2-categories?

Classification of simple transitive 2-representations for finitary 2-categories?

Is the number of simple transitive 2-representations always finite?

Homological algebra for 2-representations?

# Other results and future challenges

**Known:** Morita theory for finitary 2-categories ([M.-Miemietz]).

**Known:** Classification of isotypic 2-representations of certain fiat 2-categories ([M.-Miemietz]).

**Known:** An analogue of Schur's lemma for cell 2-representations of certain fiat 2-categories ([M.-Miemietz]).

Classification of 2-representations for finitary 2-categories?

Classification of simple transitive 2-representations for finitary 2-categories?

Is the number of simple transitive 2-representations always finite?

Homological algebra for 2-representations?

# Other results and future challenges

**Known:** Morita theory for finitary 2-categories ([M.-Miemietz]).

**Known:** Classification of isotypic 2-representations of certain fiat 2-categories ([M.-Miemietz]).

**Known:** An analogue of Schur's lemma for cell 2-representations of certain fiat 2-categories ([M.-Miemietz]).

Classification of 2-representations for finitary 2-categories?

Classification of simple transitive 2-representations for finitary 2-categories?

Is the number of simple transitive 2-representations always finite?

Homological algebra for 2-representations?

# Other results and future challenges

**Known:** Morita theory for finitary 2-categories ([M.-Miemietz]).

**Known:** Classification of isotypic 2-representations of certain fiat 2-categories ([M.-Miemietz]).

**Known:** An analogue of Schur's lemma for cell 2-representations of certain fiat 2-categories ([M.-Miemietz]).

Classification of 2-representations for finitary 2-categories?

Classification of simple transitive 2-representations for finitary 2-categories?

Is the number of simple transitive 2-representations always finite?

Homological algebra for 2-representations?

# Other results and future challenges

**Known:** Morita theory for finitary 2-categories ([M.-Miemietz]).

**Known:** Classification of isotypic 2-representations of certain fiat 2-categories ([M.-Miemietz]).

**Known:** An analogue of Schur's lemma for cell 2-representations of certain fiat 2-categories ([M.-Miemietz]).

Classification of 2-representations for finitary 2-categories?

Classification of simple transitive 2-representations for finitary 2-categories?

Is the number of simple transitive 2-representations always finite?

Homological algebra for 2-representations?

# Other results and future challenges

**Known:** Morita theory for finitary 2-categories ([M.-Miemietz]).

**Known:** Classification of isotypic 2-representations of certain fiat 2-categories ([M.-Miemietz]).

**Known:** An analogue of Schur's lemma for cell 2-representations of certain fiat 2-categories ([M.-Miemietz]).

Classification of 2-representations for finitary 2-categories?

Classification of simple transitive 2-representations for finitary 2-categories?

Is the number of simple transitive 2-representations always finite?

Homological algebra for 2-representations?

# Other results and future challenges

**Known:** Morita theory for finitary 2-categories ([M.-Miemietz]).

**Known:** Classification of isotypic 2-representations of certain fiat 2-categories ([M.-Miemietz]).

**Known:** An analogue of Schur's lemma for cell 2-representations of certain fiat 2-categories ([M.-Miemietz]).

Classification of 2-representations for finitary 2-categories?

Classification of simple transitive 2-representations for finitary 2-categories?

Is the number of simple transitive 2-representations always finite?

Homological algebra for 2-representations?

# Other results and future challenges

**Known:** Morita theory for finitary 2-categories ([M.-Miemietz]).

**Known:** Classification of isotypic 2-representations of certain fiat 2-categories ([M.-Miemietz]).

**Known:** An analogue of Schur's lemma for cell 2-representations of certain fiat 2-categories ([M.-Miemietz]).

Classification of 2-representations for finitary 2-categories?

Classification of simple transitive 2-representations for finitary 2-categories?

Is the number of simple transitive 2-representations always finite?

Homological algebra for 2-representations?

THANK YOU!!!